

**Trends in Relative Abundance and  
Size of Selected Finfishes and  
Shellfishes Along the Texas Coast:  
November 1975-December 1995**


**by  
Billy Fuls  
and  
Lawrence W. McEachron**

**Management Data Series  
No. 137  
1997**



**COASTAL FISHERIES DIVISION**

4200 Smith School Road  
Austin, Texas 78744



TRENDS IN RELATIVE ABUNDANCE AND SIZE OF SELECTED FINFISHES AND  
SHELLFISHES ALONG THE TEXAS COAST: NOVEMBER 1975-DECEMBER 1995


by

Billy Fuls and Lawrence W. McEachron

MANAGEMENT DATA SERIES

NO. 137

1997



Texas Parks and Wildlife  
Coastal Fisheries Division  
4200 Smith School Road  
Austin, Texas 78744

## TABLE OF CONTENTS

	PAGE
LIST OF TABLES . . . . .	ii
LIST OF FIGURES . . . . .	iii
LIST OF APPENDICES . . . . .	v
ACKNOWLEDGMENTS . . . . .	vi
ABSTRACT . . . . .	vii
INTRODUCTION . . . . .	1
MATERIALS AND METHODS . . . . .	1
RESULTS . . . . .	4
Gill Net . . . . .	4
Bay Bag Seine . . . . .	5
Bay Trawl . . . . .	6
Gulf Trawl . . . . .	6
Oyster Dredge . . . . .	7
Beach Seine . . . . .	7
Beach Bag Seine . . . . .	7
Intracoastal Waterway Trawl . . . . .	8
Hydrologic Data . . . . .	8
SEAMAP . . . . .	8
Summer . . . . .	8
Fall . . . . .	8
OVERVIEW . . . . .	9
LITERATURE CITED . . . . .	18
TABLES . . . . .	19
FIGURES . . . . .	56
APPENDICES . . . . .	71

## LIST OF TABLES

	PAGE
Table 1. Mean catch rates (No./h) and mean total lengths (mm) of selected fishes and blue crab caught with gill nets (all meshes combined) by bay system during spring 1976-95 . . . . .	19
Table 2. Mean catch rates (No./h) and mean total lengths (mm) of selected fishes and blue crab caught with gill nets (all meshes combined) by bay system during fall 1975-95 . . . . .	24
Table 3. Annual mean catch rate (No./ha) and mean total lengths (mm) of selected fishes and shellfishes caught with 18.3-m bag seines by bay system during 1977-95. . . . .	29
Table 4. Annual mean catch rates (No./h) and mean total lengths (mm) of select fishes and shellfishes caught with 6.1-m trawls in Texas bay systems during 1982-95 . . . . .	37
Table 5. Annual mean catch rates (No./h) and mean total lengths (mm) of select fishes and shellfishes caught with 6.1-m trawls in the Texas Territorial Sea during 1985-95 . . . . .	42
Table 6. Annual mean catch rates (No./h) and mean total lengths (mm) by size class of Eastern oyster caught with 46.0-cm wide dredges on "reef" stations in Texas bay systems during 1984-95 . . . . .	47
Table 7. Seasonal (May-Nov) mean catch rates (No./ha) and mean total lengths (mm) of select fishes and shellfishes caught with 60.9-m beach seines in 5 Texas gulf shoreline areas during 1987-95 . . . . .	48
Table 8. Seasonal (May-Nov) mean catch rates (No./ha) and mean total lengths (mm) of select fishes and shellfishes caught with 18.3-m bag seines in 5 Texas gulf shoreline areas during 1987-95 . . . . .	51
Table 9. Annual mean catch rate (No./h) and mean total lengths (mm) of select fishes and shellfishes caught with 6.1-m trawls within the Intracoastal Waterway in Texas bay systems during 1992-95 . . . . .	54

## LIST OF FIGURES

	PAGE
Figure 1.	Texas gulf shoreline and Texas Territorial Sea (TTS) . . . . . 56
Figure 2.	Spring gill net mean catch rates (No./h $\pm$ 1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1976-95. . . . . 57
Figure 3.	Fall gill net mean catch rates (No./h $\pm$ 1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1975-95 . . . . . 58
Figure 4.	Spring gill net mean total lengths (mm $\pm$ 1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1976-95 . . . . . 59
Figure 5.	Fall gill net mean total lengths (mm $\pm$ 1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1975-95 . . . . . 60
Figure 6.	Seasonal bag seine mean catch rates (No./ha) for juvenile red drum (Nov-Mar), black drum (Jun-Jul), spotted seatrout (Jul-Nov) and Atlantic croaker (Feb-May) during 1978-95. Red drum 35-75 mm, spotted seatrout 20-75 mm, black drum 35-110 mm and Atlantic croaker 30-85 mm are considered to be young-of-the-year . . . . . 61
Figure 7.	Seasonal bag seine mean total lengths (mm $\pm$ 1SE) for juvenile red drum (Nov-Mar), black drum (Jun-Jul), spotted seatrout (Jul-Nov) and Atlantic croaker (Feb-May) during 1978-95. Red drum 35-75 mm, spotted seatrout 20-75 mm, black drum 35-110 mm and Atlantic croaker 30-85 mm are considered to be young-of-the-year . . . . . 62
Figure 8.	Seasonal bag seine mean catch rates (No./ha) for juvenile brown shrimp (Apr-Jul), white shrimp (Jul-Nov) and blue crab (Mar-Jun) during 1978-95. Brown and white shrimp 33-82 mm and blue crab 13-42 mm are considered to be young-of-the-year . . . . . 63
Figure 9.	Seasonal bag seine mean total lengths (mm $\pm$ 1SE) for brown shrimp (Apr-Jul), white shrimp (Jul-Nov) and blue crab (Mar-Jun) during 1978-95. Brown and white shrimp 33-82 mm and blue crab 13-42 mm are considered to be young-of-the-year . . . . . 64
Figure 10.	Annual bay trawl catch rates (No./h $\pm$ 1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95 . . . . . 65

Figure 11.	Annual bay trawl mean total lengths (mm $\pm$ 1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95 . . . . .	66
Figure 12.	Annual gulf trawl mean catch rates (No./h $\pm$ 1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95 . . . . .	67
Figure 13.	Annual gulf trawl mean total lengths (mm $\pm$ 1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95 . . . . .	68
Figure 14.	Annual mean catch rates (No./h) for Eastern oyster spat ( $\leq$ 25 mm), small oysters (26-75 mm) and market oysters ( $\geq$ 76 mm) during 1984-95 . . . . .	69
Figure 15.	Annual mean total lengths (mm) for small and market Eastern oysters during 1984-95 . . . . .	70

## LIST OF APPENDICES

	PAGE
Appendix A. Summary of historical sampling dates, gear description, procedures, dates, number of samples collected, weighting factors, and list of species collected. . .	71
Appendix B. Summary of hydrological data collected for gill net, bay and beach bag seine, oyster dredge, bay and gulf trawl, beach seine, and ICWW trawl samples . . . . .	93
Appendix C. Summary of SEAMAP samples by year and depth zone for brown shrimp, white shrimp, pink shrimp and blue crab collected off Texas during 1982-95 . . . . .	104

## ACKNOWLEDGMENTS

We thank each member of Texas Parks and Wildlife, Coastal Fisheries Division staff who conscientiously collected and recorded data. This study was conducted with partial funding from the U.S. Department of Interior, Fish and Wildlife Service under DJ 15.605 (Project F-34-M), previous projects under PL 88-309 and the U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service under PL 99-659 (Project 2-IJ).



## ABSTRACT

The objective of coastal monitoring projects is to determine the status of marine resources for management and harvest purposes. Trends in relative abundance and size of finfishes and shellfishes have been monitored since 1975 using a standardized fishery-independent sampling program in Texas bay systems. Data were collected with bag seines along bay and gulf shorelines, gill nets along bay shorelines, beach seines along gulf shorelines, and trawls in coastal bay and Texas Territorial Sea waters. Oyster dredges were used to sample bay oyster reefs.

Data comparisons were made between 1994 and 1995 for coastwide catch rates for all gears. Coastwide spring and fall gill net catch rates for red drum declined, but were higher than the average from 1975-1995. Spotted seatrout spring catch rates decreased slightly, but remained stable for the 5th straight year in fall. Spring black drum catch rate decreased slightly, but equaled the previous year in fall; spring and fall catch rates were well above average. Coastwide seasonal bay bag seine catch rates increased for red drum, spotted seatrout and black drum and decreased for Atlantic croaker, brown shrimp, white shrimp and blue crab. Coastwide annual bay trawl catch rates equaled the previous year for brown shrimp, increased for pink shrimp and decreased for white shrimp, blue crab and Atlantic croaker. Coastwide annual Gulf trawl catch rates increased for brown shrimp and white shrimp, and decreased for blue crab and Atlantic croaker. Coastwide annual catch rate for Eastern oyster spat and small oysters increased in 1995, while market oysters declined slightly from the record high in 1994. Data collected during 1995 were used to make resource and harvest management decisions.

## INTRODUCTION

Fishery independent monitoring data are used to determine relative abundance and size of finfishes and shellfishes in Texas coastal waters to regulate and allocate harvest in Texas jurisdictional waters. To collect these data, Texas Parks and Wildlife (TPW) has used various gears systematically in Texas estuaries and the Gulf of Mexico since 1975 (Appendix A, Tables A.1-5). TPW initiated a standardized fishery independent monitoring program in 1975 using gill nets, in 1977 using bag seines, in 1982 using trawls in bays, in 1984 using oyster dredges on bay oyster reefs, in 1985 using trawls in the gulf, and in 1987 using beach seines in the Gulf to monitor and assess relative trends in abundance and size of finfishes and shellfishes. Monitoring of blue crab populations in Texas Bays was initiated in 1977 (Hammerschmidt 1982). Gill nets set during spring (9 April-23 June) and fall (9 September-23 November), and monthly bag seine, trawl, oyster dredge, and beach seine samples provide a statistically consistent and cost efficient method for obtaining population trend information on juvenile, sub-adult, and adult finfish and shellfish.

The objectives of the present study were to:

1. monitor and determine trends in species composition, size and relative abundance of selected finfishes and shellfishes in Texas coastal bay systems and in the gulf off Texas.
2. publish the results in a report which will assist resource managers to effectively manage finfishes and shellfishes.

Differences in the information in this report compared to previous reports are due to updating the data base. The present report should be considered the most accurate to date.

## MATERIALS AND METHODS

Bag seines, trawls and monofilament gill nets (Appendix A) were used in each of the 9 Texas bay systems: Sabine Lake, Galveston, East Matagorda, Matagorda, San Antonio, Aransas, Corpus Christi, upper Laguna Madre and lower Laguna Madre. Trawls, identical to those used in the bays, were used in five gulf areas of the Texas Territorial Seal (TTS)  $\leq 16.7$  km from shore: 24.1 km either side of each of the Sabine Pass jetties (Sabine), Galveston jetties (Galveston), Matagorda jetties (Port O'Connor), Aransas Pass jetties (Port Aransas), and 48.2 km north from the Texas-Mexico border (Port Isabel) (Figure 1). Oyster dredges (Appendix A) were used in the Galveston, Matagorda, San Antonio and Aransas bay systems. Bag seines, identical to those used in Texas bays, and beach seines (Appendix A) were used along gulf beach shorelines in five areas: Sabine Pass-Bolivar Peninsula, Galveston Island-Follets Island-Surfside Beach, Matagorda Peninsula, Matagorda Island and Mustang Island-South Padre Island (Figure 1).

Gill net, bag seine, and beach seine sites were randomly selected from grids (1 minute latitude by 1 minute longitude) that contained  $\geq 15.2$  m of shoreline. Each selected grid was subdivided into 144 5-second "gridlets". All "gridlets" that contained  $\geq 15.2$  m of shoreline were used to randomly choose sample sites.

Gill net sets were conducted overnight during each spring and fall season (Appendix A). The spring season began with the 2nd full week in April and extended for 10 full weeks. The fall season began with the 2nd full week in September and extended for 10 full weeks. Between three and five nets were set each week in each bay, except in East Matagorda Bay where only two overnight sets were made during each week. On no more than six nights during each season could three nets be set in a bay system. Each sampling week extended from 1 h before sunset on Sunday through 4 h after sunrise the following Sunday. Gill nets were set perpendicular to shore with the smallest mesh shoreward. Nets were set within 1 h before sunset and were retrieved within 4 h after the following sunrise. Total fishing time was recorded (nearest 0.1 h).

One half of the monthly gulf and bay bag seine samples were collected during each of the 1st-15th and the 16th-31st of the month (Appendix A). Bay and gulf bag seines were pulled parallel to the shoreline for 15.2 m; gulf bag seines were pulled in the same direction as the longshore current. The surface area sampled (nearest 0.01 ha) was estimated using distance pulled and length of extension of the bag seine. No grid was sampled more than once in a month.

One half of the monthly beach seine samples were collected during each of the 1st-15th and 16th-31st of the month (Appendix A). Beach seines were pulled parallel to gulf shorelines in the same direction as the longshore current for 30.5 m. The surface area sampled (nearest 0.01 ha) was estimated using distance pulled and length of extension of the beach seine. No grid was duplicated in a month.

Trawls were used in bays which were stratified into three zones: Zone 1 (upper bay nearest mouths of rivers), Zone 2 (lower bay farthest from rivers), and Zone 5 [Intracoastal Waterway (ICWW)]. Trawl sites in Zones 1 and 2 were randomly selected from bay grids (1-minute latitude by 1-minute longitude) that contained water  $\geq 1$  m deep in at least 1/3 of the grid and which were known to be free of obstructions. One half of the monthly trawl samples in each zone in each bay system were collected during each of the 1st-15th and 16th-31st of the month (Appendix A). In East Matagorda Bay all water was designated as Zone 1; in each of Sabine Lake, upper and lower Laguna Madre all water was designated as Zone 2. In Zones 1 and 2, trawls were towed in a circular motion near the center of each grid. Trawl sites for Zone 5 were randomly selected from all grids containing the ICWW. Each randomly selected grid was divided into 144 5-second "gridlets"; the center-most gridlet which contained the center of the ICWW within that grid was used as a starting point for the sample. Trawls in Zone 5 were pulled linearly in the channel either toward the nearest gulf pass or away from it; this direction was alternated

with each sample. All trawl tows within bays were 10 minutes in duration. No grid was duplicated in a month.

Gulf trawl sites in each area were randomly selected from Gulf grids (1-minute latitude by 1 minute longitude) in the TTS (Figure 1) that contained water  $\geq 1.8$  m deep in at least 1/3 of the grid and which was known to be free of obstructions. One half of the samples in each area were collected during each of the 1st-15th and 16th-31st of the month (Appendix A). Trawls were towed linearly, parallel to the fathom curve; direction of tow (north or south) was randomly chosen for the initial tow and alternated on subsequent tows. All tows were 10 minutes in duration. No grid was duplicated in a month.

Trawls were used during daylight in the gulf off Sabine Pass, Galveston, Port O'Connor, Port Aransas, and Port Isabel during June and November 1993 in conjunction with the Southeast Area Monitoring and Assessment Program (SEAMAP). Detailed descriptions of the gear, sample stations, and sample procedures are reported by Stuntz et al. (1985).

In each major oyster producing bay (Galveston Bay, Matagorda Bay, San Antonio Bay and Aransas Bay) oyster reef areas were mapped; areas in which Eastern oysters form reefs which are  $\geq 0.2$  m higher than adjacent bottom for a continuous distance of  $\geq 1.4$  m long and 0.4 m wide. Oyster dredge sites were randomly selected from bay grids containing defined oyster reefs. Each selected grid was divided into 144 5-second "gridlets". All gridlets that contained defined oyster reefs were used to randomly choose sample sites. One half of the oyster samples were collected during each of the 1st-15th and 16th-31st of the month (Appendix A). Dredges were pulled linearly for 30 seconds. Stations were not duplicated within a month.

Sample catch rates for each species, or category of species, were calculated by dividing total number captured by either total hours fished (gill net, trawl, and oyster dredge) or ha sampled (bag seine and beach seine). Catch rates for each bay system were then calculated by month, year or season. Bay specific catch rates were weighted for coastwide estimates (Appendix A). Fish greater than 204 mm long were eliminated from bag seine catch rate calculations based on the findings of McEachron and Green (1986). Live Eastern oysters were grouped into spat (5-25 mm), small oysters (26-75 mm), and market oysters ( $\geq 76$  mm).

Lengths (total or standard) of animals caught were recorded. In gill nets, up to 19 individuals of each species, within each mesh size, were measured on each sampling day. In trawls, 50 shrimp (length from tip of rostrum to tip of telson) of each species (brown, white, pink), 35 blue crabs (carapace width between spine tips) and up to 19 individuals of all other species were measured in each sample. For all other gears, up to 19 specimens were measured for each species in each sample collected.

Mean total lengths of individual species in gill nets were calculated for each of the four mesh sizes. Mean lengths for the combined meshes were calculated by weighing individual species mean lengths in each mesh by the

number of each species caught in each mesh. For all other gears, mean lengths of individual species were calculated from individuals measured in each sample. Coastwide total mean lengths for each species in all gears were weighted according to the catch rate in each bay system, and by bay specific and gear specific weighting factors used for coastwide catch rates.

Surface salinity (ppt), water temperature (°C) and turbidity [Nephelometric Units (NTU)] were measured at the set and pickup for each gill net and prior to each bag seine and beach seine sample (Appendix B). Bottom salinity, water temperature, and turbidity were measured prior to each trawl and oyster dredge sample (Appendix B).

## RESULTS

### Gill Net

Coastwide spring red drum (Sciaenops ocellatus) catch rate declined slightly in 1995 (0.86/h) (Table 1; Figure 2). Highest spring coastwide red drum catch rate (1.3/h) occurred in 1992, with lowest catch rates during 1977-79 (0.3/h). Average size decreased to 483 mm TL in 1995 (Table 1; Figure 4).

Coastwide fall red drum catch rate decreased in 1995 (0.7/h); highest fall coastwide catch rate for red drum (1.0/h) occurred in 1979 and 1993, with lowest catch rates (0.5/h) in 1982 and 1983 (Table 2; Figure 3). Average size decreased to 464 mm TL in 1995 (Table 2; Figure 5).

Coastwide spring spotted seatrout (Cynoscion nebulosus) catch rate decreased in 1995 (0.7/h); highest spring coastwide spotted seatrout catch rate (1.1/h) occurred in 1976, with lowest catch rates in 1979 and 1984 (0.3/h) (Table 1; Figure 2). Average size decreased to 455 mm TL in 1995 (Table 1; Figure 4).

Coastwide fall spotted seatrout catch rate in 1995 equaled 1994 (0.4/h); highest fall coastwide spotted seatrout catch rate (0.7/h) occurred in 1976, with lowest catch rate in 1979 (0.2/h) (Table 2; Figure 3). Average size increased to 440 mm TL in 1995 (Table 2; Figure 5).

Coastwide spring black drum (Pogonias cromis) catch rate decreased in 1995 (1.2/h); highest spring coastwide black drum catch rate (1.5/h) occurred in 1994; lowest occurred in 1978 (0.3/h) (Table 1; Figure 2). Average size decreased to 423 mm TL in 1995 (Table 1; Figure 4).

Coastwide fall black drum catch rate in 1995 equaled 1994 (1.4/h); highest fall coastwide black drum catch rate occurred in 1993 (1.6/h), with lowest in 1979 and 1984 (0.3/h) (Table 2; Figure 3). Average size increased to 438 mm TL (Table 2; Figure 5).

Spring and fall coastwide southern flounder (Paralichthys lethostigma) and sheepshead (Archosargus probatocephalus) catch rates varied little over

the past 10 years (<0.1-0.1/h), but are generally down from pre-1983 years (0.1-0.3/h) (Tables 1 and 2).

Coastwide spring Atlantic croaker (Micropogonias undulatus) catch rates have remained at  $\leq 0.1/h$  since 1978 (Table 1, Figure 2). The 1994 fall catch increased in 1995 (0.3/h). Highest fall catch rate on record (0.5/h) was in 1993; lowest catch rate occurred in 1975 (0.1/h) (Table 2; Figure 3).

Coastwide 1995 catch rates for all fish combined remained equal to 1994 (7.5/h) in spring, but increased during fall to 6.2/h (Tables 1 and 2).

Spring and fall coastwide catch rates of blue crab (Callinectes sapidus) have remained at  $\leq 0.1/h$  over the past 7-8 years, but are generally down from pre-1987 years (0.1-0.2/h) (Table 1 and 2). Average size increased in 1995 to 153 mm in spring and decreased to 142 mm in fall.

#### Bay Bag Seine

Annual (calendar year) bag seine catch rates for select species are listed in Table 3. Seasonal trends in catch rates and mean lengths are presented for the following select species:

Coastwide red drum catch rates increased in 1995; they were highest during November 1990-March 1991 and lowest during November 1989-March 1990 (Figure 6). Mean coastwide lengths fluctuated between 46 and 58 mm TL (Figure 7).

Coastwide spotted seatrout catch rates increased in 1995; they were highest during July through November 1991 and lowest during 1984-86 (Figure 6). Mean coastwide lengths fluctuated between 44 and 56 mm TL (Figure 7).

Coastwide black drum catch rates increased in 1995; they were highest in 1979 and 1990 (Figure 6). Mean coastwide lengths fluctuated between 54 and 93 mm TL (Figure 7).

Coastwide Atlantic croaker catch rates decreased in 1995; they were highest in 1982 and lowest in 1989 (Figure 6). Mean coastwide lengths fluctuated between 49 and 57 mm TL (Figure 7).

Coastwide brown shrimp (Penaeus aztecus) catch rates decreased in 1995; they were highest in 1987 and lowest in 1979 (Figure 8). Mean coastwide lengths fluctuated between 54 and 64 mm TL (Figure 9).

Coastwide white shrimp (P. setiferus) catch rates decreased in 1995; they were highest in 1982 and lowest in 1985 (Figure 8). Coastwide mean lengths fluctuated between 52 and 59 mm TL (Figure 9).

Coastwide blue crab catch rates decreased in 1995; they were highest in 1985 and lowest in 1989 (Figure 8). Coastwide mean lengths fluctuated between 23 and 28 mm TL (Figure 9).

Annual catch rates of other species caught in bag seines varied by species and bay (Table 3).

#### Bay Trawl

Annual coastwide bay trawl catch rates for all finfish combined increased in 1995 (273/h); they ranged from 134/h in 1984 to 318/h in 1991 (Table 4).

Coastwide brown shrimp catch rates in 1995 (41/h) equaled those in 1994; they ranged from 21/h in 1983 to 49/h in 1989 (Table 4; Figure 10). Coastwide mean length decreased in 1995 to 81 mm TL, and in previous years ranged from 83-97 mm (Table 4; Figure 11).

Coastwide white shrimp catch rates decreased in 1995 (26/h); they ranged from 20/h in 1988 and 1989 to 46/h in 1982 (Table 4; Figure 10). Mean coastwide length increased in 1995 to 92 mm TL, and in previous year ranged from 84-101 mm (Table 4; Figure 11).

Coastwide annual blue crab bay trawl catch rates reached a record low of 10/h in 1995; they have ranged up to 24/h in 1992 and 1994 (Table 4; Figure 10). Coastwide mean length decreased to a record low of 64 mm TL in 1995 (Table 4; Figure 11).

Coastwide Atlantic croaker catch rates decreased in 1995 (64/h); they ranged from 27/h in 1985 to 112/h in 1992 (Table 4; Figure 10). Coastwide mean length remained the same in 1995, but has generally declined from 1983-92 (Table 4; Figure 11).

Coastwide pink shrimp (*P. duorarum*) catch rates increased in 1995 (5/h); they were highest in 1991 and 1995 and lowest in 1982, 1984 and 1993 (Table 4).

Annual catch rates of other major species caught in bay trawls varied by species and bay (Table 4).

#### Gulf Trawl

Annual coastwide Gulf trawl catch rates for all finfish combined increased in 1995 (380/h), and have ranged from 174/h in 1985 to 406/h in 1992 (Table 5).

Coastwide brown shrimp catch rates increased in 1995 (28/h); they ranged from 9/h in 1986 and 1994 to 59/h in 1989 (Table 5; Figure 12). Coastwide mean length increased in 1995, and has ranged from 97 (1992) to 109 (1985) mm TL (Table 5; Figure 13).

Coastwide annual white shrimp catch rates increased in 1995 (12/h); they ranged from 10/h in 1990 and 1994 to 24/h in 1985 and 1986 (Table 5; Figure 12). Mean coastwide length increased in 1995, and has ranged from 105 (1986 and 1992) to 115 (1985 and 1990) mm TL (Table 5; Figure 13).

Coastwide blue crab catch rates decreased in 1995 (2/h); they ranged from 1/h in 1987-89 to 6/h in 1991 (Table 5; Figure 12). Coastwide mean length decreased in 1995 to 70 mm (carapace width), but in previous years had decreased from 127 mm in 1985 to 69 mm in 1992 (Table 5; Figure 13).

Coastwide Atlantic croaker catch rates increased from 23/h in 1985 to 162/h in 1993 then decreased to 50/h in 1995 (Table 5; Figure 12). Mean coastwide length decreased in 1995 to 119 mm TL, but in previous years had decreased from 142 mm in 1985 to 113 mm in 1993 (Table 5; Figure 13).

Coastwide annual pink shrimp catch rates decreased to 3/h in 1995; they ranged from 1-4/h in past years (Table 5).

Annual catch rates of other major species caught in Gulf trawls varied by species and Gulf area (Table 5).

#### Oyster Dredge

Coastwide catch rates of Eastern oyster (Crassostrea virginica) spat increased in 1995 (1,194/h); they ranged from 491/h in 1984 to 1,880/h in 1989 (Table 6; Figure 14).

Coastwide catch rates of small Eastern oysters increased in 1995 (2,354/h); they ranged from 1,001/h in 1986 to 2,615/h in 1991 (Table 6; Figure 14). Mean coastwide length increased in 1995 and previously ranged from 46-54 mm TL (Table 6; Figure 15).

Coastwide catch rates of market Eastern oysters decreased in 1995 (690/h); they ranged from 215/h in 1990 to 708/h in 1994 (Table 6; Figure 14). Coastwide mean length in 1995 was equal to 1994 (91 mm TL), which equaled the highest on record (Table 6; Figure 15).

#### Beach Seine

Coastwide annual catch rates and mean lengths of select finfish and shellfish species varied among species, gulf areas and years (Table 7). Striped mullet (Mugil cephalus) generally had highest catch rates.

#### Beach Bag Seine

Coastwide annual catch rates and mean lengths of select finfish and shellfish species varied among species, Gulf areas and years (Table 8). Generally, striped mullet, blue crab and white shrimp had highest catch rates;



however, Gulf menhaden (Brevoortia patronus) and Spanish mackerel (Scomberomorus maculatus) were also among the most frequently caught species in 1995.

#### Intracoastal Waterway Trawl

Coastwide annual catch rates and mean length of individual select finfish and shellfish species varied among species, bays and years (Table 9). Coastwide brown shrimp and blue crab catch rates were highest during 1992 (117/h and 77/hr, respectfully), while white shrimp catch rates were highest during 1993 (35/h). Pink shrimp coastwide catch rate was highest in 1995 (22/h). Atlantic croaker had the highest coastwide catch rate (241/h) in 1992; the San Antonio Bay system had the highest total finfish catch rate (1,670/h) in 1992.

#### Hydrologic Data

Hydrologic data varied among years, among bay systems and among gulf areas (Appendix B). Coastwide annual salinity increased during 1995 in coastal bays and in Gulf waters (Appendix B; Tables B.1, B.4, B.7, B.10, B.13 and B.16). Bay salinities were generally higher in upper Laguna Madre than in any other bay. Gulf salinities were highest off Port Isabel. Water temperatures followed seasonal trends. Coastwide annual bay and gulf bottom water temperatures were similar to 1994 values (Appendix B; Tables B.2, B.5, B.8, B.11, B.4 and B.17).

#### SEAMAP

##### Summer (June)

Catch rates of brown shrimp by depth zone ranged from 829/h in 19-37 m to 43/h in 74-91 m during 1995 (Appendix C, Table C.1). Historically, brown shrimp are predominately caught in water 19-37 m deep.

White shrimp were caught primarily in water  $\leq 18$  m deep during all years (Appendix C, Table C.1). At these depths, catch rates have ranged from 4/h-41/h in all years.

Pink shrimp were captured in waters  $\leq 55$  m deep (0-195/h) during all years (Appendix C, Table C.1). They were caught predominately in waters  $\leq 37$  m deep.

Blue crab are caught primarily in the 0-18 m zone (Appendix C, Table C.1). Catch rates at these depths ranged from 3-20/h in all years.

Fall (November)

Brown shrimp were caught in all depth zones, with highest catch rates at water depths generally >18 m (Appendix C, Table C.2). White shrimp, pink shrimp and blue crab were predominately caught in waters  $\leq$ 37 m deep.

#### OVERVIEW

TPW is mandated by the Texas Legislature and the TPW Commission to annually investigate population trends, habitat variability, socio-economics, commercial and recreational fishing impacts and any other factors or conditions which may result in increases or decreases of finfishes and shellfishes in Texas waters. Long-term trend data based on independent standardized monitoring programs are necessary to assess changes in relative abundance of these populations. Shrimp data were used to recommend dates for the annual closure of Texas gulf waters to shrimping. Oyster data were used to establish the oyster transplant season in Galveston Bay. Finfish data were used to recommend changes in fishing regulations. These data were used to develop management plans for shrimp, oysters, and blue crabs as mandated by the Texas Legislature. Additionally, these data are used routinely by "outside" scientists in the private and public sector, especially the Gulf of Mexico Fisheries Management Council and the Gulf States Marine Fisheries Commission. TPW data base was used extensively by both the Galveston Bay, and Corpus Christi Bay National Estuary Programs to determine status and trends of populations. Data in the present report are used to determine long-term trends in abundance and stability of finfishes and shellfish populations in Texas coastal waters and to implement management regulations.

Effective management of marine species populations requires knowledge of the relationship between spawning and subsequent adult abundance (Cushing 1970, Gulland 1977). Since it has been possible to detect changes in annual abundances with bag seines and gill nets, it may be possible to determine stock-recruitment relationships utilizing these gears. To determine these relationships, it is imperative that the standardized monitoring program used by TPW be maintained.

To determine effects of natural or man induced events in Texas coastal ecosystems, standardized monitoring programs used by TPW should be maintained. The following "meta events" affecting coastal waters were documented in 1995. Other unreported events may have occurred.

1. The Texas Closure (15 May-15 July; 61 days) for shrimping in state offshore Gulf waters ( $\leq$ 9 nautical miles) was coordinated with NMFS for a combined closure out to 200 nautical miles offshore (Exclusive Economic Zone) to increase yield and value for the shrimping industry.

2. The following TPW hatchery-reared fish were stocked into Texas bays: 192.3 million red drum fry, 24.2 million red drum fingerlings, 32.3 million spotted seatrout fry, 1.8 million spotted seatrout fingerlings.
3. During 1995, the TPW Artificial Reef Program placed six donated structures in the Gulf on the Freeport reef site in the shape of a star, placed seven structures on the existing Port Mansfield site, and enhanced three existing sites elsewhere.

The Artificial Reef Program received a donation of one 4-pile jacket structure from Union Pacific Resources. The full structure, the North Padre Island A-58, was located 54 nautical miles from Port Aransas in 254-ft of water. The jacket structure was donated using a new environmentally sensitive "partial mechanical removal method" developed through a cooperative agreement between the U.S. Minerals Management Service and the TPW. The jacket was cut by divers at 86-ft from the surface. No explosives were used to remove the upper jacket and maximum biological profile of the structure was left standing.

4. Airplane flights on the opening days of the spring (15 May-15 July) and fall (15 August-15 December) bay shrimping seasons revealed the Texas bay commercial shrimping fleet was active from Galveston to Corpus Christi Bay. On 15 May, 255 shrimp vessels were in Galveston Bay, 200 in Matagorda Bay, 141 in San Antonio Bay, 127 in Aransas Bay, and 110 in Corpus Christi Bay. On 15 August, 275 were in Galveston Bay, 395 in Matagorda Bay, 91 in San Antonio Bay, 89 in Aransas Bay, and 36 in Corpus Christi Bay.
5. There were 348 sea turtle strandings along the coast during 1995. Most were Kemp's Ridley (41%) and loggerhead (36%) sea turtles. Many of these strandings occurred in conjunction with Gulf nearshore shrimp trawling effort.
6. During August through November, extremely high tides and above normal salinities were observed along the coast. This resulted in numerous nearshore Gulf species being reported in bay waters. Gray snapper was especially common during this period in middle and lower Texas coastal bays. Numerous juvenile tarpon were caught in coastal bays by cast netters during September and October.
7. Aquaculture facilities continue to flourish along the coast. However, during late spring and early summer an outbreak of *Taura* virus caused massive die-offs (70-90%) in private shrimp aquaculture ponds containing *Pennaeus vannamei*. Pond water was quarantined and not released until September. Laboratory studies determined that except for post-larval white shrimp, native species were not susceptible to the virus. Severe financial losses were the result of the outbreak.
8. Controversy over three new aquaculture shrimp farms on the mid-Texas coast generated much public comment concerning degradation of water quality in receiving waters adjacent to the proposed farms. As a result,

management agencies, politicians, industry, and the public are working on finding solutions to the opposition to siting of new shrimp farms.

9. The Southern Star Shrimp Farm on the Arroyo Colorado was raising American eels and found an infection of the nematode Anquillicola crassus. This parasite has not been previously documented in the United States. This nematode was first described in Japanese eels, and later found in European eels. The eels in the aquaculture facility were marketed, and the ponds chemically treated to kill any cysts from the nematode. No evidence of this nematode was found in subsequent samples of wild eels.
10. During summer, the Gulf of Mexico "Loop Current hypoxic dead zone" off Louisiana and upper Texas was estimated at 7,000 square miles. Although about equal in size to the 1993 and 1994 events, the hypoxic zone was about double the average area documented during years prior to 1993.
11. In March, the Salt Bayou Water Control Structure was completed within the Sabine Lake ecosystem complex. This structure will assist managers in restoring a more natural hydrological regime within the Salt Bayou/Keith Lake marsh complex.
12. Tropical Storm Dean caused heavy rains and 4-ft above normal high tides in the Sabine Lake System during August.
13. During spring gill net sampling, three grass carp were caught off Atkinson Island in upper Galveston Bay near the Houston Ship Channel. Salinities ranged 3-4 ppt. Possible Spartina alterniflora was observed in the gut, but positive identification was impossible due to maceration of the gut contents.
14. Several cases of *Vibrio* were reported in individuals from the middle coast. Subsequent newspaper and television reports caused the public to become increasingly concerned about the possibility of infection during swimming or fishing. TPW and Texas Department of Health helped allay fears of the virus.
15. Maintenance dredging of Cedar Bayou, a natural pass to the Gulf between Matagorda and San Jose Islands, began in July and was completed in September. The Bayou was dredged to a depth of 2.13 m and a width of 18.3 m. After dredging, the channel opening to the Gulf migrated frequently.
16. During January, TPW samples indicated small oysters in Aransas Bay were in record abundance. Favorable salinities and temperatures in 1994 were responsible for the high abundance.
17. Annual rainfall in the Coastal Bend totaled 36.92 inches, 6 inches above normal. Corpus Christi experienced the third-wettest day (7.92 inches) ever recorded (29 October). Only 22 September 1915 (7.94 inches) and 27 June 1931 (8 inches) exceeded the 29 October daily total.

18. Brown tide persisted in the Laguna Madre (upper and lower) for the sixth consecutive year. No open bay mortalities were observed with these blooms, but fingerling production at the TPW fish hatchery at Flour Bluff was adversely affected; low dissolved oxygen was recorded in ponds. It was reported that clearer water was seen in some areas of Baffin Bay during May and June.
19. Commercial black drum landings increased to record highs in upper Laguna Madre. About 1.3 million pounds were reported landed by trotline fishermen. One dealer recorded more than 100,000 lbs being landed in September from Baffin Bay alone; all fish were shipped to San Antonio and Houston.
20. The exotic bivalve mussel Perna perna, which has been reported from mid-Texas coast areas, was discovered in the Brownsville Ship Channel and on channel markers extending about 8 miles to the north in lower Laguna Madre. This nonindigenous mussel was first documented in 1992 on the Brazos-Santiago jetties.
21. The Texas Department of Health, Seafood Safety Division published a Fish Advisories & Bans report which listed the southern portion of the lower Laguna Madre closed to the harvest of shellfish during 26 September to 27 October due to unsafe levels of red tide (Gymnodinium breve). Densities ranged from 1-601 cells/ml.
22. Kill Investigations:

In January, about 88,000 spider crab (Libinia sp.) exoskeletons washed ashore from Sabine Pass to San Luis Pass. University of Texas Marine Science Institute (Port Aransas) experts determined the cause to be molting. Similar occurrences in Long Island Sound were the result of mating pods undergoing synchronous molting. It is theorized, synchronous molting caused the Texas event.

In April, about 1,200 blue catfish (some > 1 meter in TL), thousands of Atlantic croaker, and a few red drum were killed at Gulf States Utilities (Entergy) discharge canal adjacent to Bessie Heights marsh in Sabine Lake. No cause was determined.

In July, seismic activity in St. Charles Bay killed 8,551 fish consisting of several species. Restitution of \$25,138 was made to TPW.

A large number of red drum (>30 inches TL) were found in July within an area of upper Laguna Madre called the "Grave Yard". Investigation revealed that the fish were most likely caught during illegal commercial fishing activities and discarded to avoid detection.

During August, extensive fish kills (millions of fishes) of mostly hardhead catfish and Atlantic croaker occurred near Sabine Pass. Gulf nearshore discolored water was analyzed by Dr. Bill Wardle (Texas A & M,

Galveston). He found Prorocentrum minima, a dinoflagellate species that has been reported to produce ichthyotoxins.

On 25 August, about 50 million Gulf menhaden were killed in the lower Colorado River at the junction between the river and the Gulf of Mexico. The kill was caused by low dissolved oxygen brought about by high water temperature, low wind, and little tidal movement.

In mid-September, a large fish kill occurred in the Arroyo Colorado between Port Harlingen and Rio Hondo. About two million Gulf menhaden died. Subsequent water sampling revealed no dissolved oxygen below a water depth of three feet.

On 23 September, an estimated 16 million Gulf menhaden died along 3.5 miles of shoreline in the lower Laguna Madre at Port Isabel. The kill coincided with the passage of an unusually strong cold front. Water samples revealed the red tide organism (G. breve) at levels of 601 cells/ml or less.

Following the passage of Hurricane Opal in October, a fish kill occurred in a dead-end bar-ditch between Sea Rim State Park's Marsh Unit and Fence Lake within the Sabine Lake area. Millions (about 90 mm long) of Gulf menhaden, hundreds of blue catfish, some flounder, and some large white shrimp (9-10 count) were killed. Low dissolved oxygen was the suspected cause of death.

Investigation of a hardhead catfish kill began off Sabine Pass on 4 October and proceeded through 30 November. Concentrations of dead and dying hardhead catfish extended throughout the Galveston Bay and Brazos River areas to Matagorda Bay and Pass Cavallo. About one million animals were estimated to have died. Bull red drum and black drum were also affected during the kill, and were found along the Texas City Dike and along Gulf beaches in the Galveston area. During the assessment, many fish were observed swimming erratically, displaying physical abnormalities and possessing large external parasite loads. Gross examination of a freshly killed red drum revealed the following anomalies: eyes deeply recessed in the eye cavities, eroded anus, external infestation by parasitic isopods, argulid parasites, and caligid copepods. Whether the external parasites were responsible for the death of the red drum was not determined. Samples of hardhead catfish were sent to Texas A & M and labs in Florida for analysis. A kill of a similar magnitude occurred off Florida and was caused by an amoebic infestation of the kidney and liver.

### 23. Oil/chemical spills

During February, about 1,680 gallons of crude oil spilled from a broken pipeline covering 200 acres of interior marsh on Smith Point in Galveston Bay. Blue crabs, plovers, clapper rails, and scaups were found oiled in the spill area.

On 5 February, the Norwegian tanker *Berge Banker* spilled 900 barrels of crude oil in the nearshore Gulf about 60 miles east of Freeport after colliding with another vessel. By 21 February, oil had reached Matagorda Island and was being cleaned up by oil spill response contractors. Fewer than 200 birds were found to be "oiled." No fish were observed killed.

In April, a cumene spill occurred at Ingleside when a barge and tanker collided. Little or no effects on living resources (wetlands & rookeries) were noted following the spill.

#### 24. Regulations and Legislation:

##### Adopted 23 March:

State-wide hunting and fishing regulations. (TPW Commission adopted the repeal of 31 TAC 65.1 to 65.91 and a new 31 TAC 65.1 to 65.91 regarding the statewide hunting and fishing proclamation with amendments to the proposals in Exhibit A, Agenda item No. 1):

Snook: Changed daily bag limits from 3 to 1, possession limits from 6 to 2, minimum size limits from 20 inches to 24 inches. Additional conservation measures were needed to help maximize the snook resource and opportunities for harvest. Spawning success of snook will be increased with a larger minimum size by allowing more snook to reach sexual maturity before becoming vulnerable to harvest. In addition, a lower bag limit will more equitably distribute those snook available to be caught. These regulation changes should encourage more catch and release on this limited stock.

Finfish in Commercial Bait Shrimp Trawls: Established an exemption for persons shrimping under a commercial bait shrimp license during the period 1 July through 31 August from the provision requiring the non-game finfish portion of the trawl catch not to exceed 50% by weight of the total trawl catch, except for those species regulated by bag and size limits. This rule allows up to 1200 live non-game finfish not regulated by bag and size limits to be retained. This rule change will provide additional live fish as bait for sport anglers during that summer period when bait shrimp are less abundant. In addition, the bycatch of bait shrimp trawls will be reduced because fewer trawls will be needed to retain the same amount of bait fish. During other periods of the year, when bait shrimp are more abundant, live bait fish can be obtained in amounts up to 50% of the weight of the total trawl catch.

Fee changes regarding red drum licenses: The proposed fee charged for an Exempt Red Drum Tag was \$6 and a Duplicate Exempt Red Drum Tag was \$6, and authorized license deputies to retain \$1 as a fee for collecting each tag fee and issuing each tag. It also established that the holder of an Exempt Red Drum Tag is prohibited from possessing both an Exempt Red Drum Tag and Bonus Red Drum Tag. Holders of an Exempt Red Drum Tag were added to the Special

Regulation regarding possession of red drum over the stated maximum size limit.

Adopted 1 June:

Shrimp management proclamation (agenda item 4). Regulation changes repealed the 1 ½-inch mesh size scheduled to be implemented 1 September 1995 for commercial bait shrimp trawls, commercial bay shrimp trawls, and commercial Gulf trawls used to catch seabobs because limited entry plan was in place. (TPW adopted amendment to 31 TAC 57.661 concerning trawl mesh sizes for selected portions of the shrimp fishery).

Adopted 6 July:

Regulations concerning the use of sand pumps for taking aquatic organisms. The sand pump was made a legal device for the taking of ghost shrimp (Callichris islagrande) for bait, provided the gear was operated manually only, that there was a conservative bag limit on the take of ghost shrimp, and that it would be used for personal use only. (TPW Commission adopted 31 TAC 65.3 and 65.78).

Red Snapper regulations (Agenda item 4). Regulation changes for red snapper caught in state waters will correspond to those regulations implemented by the Gulf of Mexico Fishery Management Council on 1 January 1995, for red snapper caught in federal waters where most of the red snapper fishery occurs. Establishing rules compatible with those in place in federal waters will reduce angler confusion, enhance law enforcement, and aid in the recovery of this overfished stock. (TPW Commission adopts amendments to 31 TAC 65.72).



Adopted 2 November:

New fees effective FY 1996:

Resident Combination Hunting and Fishing	\$32.00
"Super" Resident Combination Hunting and Fishing (includes all 7 state stamps)	\$49.00
"All Purpose" Resident Combination Hunting and Fishing (includes all 7 state stamps, annual public hunting permit, Texas Conservation Passport)	\$100.00
Resident Fishing	\$19.00
Special Resident Fishing (blind residents, qualified resident disabled veterans, resident commercial fishermen)	\$6.00
Special Resident Fishing (resident seniors 65 after 9-1-95)	\$9.00
Temporary (14-day) Resident Fishing	\$12.00
Temporary (3-day) Resident Fishing	\$8.00
Lifetime Resident Combination Hunting and Fishing (includes exemption from all state hunting and fishing stamps)	1,000.00
Lifetime Resident Fishing (includes exemption from all state fishing stamps)	\$600.00

Regulation 57.801 concerning EEZ. Rule will function by allowing better and more timely management of species which migrate between state and federal waters off Texas and improve law enforcement while reducing confusion for Gulf anglers. Per implementation of S.B. 733 by the 74th Legislature, the item allows the Commission to authorize the executive director to modify state coastal fisheries regulations when necessary to make them consistent with federal regulations in the EEZ (specified waters off the Texas coast).

Senate Bill 733:

Texas Parks and Wildlife implemented Proclamation 57.081 which establishes authority for the Executive Director to take action as necessary to modify state coastal fisheries regulations to conform with federal regulations in the Exclusive Economic Zone. This authority has been used once to date for closure of the commercial red snapper fishery.

Senate Bill 750:

The Shrimp License Management program for Commercial Bay and Bait Shrimp Boat Licenses was established for license year beginning 1

September 1995. Initial eligible license participants for the Shrimp License Management Program for commercial bay and bait shrimp boat licenses were identified. In conjunction with the initial license year and beginning license period, as designated under Senate Bill 750, a Shrimp License Management Review Board was established through an election procedure; individual appeal cases concerning eligibility into the license management program were reviewed. The Voluntary Buy-back fund was established and the additional fee increases associated with licenses that allow handling of shrimp in some manner were instituted for license year beginning 1 September 1995. All monies associated with these fee increases were placed in the established fund.

The Shrimp License Management Review Board was created within the required deadline and met routinely to deal with special cases concerning license eligibility and to consult with the department on other issues surrounding the license management system. The voluntary Buy-Back Program was developed.

House Bill 1956:

Removed requirements to produce several reports to the Legislature and the Governor. These special reports are not being produced, but production of similar reports containing like information is ongoing.

## LITERATURE CITED

- Cushing, D. H. 1970. Fisheries biology: a study in population dynamics. University of Wisconsin Press. Madison, Wisconsin.
- Gulland, J. A. 1977. The management of marine fishes. University of Washington Press. Seattle, Washington.
- Hammerschmidt, P. C. 1982. Population trends and commercial harvest of the blue crab (Callinectes sapidus, Rathbun) in Texas bays, September 1974-August 1979. Management Data Series Number 38. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- Matlock, G. C., and M. F. Osborn (Ferguson). 1982. Shallow-water surface areas and shoreline distances on the Texas coast. Management Data Series Number 37. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- McEachron, L. W., and A. W. Green. 1985. Trends in relative abundance and size of selected finfishes along the Texas coast: November 1975-June 1984. Management Data Series Number 79. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- McEachron, L. W., and A. W. Green. 1986. Assessment of annual relative abundance and mean length of six marine fishes in Texas coastal waters. Proceedings of the 38th Southeastern Association of Fish and Wildlife Agencies. 38:506-519.
- Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. N. Lea, and W. B. Scott. 1991. Common and scientific names of fishes from the United States and Canada. American Fisheries Society Special Publication 20. Bethesda, Maryland.
- Stuntz, W. E., C. E. Bryan, K. Savastano, R. S. Waller, and P. A. Thompson. 1985. SEAMAP environmental and biological atlas of the Gulf of Mexico, 1982. Gulf States Marine Fisheries Commission. Ocean Springs, Mississippi.

Table 1. Mean catch rates (No./h) and mean total lengths (mm) of selected fishes and blue crab caught with gill nets (all meshes combined) by bay system during spring 1976-95. Blank indicates no measurement taken; ND = no data.

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aranzas	Christi	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
Red drum	1976	ND	310	ND	429	1.0	410	1.0	451	0.6	412	0.1	509	1.2	458	0.7	435				
	1977	ND	450	0.2	418	0.1	467	0.3	380	0.4	409	0.1	438	0.5	442	0.3	426				
	1978	ND	394	0.4	429	0.5	485	0.2	400	0.4	443	0.2	495	0.5	462	0.3	460				
	1979	ND	480	0.1	466	0.2	421	0.4	414	0.2	423	0.3	479	0.3	452	0.3	448				
	1980	ND	449	0.4	451	1.1	387	0.7	400	0.4	373	1.0	430	0.6	438	0.8	418				
	1981	ND	431	0.2	465	0.2	408	0.6	396	0.4	399	0.3	424	1.0	438	0.4	420				
	1982	ND	474	0.4	436	0.5	425	0.4	408	0.4	430	0.5	469	1.0	497	0.6	464				
	1983	ND	474	1.0	475	0.6	411	0.7	402	0.5	385	0.4	427	0.8	479	0.6	444				
	1984	ND	482	0.7	446	0.1	430	0.2	513	0.3	439	0.8	457	0.7	514	0.5	473				
	1985	ND	538	0.5	514	0.2	457	0.2	465	0.4	463	0.6	457	0.3	505	0.6	508	0.4	500		
	1986	0.4	520	1.4	454	0.8	463	0.6	454	0.6	395	0.7	463	1.0	493	0.8	474				
	1987	0.2	516	0.6	501	0.9	465	0.7	451	0.6	459	0.6	463	1.1	508	0.7	483				
	1988	0.3	498	0.7	473	0.7	434	1.1	470	0.5	436	0.6	495	1.2	499	0.8	481				
	1989	0.5	480	0.7	478	1.7	492	0.6	452	0.7	438	0.5	469	0.9	517	0.7	476				
	1990	0.5	509	0.8	568	0.4	483	0.3	474	0.5	494	1.0	505	0.8	534	0.5	515				
	1991	0.5	581	0.5	532	0.3	495	0.3	447	0.4	472	0.9	476	1.2	509	0.5	504				
	1992	0.7	470	1.2	465	1.3	387	1.3	429	1.6	402	1.2	481	1.5	494	1.3	450				
	1993	0.4	529	1.2	529	2.6	514	0.9	426	1.6	439	1.2	509	1.3	511	1.1	490				
	1994	0.4	507	0.5	536	1.6	470	1.3	458	1.2	471	0.6	529	1.8	549	0.9	511				
1995	0.5	456	0.7	447	0.9	459	1.4	447	0.8	445	0.6	488	0.9	542	0.9	483					
Spotted seatrout	1976	ND	530	ND	422	0.3	422	0.5	382	3.3	465	0.4	405	3.4	457	1.1	453				
	1977	ND	516	2.0	434	0.2	381	0.9	392	1.0	422	0.4	372	1.2	442	0.8	422				
	1978	ND	523	0.4	441	0.6	409	1.4	408	0.1	435	0.5	437	0.9	474	1.4	503	0.7	456		
	1979	ND	515	0.4	426	0.3	490	0.1	436	0.4	507	0.3	524	0.4	442	0.6	525	0.3	495		
	1980	ND	419	0.8	402	0.6	426	0.9	402	0.2	465	0.3	506	0.5	473	0.9	497	0.5	449		
	1981	ND	483	1.8	416	0.4	406	0.7	453	0.8	468	0.5	445	0.4	423	2.2	471	0.8	456		
	1982	ND	491	0.9	454	0.5	489	0.8	440	0.7	435	0.8	489	0.8	481	2.5	485	0.9	472		
	1983	ND	510	1.7	441	0.7	452	0.8	444	0.6	447	0.7	478	0.7	509	1.3	500	0.7	476		
	1984	ND	498	0.7	468	0.3	439	0.3	483	0.2	435	0.2	473	0.7	475	0.3	472				
	1985	ND	506	0.6	467	0.3	424	0.3	457	0.4	430	0.4	471	1.4	485	0.5	473				
	1986	0.3	460	1.0	432	0.5	441	0.4	426	0.4	430	1.0	447	1.5	488	0.7	456				
	1987	0.2	339	0.6	449	0.7	436	0.4	434	0.4	447	0.5	456	0.9	478	0.4	490	1.9	508	0.7	474
	1988	0.2	386	0.7	459	0.8	430	0.5	435	0.5	458	0.8	478	0.4	507	1.6	498	0.7	470		
	1989	0.2	441	0.6	481	0.5	428	0.6	459	0.6	463	0.7	487	0.4	514	1.1	485	0.6	474		
	1990	0.1	441	0.5	457	0.6	432	0.6	480	0.5	442	1.1	447	0.2	468	1.3	455	0.6	456		
	1991	0.1	467	0.5	498	0.4	431	0.8	440	1.0	467	1.0	460	0.6	447	1.9	461	0.8	455		
	1992	0.2	406	0.7	446	0.4	440	0.4	449	0.7	443	1.3	463	0.6	529	1.9	483	0.8	467		
	1993	0.3	415	0.5	460	0.5	501	0.6	428	0.7	477	0.6	456	1.1	440	0.5	507	1.9	459	0.8	459
	1994	0.3	408	0.7	460	0.8	496	0.7	418	0.8	438	0.9	447	1.0	454	1.8	483	0.9	458		
1995	0.1	462	0.7	455	0.5	489	0.4	431	0.9	446	0.6	448	0.9	471	1.0	467	0.7	455			
Black drum	1976	ND	290	ND	418	0.8	418	1.0	306	0.9	389	0.6	360	0.5	387	0.7	366				
	1977	ND	388	0.3	262	0.5	519	1.0	314	1.2	316	0.5	347	0.4	377	0.9	428	0.7	374		
	1978	ND	439	0.4	345	0.2	300	0.1	306	0.4	325	0.4	325	0.1	398	0.8	395	0.3	373		
	1979	ND	292	0.7	328	0.5	415	<1	370	0.3	323	0.1	375	0.3	371	0.9	413	0.4	371		
	1980	ND	314	1.0	272	0.9	355	0.5	263	1.0	320	0.3	352	0.7	384	0.4	452	0.6	341		

Table 1. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake	Galveston	Matacorda	Matacorda	San Antonio	Aransas	Christi	Upper Laguna Madre	Lower Laguna Madre	Coastwide	No./h	Length	No./h	Length	No./h	Length	No./h	Length		
Black drum (Cont'd.)	1981	ND	0.8	418	0.8	312	0.3	301	0.4	352	0.8	362	0.1	379	1.1	390	0.9	391	0.7	381	
	1982	ND	0.6	343	0.8	294	0.5	363	0.7	317	1.1	300	0.4	339	0.7	374	1.2	400	0.8	347	
	1983	ND	0.9	337	2.7	365	0.6	355	0.6	323	0.2	340	0.9	371	1.0	400	1.6	441	1.0	372	
	1984	ND	0.6	373	1.0	391	0.2	414	0.2	460	0.1	559	0.5	414	0.6	442	0.6	459	0.4	417	
	1985	ND	0.5	346	0.4	313	0.2	476	0.1	426	0.2	396	0.2	342	0.8	361	0.4	372	0.4	374	
	1986	0.3	0.5	383	0.6	345	0.3	402	0.1	313	0.4	316	0.6	369	0.7	418	0.4	464	0.4	387	
	1987	0.1	0.5	368	0.6	320	0.4	366	0.2	392	0.5	352	0.5	459	1.1	453	0.5	458	0.7	409	
	1988	0.1	410	0.4	380	0.7	376	0.4	390	0.4	339	0.4	375	0.8	444	0.7	397	0.3	451	0.5	396
	1989	0.2	326	0.6	350	1.8	378	0.4	412	0.3	363	0.6	371	0.4	406	1.0	426	0.5	408	0.6	386
	1990	0.2	378	0.5	372	1.5	393	0.8	341	0.3	330	0.7	336	0.6	411	1.4	418	0.7	410	0.7	381
	1991	0.3	318	0.6	356	1.4	347	0.8	354	0.5	294	1.1	308	0.4	361	3.0	366	1.2	369	1.0	350
	1992	0.2	366	0.5	370	1.3	391	0.4	339	0.8	388	0.7	335	1.6	374	3.0	349	2.3	408	1.2	372
	1993	0.3	360	0.4	377	0.4	345	0.4	374	1.8	449	1.6	417	3.4	400	2.4	376	2.2	397	1.4	401
	1994	0.4	376	0.4	415	0.1	363	0.6	418	1.2	489	1.0	469	2.1	431	4.3	398	3.2	429	1.5	427
	1995	0.2	330	0.4	381	0.3	332	0.9	418	0.7	400	1.0	444	1.0	454	4.2	420	1.8	442	1.2	423
	Sheeps-head	1976	ND	0.0	ND	ND	0.1	420	0.1	420	0.3	341	0.6	342	0.0	367	0.3	318	0.2	345	
		1977	ND	<.1	338	<.1	234	0.1	280	0.2	308	<.1	232	0.1	294	0.1	380	0.1	336	0.1	311
1978		ND	0.0	0.4	296	0.4	254	<.1	278	0.1	313	0.2	354	0.2	356	0.2	394	0.2	358	0.1	350
1979		ND	<.1	305	0.1	297	<.1	391	<.1	402	0.1	320	0.5	362	0.1	370	0.2	340	0.1	350	
1980		ND	<.1	353	0.3	347	0.1	324	0.1	320	0.2	352	0.2	322	0.2	369	0.3	343	0.2	345	
1981		ND	<.1	393	0.2	326	<.1	453	0.6	335	0.3	349	0.1	319	0.2	390	0.6	325	0.2	342	
1982		ND	0.1	332	0.0	330	0.1	330	0.2	354	<.1	326	0.2	333	0.2	361	0.6	326	0.2	336	
1983		ND	0.1	313	0.4	311	0.1	373	0.2	392	0.1	349	0.3	370	0.2	392	0.4	342	0.2	354	
1984		ND	0.1	351	0.3	354	<.1	387	0.2	398	<.1	401	0.1	379	0.1	385	0.2	348	0.1	370	
1985		ND	<.1	352	0.2	372	<.1	337	<.1	409	<.1	382	0.1	424	<.1	427	0.1	353	0.1	382	
1986		<.1	372	0.2	356	<.1	369	<.1	369	0.1	417	<.1	305	0.1	388	<.1	427	0.1	370	<.1	382
1987		<.1	364	<.1	361	0.2	314	<.1	340	<.1	447	<.1	342	<.1	342	<.1	403	0.2	372	<.1	366
1988		0.0	<.1	405	0.1	350	<.1	357	<.1	342	0.1	348	0.1	348	0.1	407	0.1	369	<.1	366	
1989		<.0	529	0.1	384	0.3	324	<.1	371	<.1	379	<.1	350	0.2	412	0.2	371	0.2	371	0.1	374
1990		<.1	364	<.1	378	0.3	364	<.1	400	<.1	444	<.1	372	0.2	388	<.1	358	0.1	396	0.1	387
1991		<.1	354	<.1	381	0.2	343	<.1	359	<.1	491	<.1	304	<.1	367	<.1	406	0.1	389	<.1	382
1992		<.1	278	<.1	346	0.1	356	0.1	367	0.1	415	<.1	348	0.1	436	0.1	434	0.2	379	0.1	390
1993	<.1	343	<.1	376	0.2	360	0.1	408	0.1	355	<.1	408	0.1	422	<.1	427	0.1	394	0.1	392	
1994	<.1	353	<.1	374	0.2	413	<.1	372	<.1	338	<.1	374	<.1	435	<.1	429	0.1	375	<.1	376	
1995	<.1	309	<.1	389	0.1	428	0.1	407	0.3	359	0.1	363	0.3	446	<.1	426	0.1	372	0.1	393	
Southern flounder	1976	ND	0.0	ND	ND	0.0	328	0.0	328	0.1	335	0.0	350	0.0	350	0.2	350	0.2	345		
	1977	ND	<.1	351	0.1	358	<.1	358	<.1	208	0.1	358	<.1	430	0.0	345	<.1	345	<.1	347	
	1978	ND	<.1	249	0.1	352	<.1	330	0.1	279	<.1	338	0.1	338	<.1	345	0.1	344	<.1	323	
	1979	ND	<.1	451	0.1	348	<.1	290	0.1	388	<.1	291	0.1	373	<.1	320	0.2	366	0.1	354	
	1980	ND	0.1	344	0.1	325	0.1	307	<.1	316	<.1	292	0.1	316	<.1	364	0.1	366	0.1	330	
	1981	ND	<.1	244	<.1	340	<.1	270	<.1	291	<.1	368	0.1	332	0.1	348	0.1	338	<.1	322	
	1982	ND	0.1	303	<.1	319	<.1	307	<.1	305	0.1	299	0.1	361	0.1	337	0.1	350	0.1	332	
	1983	ND	0.1	366	0.1	318	0.1	327	<.1	333	<.1	329	0.1	385	0.1	359	0.1	371	0.1	357	
1984	ND	0.1	338	0.1	388	<.1	317	<.1	321	<.1	310	0.1	377	<.1	344	<.1	355	<.1	342		
1985	ND	0.1	349	0.1	348	<.1	346	0.1	349	<.1	347	0.1	353	0.1	346	0.1	336	0.1	354		
1986	<.1	345	<.1	345	0.2	329	<.1	358	<.1	316	<.1	357	<.1	395	0.1	354	0.1	370	0.1	344	
1987	<.1	364	<.1	339	0.1	330	<.1	304	0.1	345	<.1	336	<.1	333	0.1	407	<.1	401	<.1	353	

Table 1. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aranas	Christi	Upper Laguna Madre	Lower Laguna Madre	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Southern flounder (Cont'd.)	1988	<.1	292	0.1	349	<.1	354	<.1	350	<.1	334	<.1	353	<.1	400	<.1	360	<.1	359	<.1	
	1989	<.1	288	<.1	347	<.1	318	<.1	317	<.1	340	<.1	381	<.1	402	<.1	392	<.1	349	<.1	
	1990	<.1	309	<.1	351	<.1	354	<.1	350	<.1	311	<.1	347	<.1	333	<.1	410	<.1	358	<.1	
	1991	<.1	329	0.1	322	<.1	348	<.1	348	<.1	326	0.1	343	0.1	363	0.1	358	0.1	346	0.1	
	1992	<.1	319	0.1	366	<.1	346	<.1	373	<.1	355	<.1	377	<.1	438	<.1	394	<.1	374	<.1	
	1993	<.1	364	<.1	360	0.1	395	<.1	357	<.1	417	0.1	398	<.1	453	<.1	349	<.1	374	<.1	
	1994	<.1	334	<.1	343	0.1	378	<.1	362	<.1	332	<.1	362	<.1	332	<.1	380	<.1	352	<.1	
	1995	<.1	315	<.1	331	0.1	400	<.1	369	<.1	370	<.1	398	<.1	362	<.1	379	<.1	367	<.1	
	Atlantic croaker	1976	ND	298	0.2	298	0.1	277	0.2	332	0.0	277	1.0	277	0.0	277	0.8	333	0.3	306	0.3
		1977	ND	268	0.1	255	0.0	255	<.1	227	<.1	285	1.0	264	0.4	297	0.2	269	0.2	271	0.2
1978		ND	247	<.1	270	<.1	293	<.1	250	<.1	248	0.1	281	0.2	281	0.1	276	0.1	268	0.1	
1979		ND	260	<.1	257	<.1	263	0.0	260	0.0	0.0	0.1	265	0.1	298	0.2	308	0.1	279	0.1	
1980		ND	268	0.1	250	0.0	276	<.1	254	<.1	240	0.1	272	0.2	312	0.1	286	0.1	286	0.1	
1981		ND	264	0.1	250	0.1	270	0.0	265	<.1	289	0.1	266	0.1	302	0.1	277	0.1	282	0.1	
1982		ND	268	0.1	258	<.1	270	<.1	265	<.1	285	0.1	295	0.2	313	0.4	347	0.1	308	0.1	
1983		ND	268	0.1	278	<.1	273	<.1	277	<.1	286	0.2	285	0.2	289	0.4	314	0.1	286	0.1	
1984		ND	265	<.1	322	<.1	225	<.1	298	<.1	260	<.1	262	<.1	304	<.1	285	<.1	266	<.1	
1985		ND	273	<.1	318	<.1	260	<.1	184	<.1	115	0.1	265	0.1	267	0.1	261	0.1	266	0.1	
1986		0.1	259	0.4	271	0.1	250	<.1	245	<.1	292	0.3	255	0.2	297	0.1	288	0.1	272	0.1	
1987		<.1	263	0.2	242	<.1	236	<.1	268	<.1	246	<.1	282	<.1	319	<.1	251	0.1	263	0.1	
1988		0.1	259	0.1	226	<.1	226	<.1	268	<.1	260	0.1	261	0.1	337	<.1	296	<.1	276	0.1	
1989		0.1	268	0.1	280	<.1	250	0.0	268	0.0	262	<.1	284	<.1	342	<.1	283	<.1	274	<.1	
1990		<.1	278	0.1	264	<.1	268	<.1	283	<.1	276	<.1	267	<.1	245	<.1	272	<.1	269	<.1	
1991	0.1	297	0.1	256	<.1	237	<.1	239	<.1	252	0.1	261	<.1	269	0.1	267	<.1	263	<.1		
1992	0.1	263	0.2	270	<.1	257	0.0	277	0.0	232	0.1	204	<.1	290	0.1	266	0.1	261	0.1		
1993	0.1	286	0.2	256	<.1	237	<.1	277	<.1	267	0.1	265	<.1	295	0.1	272	0.1	264	0.1		
1994	0.1	297	0.1	272	<.1	266	<.1	263	<.1	257	0.1	279	<.1	328	<.1	298	0.1	279	0.1		
1995	<.1	278	0.2	281	<.1	288	<.1	276	<.1	273	<.1	300	<.1	382	<.1	321	<.1	275	<.1		
Gafftop-sail catfish	1976	ND	504	6.4	504	0.5	494	2.3	456	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.8	496	1.8	
	1977	ND	480	0.2	506	0.9	556	3.3	538	3.1	506	0.0	0.0	0.0	0.0	0.0	0.0	1.0	524	1.0	
	1978	ND	539	0.3	546	1.1	546	1.8	496	0.1	545	<.1	436	0.0	0.0	0.0	0.0	0.5	521	0.5	
	1979	ND	520	0.5	534	0.4	553	0.4	534	0.5	544	0.2	551	0.0	0.0	0.0	0.0	0.3	539	0.3	
	1980	ND	511	0.2	566	0.5	554	1.2	547	0.4	552	0.1	598	0.0	0.0	0.0	0.0	0.3	546	0.3	
	1981	ND	514	0.3	480	0.8	541	0.5	537	1.4	541	0.1	521	0.0	577	0.0	0.0	0.4	536	0.4	
	1982	ND	513	0.2	496	0.4	544	1.4	540	0.9	542	0.3	530	<.1	534	<.1	372	0.5	535	0.5	
	1983	ND	544	<.1	475	0.3	537	2.0	530	0.9	537	0.1	536	<.1	575	0.0	0.0	0.5	534	0.5	
	1984	ND	527	<.1	580	1.0	529	1.1	530	0.6	550	0.2	532	<.1	472	<.1	211	0.4	533	0.4	
	1985	ND	532	<.1	467	0.4	517	0.8	537	0.1	557	0.1	507	<.1	413	<.1	388	0.2	530	0.2	
	1986	0.2	490	0.4	468	0.3	533	0.5	554	0.4	529	0.4	534	0.0	374	0.0	0.0	0.3	528	0.3	
	1987	<.1	509	0.4	552	0.1	507	0.2	565	0.1	567	0.2	550	<.1	532	<.1	518	0.2	551	0.2	
	1988	0.1	538	0.2	511	0.1	530	0.3	563	0.3	562	0.2	550	0.0	0.0	<.1	428	0.2	537	0.2	
	1989	<.1	494	0.3	536	0.1	535	0.6	530	0.4	569	0.1	533	<.1	536	<.1	536	0.2	539	0.2	
	1990	<.1	518	0.8	528	0.2	460	0.8	534	0.6	555	0.4	554	0.0	0.0	0.0	0.0	0.4	537	0.4	
1991	<.1	520	0.2	504	0.2	528	0.5	531	0.7	527	0.4	530	<.1	546	0.0	0.0	0.4	537	0.4		
1992	<.1	519	0.1	521	0.2	556	0.3	556	0.6	578	0.1	530	0.0	0.0	<.1	508	0.2	542	0.2		
1993	<.1	457	0.5	494	0.2	581	0.5	543	0.8	563	0.3	503	0.0	0.0	<.1	405	0.3	535	0.3		
1994	<.1	518	0.1	495	0.2	569	0.8	545	1.2	571	0.2	548	0.0	0.0	0.0	0.0	0.3	554	0.3		
1995	<.1	508	0.4	498	0.3	543	0.3	516	0.7	557	0.2	529	0.1	0.0	<.1	210	0.2	533	0.2		

Table 1. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake	Galveston	Mataforda	Mataforda	San Antonio	Aransas	Christi	Upper Laguna Madre	Lower Laguna Madre	Coastwide	Upper Laguna Madre	Lower Laguna Madre	Coastwide	Upper Laguna Madre	Lower Laguna Madre	Coastwide	Upper Laguna Madre	Lower Laguna Madre	Coastwide	
		No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	
Gulf menhaden	1976	ND	261	ND	0.1	250	0.1	275	0.0	247	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	261
	1977	ND	251	0.7	299	0.1	245	0.1	233	0.3	247	2.6	255	<.1	282	<.1	229	<.1	229	0.9	253
	1978	ND	242	<.1	194	0.2	245	1.2	258	0.0	241	0.2	263	1.2	264	0.0	246	0.0	246	0.4	256
	1979	ND	251	0.0	132	0.1	251	<.1	132	<.1	241	0.1	255	0.2	260	0.0	253	0.0	253	0.3	251
	1980	ND	193	0.0	193	<.1	252	0.1	287	<.1	271	<.1	257	0.6	269	<.1	253	0.0	253	0.1	265
	1981	ND	260	0.0	254	0.2	254	0.1	252	0.2	254	0.1	243	0.1	246	0.1	244	0.1	244	0.2	255
	1982	ND	254	0.0	248	<.1	248	0.3	252	0.1	249	<.1	248	0.4	268	<.1	303	<.1	303	0.2	257
	1983	ND	252	0.0	244	0.2	243	0.1	244	0.1	244	0.1	248	0.1	304	0.1	252	0.3	252	0.3	252
	1984	ND	254	0.0	251	0.1	251	0.2	279	0.2	246	0.1	257	<.1	284	<.1	265	0.2	265	0.2	256
	1985	ND	253	<.1	281	0.5	243	0.3	243	0.4	250	0.6	250	<.1	244	0.8	260	0.5	260	0.5	252
	1986	0.1	279	1.3	226	0.1	242	0.1	244	0.2	245	0.4	258	<.1	252	<.1	253	0.4	252	0.4	251
	1987	<.1	348	1.2	245	<.1	241	0.0	226	0.0	226	0.2	242	<.1	240	0.1	253	0.3	245	0.3	245
	1988	<.1	278	0.1	244	0.0	244	<.1	278	<.1	236	<.1	253	<.1	257	<.1	290	0.1	249	0.1	249
	1989	<.1	269	1.4	249	0.0	242	<.1	226	0.0	187	0.1	235	0.0	257	0.0	248	0.0	248	0.3	248
	1990	<.1	270	1.6	242	<.1	216	0.1	263	<.1	255	<.1	237	<.1	308	<.1	239	0.4	242	0.4	242
	1991	<.1	253	0.3	252	<.1	216	0.1	239	<.1	281	0.1	255	0.0	251	0.0	241	0.1	247	0.1	247
	1992	<.1	266	0.7	257	0.0	207	0.1	245	0.1	251	0.1	275	<.1	252	<.1	279	0.2	257	0.2	257
	1993	<.1	256	1.5	247	0.0	257	<.1	217	0.0	251	<.1	242	<.1	312	<.1	282	0.3	247	0.3	247
	1994	0.1	267	0.5	260	0.0	235	<.1	254	<.1	262	0.1	253	<.1	238	<.1	295	0.1	258	0.1	258
	1995	<.1	275	0.2	257	<.1	252	0.1	255	<.1	265	<.1	206	0.0	266	<.1	269	0.1	257	0.1	257
Striped mullet	1976	ND	385	ND	0.2	322	0.2	338	0.6	366	0.0	0.0	0.0	375	0.0	0.0	0.0	0.0	0.2	358	
	1977	ND	322	0.0	314	0.2	314	0.9	317	0.8	319	0.1	340	0.2	368	0.2	345	0.2	345	0.3	323
	1978	ND	338	0.0	327	0.4	336	0.2	334	0.2	327	0.2	366	<.1	327	0.1	354	0.2	354	0.2	338
	1979	ND	320	0.1	336	0.1	341	0.7	343	0.2	339	0.1	333	0.1	404	0.1	354	0.2	341	0.2	341
	1980	ND	343	<.1	338	0.4	335	0.2	328	0.1	337	0.1	320	0.2	379	0.2	356	0.2	356	0.2	343
	1981	ND	318	0.1	345	<.1	336	<.1	341	0.1	336	0.1	321	0.2	353	0.2	353	0.1	344	0.1	344
	1982	ND	344	0.2	295	0.2	326	0.2	330	0.2	333	0.2	344	0.2	359	0.3	361	0.2	341	0.2	341
	1983	ND	350	0.1	346	0.1	346	0.2	341	0.2	341	0.1	351	0.2	367	0.2	368	0.2	368	0.2	352
	1984	ND	344	0.2	340	0.3	328	0.2	337	0.4	337	0.1	336	0.6	352	0.5	347	0.3	347	0.3	342
	1985	ND	340	0.2	339	0.3	332	0.1	328	0.3	340	0.1	338	0.2	380	0.1	339	0.2	339	0.2	342
	1986	<.1	326	0.2	321	0.2	330	0.1	368	0.2	336	0.1	340	0.1	368	0.1	341	0.1	340	0.1	340
	1987	<.1	312	0.2	366	0.1	319	0.2	348	0.2	354	0.1	356	0.1	402	0.2	359	0.2	359	0.2	357
	1988	<.1	327	0.1	344	0.2	333	0.1	350	0.1	343	0.1	344	0.1	371	0.1	364	0.1	364	0.1	348
	1989	<.1	323	0.2	348	0.4	339	0.2	356	0.2	356	0.1	344	0.1	400	0.1	372	0.2	372	0.2	354
	1990	<.1	325	0.2	341	0.3	342	0.4	342	0.2	340	0.2	340	0.4	389	0.4	353	0.3	353	0.3	354
	1991	<.1	325	0.1	347	0.2	341	0.2	343	0.3	335	0.1	343	0.2	386	0.1	377	0.2	377	0.2	350
	1992	<.1	310	0.1	352	0.3	340	0.3	342	0.4	352	0.2	355	0.2	389	0.2	374	0.2	374	0.2	355
	1993	<.1	331	0.1	358	0.3	371	0.2	355	0.3	379	0.4	379	0.2	379	0.1	354	0.2	379	0.1	353
	1994	0.1	343	0.1	347	0.1	381	0.3	359	0.2	368	0.1	365	0.2	386	0.2	383	0.2	383	0.2	362
	1995	<.1	341	0.3	356	0.2	366	0.5	364	0.2	366	0.1	359	0.2	403	0.1	376	0.2	376	0.2	361
Total finfishes	1976	ND	429	ND	5.2	394	7.6	391	9.5	415	6.2	332	1.1	378	7.1	419	7.3	408	7.3	408	
	1977	ND	316	4.3	395	5.9	442	8.2	428	8.1	428	7.6	297	3.8	366	4.3	377	3.95	377	6.7	395
	1978	ND	357	2.4	359	4.8	437	7.7	409	2.0	406	3.4	343	4.6	365	5.0	406	4.6	406	4.6	390
	1979	ND	345	2.5	336	3.4	409	3.2	453	3.2	433	2.7	393	3.2	360	3.2	411	3.8	387	4.1	387
	1980	ND	380	4.2	347	5.4	428	5.2	422	3.1	405	2.8	387	3.9	368	3.5	419	4.3	400	4.3	400
	1981	ND	4.6	369	5.5	363	5.3	408	6.1	417	6.0	432	2.8	634	4.2	353	6.5	406	5.2	396	
	1982	ND	8.1	378	4.7	368	5.3	405	6.8	411	5.8	417	4.6	400	4.5	367	8.8	394	6.4	397	
	1983	ND	9.0	369	7.6	384	4.5	417	7.2	422	5.5	404	5.5	397	5.0	373	7.5	409	6.6	394	
	1984	ND	6.2	389	3.7	397	4.3	449	5.6	431	3.9	432	4.8	397	3.2	369	4.6	412	4.7	410	
	1985	ND	7.6	381	3.8	408	5.2	446	4.1	479	3.6	452	5.0	368	3.6	350	5.2	384	5.1	404	

Table 1. (Cont'd.)

Species	Year	Sabine Lake		Galveston		East		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
Total	1986	4.9	432	9.3	377	5.4	381	5.0	425	3.5	422	3.2	418	5.7	371	2.9	387	5.2	425	5.3	398
finfishes	1987	2.0	517	8.7	373	4.3	384	4.0	430	2.9	420	3.4	431	3.8	420	3.0	432	5.9	434	4.8	408
(Cont'd.)	1988	2.5	472	6.7	385	4.6	401	4.5	411	4.7	444	3.0	436	6.4	390	3.2	407	5.4	436	4.8	411
	1989	2.6	474	9.0	365	7.4	396	5.1	428	6.4	437	4.2	403	4.4	402	2.8	432	4.7	425	5.5	403
	1990	2.5	485	10.5	367	8.2	403	6.6	432	6.1	448	5.1	410	6.8	405	3.5	405	5.2	424	6.5	405
	1991	3.1	474	6.9	367	11.7	358	6.4	415	6.1	437	6.0	400	5.8	405	5.3	381	7.2	409	6.4	398
	1992	2.6	445	8.4	395	8.8	423	6.3	407	5.9	448	7.1	412	7.0	410	5.7	409	8.4	431	7.0	414
	1993	2.4	480	9.8	387	8.7	459	7.0	424	8.6	467	6.9	453	9.7	419	4.8	427	8.3	428	7.8	425
	1994	2.7	451	6.6	394	6.8	467	7.2	419	9.0	444	7.2	438	7.4	425	7.7	428	9.7	454	7.5	429
	1995	2.3	463	2.4	390	7.4	460	7.8	400	9.3	422	6.8	411	6.7	421	7.2	417	6.3	435	7.5	412
Blue crab	1983	ND		0.2	151	0.3	154	0.1	151	0.2	142	0.3	142	0.2	151	0.1	156	0.2	145	0.2	147
	1984	ND		0.2	150	0.4	135	0.1	143	0.2	137	0.2	142	0.3	147	0.3	145	0.2	142	0.2	144
	1985	ND		0.3	149	0.5	151	0.2	144	0.3	136	0.2	141	0.2	149	0.3	141	0.2	158	0.2	147
	1986	0.2	146	0.3	151	0.6	133	0.2	140	0.1	135	0.1	144	0.1	154	<1	147	0.1	148	0.2	145
	1987	0.3	152	0.3	139	0.3	138	0.1	138	0.2	140	0.1	155	0.1	151	<1	137	0.1	142	0.1	141
	1988	0.3	154	0.1	148	0.1	159	<1	135	<1	141	<1	150	0.1	145	<1	115	0.1	152	0.1	147
	1989	0.2	157	0.1	137	0.4	128	<1	136	<1	128	<1	131	<1	149	<1	72	<1	147	0.1	136
	1990	0.2	154	0.2	141	0.2	129	<1	138	0.2	135	0.1	135	0.2	140	<1	114	0.1	139	0.1	138
	1991	0.1	141	0.2	132	0.4	135	0.2	144	0.1	136	0.1	144	0.1	140	<1	105	0.1	152	0.1	138
	1992	0.1	151	0.2	153	0.1	135	<1	144	0.1	133	0.1	142	0.3	150	0.4	146	0.1	146	0.1	147
	1993	0.2	161	0.1	144	0.2	162	0.1	147	0.1	148	0.1	152	0.2	148	0.1	147	<1	136	0.1	149
	1994	0.1	155	<1	144	0.1	160	<1	143	0.1	139	<1	149	0.1	158	<1	102	<1	129	<1	140
	1995	0.1	164	<1	147	0.3	165	<1	151	<1	161	<1	167	<1	152	<1	141	<1	107	<1	153



Table 2. Mean catch rates (No./h) and mean total lengths (mm) of selected fishes and blue crab caught with gill nets (all meshes combined) by bay system during fall 1975-95. Blank indicates no measurement taken; ND = no data.

Species	Year	East																			
		Sabine Lake		Galveston		Matagorda		Matagorda		San Antonio		Arkansas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
Red drum	1975	0.8	382	1.1	403	ND	487	1.2	337	0.9	326	1.1	339	0.4	330	0.3	424	0.7	474	0.9	373
	1976	ND	ND	1.0	509	1.1	487	0.5	415	1.6	406	0.5	395	0.5	460	0.4	442	1.3	465	0.9	452
	1977	ND	ND	0.6	445	0.9	390	0.8	435	1.0	386	0.6	392	0.5	427	0.2	364	0.4	448	0.6	416
	1978	ND	ND	0.3	429	0.7	376	1.1	395	0.6	384	1.0	401	0.4	429	0.3	455	0.4	493	0.6	412
	1979	ND	ND	0.8	386	0.7	403	1.4	353	1.9	376	0.9	378	0.8	352	0.5	387	0.5	449	1.0	378
	1980	ND	ND	0.5	436	0.8	473	0.6	434	0.9	411	1.1	386	0.7	370	0.5	454	0.7	449	0.7	419
	1981	ND	ND	0.5	429	0.7	405	0.6	390	0.7	373	0.8	403	0.6	396	0.3	515	0.8	488	0.6	422
	1982	ND	ND	0.6	440	0.9	401	0.6	390	0.5	360	0.4	386	0.3	417	0.2	456	0.5	440	0.5	412
	1983	ND	ND	0.6	436	0.8	394	0.5	418	0.6	407	0.4	410	0.3	448	0.2	486	0.7	509	0.5	440
	1984	ND	ND	0.9	451	1.1	551	0.4	381	0.6	383	0.5	377	0.8	400	0.7	457	0.7	472	0.7	433
	1985	ND	ND	0.9	421	1.3	420	0.8	394	1.3	385	0.9	427	0.7	436	0.3	460	0.9	478	0.9	423
	1986	0.4	481	0.7	468	0.9	453	0.8	403	1.2	441	0.9	454	0.5	450	0.4	486	0.9	495	0.8	456
	1987	0.4	449	0.5	459	0.9	446	0.8	372	1.0	473	0.6	459	0.4	424	0.3	527	1.5	532	0.7	467
	1988	0.5	399	0.8	437	1.5	486	0.9	418	1.1	457	0.9	454	0.5	458	0.3	520	1.3	522	0.8	463
	1989	0.4	449	0.6	479	1.1	511	0.4	402	1.1	468	0.7	423	0.6	476	0.3	533	1.1	521	0.7	475
	1990	0.4	500	0.3	488	0.8	497	0.5	408	1.1	458	1.0	477	0.8	432	0.7	553	1.0	534	0.7	482
	1991	1.1	412	0.5	393	0.9	380	0.6	402	1.3	375	1.0	442	1.5	451	0.6	517	1.5	514	0.9	441
	1992	0.5	531	0.7	482	2.0	494	0.8	419	0.7	453	1.4	435	1.0	477	0.7	502	1.3	479	0.9	465
	1993	0.3	484	0.4	482	1.9	526	0.9	439	1.6	480	1.7	490	1.0	500	0.7	555	1.4	531	1.0	496
	1994	0.6	426	0.6	437	1.9	478	0.6	447	1.0	470	1.0	468	0.6	471	0.4	568	1.1	539	0.8	481
1995	0.6	454	0.5	474	2.1	470	0.5	412	0.7	421	0.9	458	0.6	467	0.3	521	0.9	506	0.7	464	
Spotted seatrout	1975	0.1	413	0.2	447	ND	451	0.6	419	1.0	389	0.6	474	0.4	479	0.2	455	0.8	413	0.5	428
	1976	ND	ND	0.3	463	0.9	451	0.4	437	0.7	427	0.2	448	0.6	387	0.2	455	2.4	431	0.7	433
	1977	ND	ND	0.3	501	0.3	461	0.4	455	0.5	387	0.1	485	0.3	483	0.6	412	0.8	464	0.4	449
	1978	ND	ND	0.3	544	0.3	400	0.8	406	0.5	387	0.2	417	0.1	413	0.4	431	0.5	437	0.4	432
	1979	ND	ND	0.2	449	0.1	395	0.6	418	0.2	439	0.1	476	0.2	413	0.1	434	0.4	472	0.2	438
	1980	ND	ND	0.4	476	0.2	418	0.3	406	0.3	435	0.2	446	0.3	465	0.2	434	0.5	490	0.3	458
	1981	ND	ND	0.3	483	0.8	419	0.4	437	0.3	428	0.2	442	0.4	437	0.2	469	0.7	486	0.4	457
	1982	ND	ND	0.3	456	0.4	468	0.4	430	0.4	428	0.2	446	0.2	458	0.4	435	0.5	453	0.3	445
	1983	ND	ND	0.3	464	0.5	420	0.3	438	0.5	425	0.2	459	0.3	435	0.3	459	0.6	476	0.4	452
	1984	ND	ND	0.4	465	0.3	459	0.2	430	0.2	420	0.1	453	0.2	467	0.1	400	0.4	458	0.3	453
	1985	ND	ND	0.3	470	0.3	418	0.4	439	0.2	430	0.2	438	0.2	432	0.2	443	0.6	475	0.3	453
	1986	0.2	395	0.4	438	0.4	444	0.5	419	0.4	432	0.3	442	0.4	464	0.4	432	0.2	443	0.4	446
	1987	0.1	410	0.2	459	0.5	425	0.6	425	0.3	422	0.3	452	0.5	461	0.2	456	0.7	461	0.4	446
	1988	0.1	420	0.5	444	0.7	432	0.3	439	0.4	438	0.3	430	0.4	442	0.2	428	0.9	479	0.4	449
	1989	0.1	430	0.3	441	0.4	447	0.2	435	0.4	457	0.3	446	0.4	475	0.1	464	0.6	460	0.3	453
	1990	<1	399	0.2	460	0.5	461	0.2	427	0.2	479	0.3	459	0.5	474	0.1	505	0.5	477	0.3	467
	1991	0.1	378	0.2	442	0.3	473	0.5	406	0.4	415	0.3	436	0.6	449	0.4	482	0.8	466	0.4	443
	1992	0.1	392	0.3	418	0.5	452	0.4	417	0.2	436	0.4	457	0.6	463	0.5	508	0.8	443	0.4	448
	1993	0.1	450	0.3	446	0.9	441	0.9	428	0.4	430	0.3	441	0.9	427	0.5	468	0.6	447	0.4	444
	1994	0.1	398	0.4	434	0.8	465	0.3	417	0.5	431	0.4	434	0.6	442	0.5	448	0.6	429	0.4	436
1995	0.1	397	0.4	439	1.0	474	0.3	438	0.5	431	0.4	429	0.4	454	0.4	451	0.5	432	0.4	440	
Black drum	1975	0.5	294	0.4	366	ND	344	0.9	326	0.5	315	0.8	290	0.4	358	1.2	422	1.0	454	0.7	367
	1976	ND	ND	0.3	337	0.7	305	0.9	344	1.2	325	0.6	376	0.3	366	1.0	503	2.4	419	0.9	388
	1977	ND	ND	0.4	384	0.5	371	0.5	338	0.7	336	0.4	341	0.3	365	0.8	406	2.2	410	0.7	383

Table 2. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Christi	Madre	Madre	Madre	Madre	Madre	Madre	Madre	Madre	Madre	Madre	Madre		
		No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h	No./h		
Black drum (Cont'd.)	1978	ND	0.4	383	1.0	346	0.5	383	0.3	306	0.5	311	0.1	383	0.8	425	0.4	377	0.5	372	
	1979	ND	0.2	398	0.1	410	0.2	404	0.4	361	0.3	380	0.4	308	0.4	391	0.5	423	0.3	387	
	1980	ND	0.8	391	0.9	341	0.7	306	1.2	298	0.9	340	0.5	370	0.6	365	1.0	400	0.8	352	
	1981	ND	0.3	408	0.4	343	0.4	357	1.5	315	0.5	341	0.4	357	0.5	390	0.8	384	0.5	369	
	1982	ND	0.6	355	2.4	346	0.6	352	1.0	296	1.1	337	0.6	369	0.9	388	1.9	387	1.0	356	
	1983	ND	0.2	381	1.0	361	0.6	375	0.6	328	0.6	345	0.7	406	0.5	422	0.9	418	0.6	381	
	1984	ND	0.5	405	0.7	348	0.2	386	0.3	269	0.2	329	0.2	376	0.4	438	0.5	442	0.3	389	
	1985	ND	0.8	379	0.6	363	0.4	357	0.3	295	0.4	325	0.4	363	0.9	329	0.5	435	0.5	379	
	1986	0.4	360	0.7	380	0.6	303	0.6	351	0.4	342	0.5	357	0.3	388	0.5	417	0.5	441	0.5	379
	1987	0.3	378	0.4	376	1.5	376	0.4	384	0.3	364	0.5	370	0.2	384	0.4	403	0.6	465	0.4	393
	1988	0.2	355	0.5	387	1.2	339	0.7	346	1.0	334	0.7	330	0.7	337	1.5	405	0.6	422	0.8	368
	1989	0.5	324	2.0	384	1.4	358	0.8	351	1.0	337	1.4	373	1.3	416	1.5	421	1.2	401	1.3	383
	1990	0.3	342	0.4	375	0.8	368	0.6	362	1.0	298	1.0	334	0.6	398	1.0	431	1.0	423	0.8	372
	1991	0.3	347	0.5	382	1.0	364	0.6	375	1.3	369	0.7	321	0.9	340	2.2	359	1.8	367	1.0	361
	1992	0.4	394	0.5	402	1.1	422	0.7	394	0.3	352	0.9	372	0.8	372	1.4	363	2.2	366	0.9	375
	1993	0.3	372	0.6	400	1.0	456	0.8	430	1.0	449	1.6	439	1.7	424	3.8	422	2.7	401	1.6	421
	1994	0.5	370	0.5	415	0.3	442	0.7	438	0.9	453	0.9	429	0.9	432	4.9	430	2.2	445	1.4	433
	1995	1.0	347	0.6	358	0.5	461	0.6	354	0.6	398	1.1	408	1.5	472	4.5	461	2.3	480	1.4	438
	Sheeps-head	1975	0.0	<.1	362	ND	ND	0.1	316	0.2	291	1.1	296	0.2	376	0.3	409	0.1	352	0.3	323
		1976	ND	<.1	331	0.2	308	0.2	273	0.4	329	1.0	255	0.1	328	0.2	360	0.4	341	0.3	297
1977		ND	<.1	342	0.3	316	0.1	314	0.2	321	0.5	267	0.2	335	0.2	406	0.3	356	0.2	323	
1978		ND	0.1	308	0.2	307	0.1	342	0.5	371	0.6	306	0.2	361	0.2	376	0.1	300	0.2	337	
1979		ND	<.1	335	0.2	352	0.1	312	0.4	362	0.8	318	0.2	339	0.1	395	0.2	349	0.2	338	
1980		ND	0.1	283	0.1	309	<.1	353	0.7	296	0.6	307	0.2	361	0.2	382	0.4	330	0.3	316	
1981		ND	<.1	321	0.1	277	0.2	322	0.3	335	0.2	322	0.1	343	0.1	382	0.3	332	0.2	327	
1982		ND	0.1	330	0.3	332	0.1	313	0.1	296	0.2	350	0.1	365	0.1	383	0.3	330	0.1	339	
1983		ND	<.1	342	0.5	345	0.1	338	0.2	302	0.1	355	0.1	361	0.2	395	0.3	340	0.2	346	
1984		ND	<.1	369	0.3	383	<.1	369	<.1	427	<.1	436	<.1	383	0.1	417	0.1	333	0.2	346	
1985		ND	<.1	380	0.2	379	<.1	374	<.1	362	<.1	326	<.1	352	<.1	435	0.1	369	<.1	369	
1986		<.1	340	<.1	359	0.1	297	0.1	336	0.1	329	0.1	304	0.1	359	<.1	407	0.1	351	0.1	336
1987		<.1	402	<.1	381	0.1	366	0.1	352	0.1	371	0.2	360	0.1	340	<.1	386	0.2	342	0.1	355
1988		0.0	<.1	368	0.1	340	<.1	358	0.1	346	0.1	304	<.1	354	<.1	398	<.1	382	0.1	359	
1989		<.1	299	0.1	371	0.2	343	<.1	324	0.2	341	0.1	329	0.1	361	<.1	422	0.2	371	0.1	357
1990		<.1	303	<.1	418	0.3	354	<.1	332	<.1	417	<.1	360	<.1	367	<.1	422	0.1	403	<.1	385
1991		<.1	336	<.1	435	0.1	392	<.1	359	<.1	365	<.1	353	<.1	413	<.1	446	0.1	384	<.1	387
1992		<.1	367	<.1	362	0.1	392	0.2	368	<.1	320	0.1	307	<.1	379	<.1	445	0.1	398	0.1	363
1993		<.1	329	<.1	372	0.2	389	0.1	363	0.1	328	0.1	315	<.1	407	<.1	486	0.1	412	0.1	369
1994		<.1	310	0.1	426	0.2	390	0.1	366	0.2	371	0.1	365	<.1	406	<.1	453	0.1	377	0.1	383
1995	0.1	341	<.1	356	0.2	412	0.1	354	0.2	377	0.1	377	<.1	352	<.1	339	0.1	375	0.1	373	
Southern flounder	1975	0.1	<.1	317	ND	ND	0.1	323	0.1	250	0.1	309	0.2	380	0.1	448	0.1	338	0.1	342	
	1976	ND	<.1	365	0.5	321	<.1	296	0.2	363	0.1	304	0.2	351	0.1	347	0.1	389	0.1	348	
	1977	ND	0.2	331	0.2	342	<.1	322	0.2	312	0.2	368	0.1	383	<.1	491	<.1	353	0.1	342	
	1978	ND	0.1	359	0.1	354	<.1	310	0.1	310	0.1	377	0.2	372	<.1	354	<.1	335	0.1	352	
	1979	ND	<.1	348	0.1	331	0.1	338	0.2	388	0.1	336	0.1	347	0.1	396	0.2	366	0.2	363	
	1980	ND	0.2	345	0.3	369	0.2	330	0.1	325	0.1	359	0.2	367	<.1	363	0.2	400	0.1	354	
	1981	ND	0.1	326	0.1	351	0.1	335	0.1	311	0.1	356	0.1	348	0.1	387	0.1	358	0.1	346	
	1982	ND	0.2	345	0.3	354	0.1	350	0.2	311	0.1	360	0.1	353	0.1	349	0.2	354	0.2	346	
1983	ND	0.1	348	0.2	350	0.1	324	0.2	342	0.2	342	<.1	367	0.1	345	0.1	389	0.1	351		

Table 2. (Cont'd.)

Species	Year	Sabine Lake		Galveston		East		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Southern flounder (Cont'd.)	1984	ND		0.1	341	0.2	364	<.1	328	0.1	322	0.1	323	0.1	326	0.1	293	0.1	326	
	1985	ND		0.1	340	0.2	370	0.1	333	0.1	330	0.1	336	0.1	337	0.2	347	0.1	339	
	1986	0.1	299	0.1	363	0.1	376	0.1	346	0.1	377	<.1	371	0.1	368	0.2	363	0.1	361	
	1987	0.1	335	0.1	336	0.1	350	0.1	308	0.1	345	0.1	394	0.1	391	<.1	381	0.1	402	
	1988	<.1	346	0.1	350	0.2	353	0.1	342	0.1	340	0.1	372	<.1	372	<.1	419	0.1	387	
	1989	<.1	324	0.1	349	0.2	362	0.1	328	0.1	353	0.1	342	<.1	342	0.1	392	0.1	352	
	1990	<.1	325	0.1	326	0.2	340	0.1	324	0.1	324	0.1	344	0.1	333	0.1	279	0.1	340	
	1991	<.1	313	<.1	354	0.1	371	0.1	332	0.1	354	0.1	366	0.1	384	0.1	365	0.1	360	
	1992	<.1	330	0.1	356	0.3	375	0.1	352	<.1	370	0.1	385	0.1	379	<.1	461	0.1	374	
	1993	<.1	350	0.1	379	0.2	426	0.1	364	0.1	411	0.1	411	0.1	388	<.1	352	0.1	387	
	1994	<.1	373	0.1	361	0.2	401	0.1	378	0.1	378	0.1	386	0.1	383	<.1	416	<.1	377	
	1995	0.1	349	<.1	360	0.1	407	0.1	357	0.1	351	0.1	390	0.1	382	<.1	420	<.1	372	
	Atlantic croaker	1975	0.0		<.1	245	ND		0.0	312	0.1	312	0.2	338	0.4	321	0.1	314	0.1	343
		1976	ND		0.2	262	0.1	248	0.3	263	0.4	296	0.2	314	0.6	320	0.5	329	0.3	326
		1977	ND		0.1	291	0.1	275	0.2	274	0.2	290	0.8	307	0.6	350	0.7	345	0.2	340
1978		ND		0.1	274	0.1	248	0.2	255	0.1	242	0.5	314	0.4	296	0.4	283	<.1	331	
1979		ND		<.1	271	0.2	248	0.1	287	0.2	270	0.2	303	0.5	305	0.2	316	0.2	331	
1980		ND		0.2	284	0.1	262	0.2	261	0.1	264	0.2	320	1.7	320	0.1	302	0.2	298	
1981		ND		0.2	279	0.2	254	0.1	273	0.2	268	0.7	328	0.8	320	0.2	323	0.4	320	
1982		ND		0.4	282	0.4	256	0.1	277	0.2	278	0.4	328	1.0	327	0.4	338	0.3	330	
1983		ND		0.3	275	0.4	261	0.2	263	0.5	286	0.3	309	1.0	320	0.1	312	0.5	314	
1984		ND		0.2	274	0.2	259	0.2	259	0.2	252	0.1	261	0.5	274	0.1	264	0.2	270	
1985		ND		0.6	272	0.4	258	0.1	254	0.1	261	0.3	268	0.6	279	0.2	307	0.3	281	
1986		0.2	296	0.4	281	0.1	261	0.2	256	0.2	256	0.3	280	1.4	305	0.1	322	0.3	299	
1987		0.1	287	0.8	288	0.1	252	0.3	253	<.1	253	0.2	283	1.5	323	0.1	321	0.3	322	
1988		0.2	276	0.6	291	0.1	267	0.3	255	0.2	255	0.3	318	0.8	317	0.1	357	0.3	318	
1989		0.1	284	0.6	271	0.2	257	0.2	250	0.2	262	0.2	266	0.3	317	<.1	324	<.1	308	
1990	0.2	283	0.4	286	0.2	270	0.1	261	<.1	260	0.1	261	0.3	290	<.1	298	0.1	264		
1991	0.1	271	0.2	274	0.1	290	0.2	260	0.2	251	0.2	262	0.4	283	<.1	269	1.4	279		
1992	0.2	293	0.4	269	0.1	278	0.1	258	0.1	268	0.3	278	1.0	299	0.1	328	0.7	291		
1993	0.1	286	1.4	273	0.2	276	0.1	265	0.2	267	0.1	281	1.0	313	<.1	306	0.3	300		
1994	0.1	277	0.3	283	0.1	295	0.1	270	0.2	265	0.1	293	1.0	310	0.1	336	0.1	324		
1995	0.1	272	0.4	284	0.3	301	0.1	271	0.3	272	0.4	285	0.6	343	0.1	331	0.1	321		
Gafftop-sail catfish	1975	<.1	530	0.0		ND		0.1	571	<.1	493	<.1	552	0.1	575	0.0	0.0	0.0	567	
	1976	ND		0.1	482	0.0		0.2	526	0.4	498	<.1	587	<.1	475	0.0	0.0	0.1	509	
	1977	ND		<.1	516	0.0		<.1	499	0.2	526	<.1	385	<.1	600	0.1	529	0.0	516	
	1978	ND		0.0		0.0		<.1	514	<.1	543	0.0		0.1	551	0.0		0.1	534	
	1979	ND		0.0		0.2	542	0.0		0.1	499	<.1	533	0.0		0.0		0.1	511	
	1980	ND		0.1	550	0.0		<.1	478	0.3	509	0.1	522	0.1	517	0.0	0.0	0.1	525	
	1981	ND		0.1	492	0.0		<.1	505	<.1	542	0.1	511	0.1	523	0.0	0.1	379	0.1	507
	1982	ND		<.1	423	<.1	616	<.1	520	0.3	527	0.1	533	<.1	545	<.1	541	0.0	517	
	1983	ND		<.1	492	0.1	473	0.1	544	0.3	514	0.1	532	0.0		<.1	408	0.1	514	
	1984	ND		<.1	517	0.1	474	0.1	510	0.3	507	0.1	521	0.1	488	0.0		0.1	509	
1985-1988	1985	ND		0.1	525	0.1	482	<.1	498	0.1	546	0.1	556	0.1	519	<.1	556	<.1	528	
	1986	0.1	462	<.1	521	<.1	473	<.1	474	0.2	485	0.1	532	<.1	514	0.0		0.1	495	
	1987	<.1	423	<.1	491	0.1	527	<.1	512	<.1	519	0.1	542	<.1	528	0.0		0.1	514	
	1988	<.1	370	<.1	515	<.1	534	0.2	521	0.1	544	0.1	538	0.1	521	<.1	495	<.1	525	
	1989	<.1	321	<.1	480	0.1	485	0.2	509	0.1	549	0.1	547	0.1	384	0.0		0.1	524	

Table 2. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake		Galveston		Matagorda		Matagorda		San Antonio		Arkansas		Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
Gafftop-sail catfish (Cont'd.)	1990	<.1	465	0.1	504	0.1	499	0.2	499	0.2	509	<.1	583	0.1	549	<.1	598	<.1	429	0.1	513
	1991	<.1	469	0.1	502	0.1	518	<.1	476	<.1	562	<.1	569	<.1	472	0.0	0.0	<.1	479	<.1	299
	1992	<.1	464	0.1	444	0.1	556	0.1	519	0.1	565	<.1	541	<.1	496	<.1	495	<.1	406	0.1	508
	1993	0.0	513	0.1	566	0.1	501	0.1	501	0.1	538	<.1	585	0.1	473	0.0	0.0	<.1	414	0.1	524
	1994	<.1	409	0.1	441	0.1	501	0.2	516	0.2	541	0.1	561	0.1	511	0.0	0.0	<.1	419	0.1	509
	1995	<.1	380	0.1	408	0.2	511	0.2	503	0.2	522	0.1	565	0.1	462	0.0	0.0	<.1	340	0.1	498
Gulf menhaden	1975	0.0	ND	ND	272	0.5	272	1.7	302	0.4	221	0.2	307	0.5	284	0.3	280	0.1	312	0.5	286
	1976	ND	246	0.1	240	0.3	246	0.3	246	0.3	275	0.1	277	0.5	275	0.2	304	0.1	275	0.8	255
	1977	ND	246	<.1	248	0.2	244	0.2	244	0.1	240	<.1	237	2.0	254	1.4	258	0.1	211	1.0	249
	1978	ND	249	0.5	249	<.1	241	<.1	241	0.1	239	0.6	242	1.4	250	0.2	254	0.0	0.0	0.4	248
	1979	ND	249	0.1	249	0.1	231	0.4	250	<.1	235	0.1	251	0.3	251	0.1	251	<.1	294	0.1	252
	1980	ND	253	0.0	253	0.0	260	<.1	260	0.1	255	0.1	245	<.1	243	0.6	249	0.1	325	0.2	254
	1981	ND	259	0.7	259	<.1	260	0.1	246	0.1	242	0.1	238	0.3	255	0.7	262	0.1	273	0.3	258
	1982	ND	251	0.6	251	<.1	310	<.1	246	0.1	243	<.1	238	0.8	255	0.1	264	<.1	239	0.2	252
	1983	ND	257	1.7	257	0.1	248	<.1	249	0.2	239	0.2	246	0.2	258	<.1	290	<.1	250	0.5	255
	1984	ND	256	1.0	256	0.2	255	0.4	248	0.4	246	0.6	251	0.5	254	0.2	273	0.2	295	0.5	259
	1985	ND	249	1.5	249	<.1	233	0.1	254	0.1	249	0.1	263	0.5	260	0.2	281	0.1	279	0.4	253
	1986	0.2	246	1.5	244	0.1	233	0.3	239	0.1	244	0.1	249	0.8	263	<.1	249	<.1	262	0.5	247
	1987	0.1	244	1.8	250	0.0	206	0.1	244	<.1	278	<.1	250	0.2	259	<.1	256	<.1	278	0.4	250
	1988	0.2	268	0.8	244	<.1	206	0.2	233	0.1	241	<.1	252	0.1	264	<.1	249	0.1	317	0.2	247
	1989	0.2	253	0.8	245	<.1	236	0.2	231	<.1	240	<.1	276	<.1	252	0.0	0.0	<.1	253	0.2	244
	1990	0.1	256	1.3	253	<.1	247	0.6	224	0.1	251	0.1	214	<.1	294	0.0	0.0	<.1	226	0.4	247
	1991	0.3	255	1.4	257	0.0	217	<.1	217	<.1	239	<.1	229	<.2	256	<.1	287	<.1	240	0.3	256
	1992	<.1	299	1.3	257	<.1	232	0.1	239	0.1	245	<.1	257	0.1	271	<.1	266	0.1	237	0.3	256
	1993	0.4	283	1.0	254	<.1	255	0.2	269	<.1	300	0.0	0.0	0.1	239	<.1	281	0.3	301	0.3	262
1994	0.2	240	0.5	254	<.1	210	0.1	249	<.1	266	<.1	268	0.2	256	<.1	96	<.1	282	0.1	254	
1995	0.2	250	2.5	254	<.1	237	0.1	245	0.1	256	<.1	268	0.1	230	<.1	271	<.1	316	<.1	254	
Striped mullet	1975	<.1	390	0.3	331	0.3	331	0.4	347	0.6	322	2.5	328	1.0	382	0.3	358	0.5	345	0.7	339
	1976	ND	346	0.3	346	0.2	320	0.3	349	1.6	331	0.5	360	0.3	342	0.6	402	2.0	397	0.7	367
	1977	ND	345	0.2	345	0.2	380	0.4	330	0.9	343	0.3	321	0.4	371	0.3	396	0.6	354	0.4	348
	1978	ND	423	0.2	423	0.6	330	0.6	342	0.5	322	1.1	336	0.1	336	0.1	364	0.3	387	0.4	347
	1979	ND	351	0.1	351	0.1	338	0.3	340	0.7	344	0.7	344	0.3	353	0.6	410	0.3	365	0.4	357
	1980	ND	363	0.2	363	<.1	319	0.2	343	0.6	357	0.6	357	0.3	340	0.3	360	0.4	346	0.3	353
	1981	ND	395	0.1	395	0.1	349	0.1	332	0.6	341	0.5	334	0.3	353	0.3	364	0.9	363	0.4	352
	1982	ND	376	0.2	376	0.4	329	0.3	330	0.4	341	0.8	331	0.2	345	0.1	348	0.4	372	0.4	347
	1983	ND	370	0.2	370	0.2	335	0.2	339	0.3	334	0.5	350	0.2	347	0.3	383	0.6	375	0.3	358
	1984	ND	362	0.4	362	0.7	328	0.3	331	0.5	350	0.6	342	0.4	357	0.5	376	0.4	356	0.5	352
	1985	ND	338	0.2	338	0.2	326	0.2	323	0.3	355	0.3	343	0.2	342	0.3	397	0.3	375	0.3	354
	1986	<.1	328	0.1	377	0.3	328	0.1	337	0.4	369	0.2	356	0.2	358	<.1	370	0.6	359	0.2	359
	1987	<.1	325	0.2	375	0.4	333	0.7	319	1.1	360	0.6	348	0.3	338	0.2	391	0.4	382	0.5	351
	1988	<.1	331	0.2	362	0.4	344	0.4	326	0.4	347	0.4	365	0.3	370	0.4	409	0.4	396	0.3	366
	1989	<.1	329	0.2	349	0.2	334	0.2	328	0.3	350	0.4	348	0.2	359	0.3	394	0.4	366	0.3	357
	1990	0.1	334	0.4	341	0.3	368	0.2	344	0.8	369	0.7	358	0.2	353	0.3	387	0.4	383	0.4	361
	1991	0.1	331	0.2	333	0.6	366	0.1	343	0.8	364	0.5	351	0.3	368	0.1	383	0.4	401	0.3	363
	1992	<.1	328	0.3	376	0.3	387	0.4	330	0.2	350	0.7	364	0.4	360	0.2	389	0.3	383	0.4	362
	1993	0.6	328	0.9	364	0.7	377	0.5	352	0.7	374	0.9	365	0.4	376	0.6	422	0.4	402	0.7	373
1994	0.1	353	0.6	372	0.4	384	0.6	347	0.3	358	0.7	365	0.2	379	0.1	398	0.2	400	0.4	367	
1995	0.2	353	0.4	371	0.3	397	0.4	356	0.4	347	0.5	370	0.3	368	0.4	421	0.2	375	0.3	374	

Table 2. (Cont'd.)

Species	Year	Sabine Lake		Galveston		East		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
Total finfishes	1975	3.0	383	5.1	396	ND	ND	6.6	355	4.9	339	7.9	345	5.7	343	4.3	374	4.8	394	5.5	365
	1976	ND		7.2	334	4.0	385	4.9	388	9.1	365	5.0	363	5.0	349	5.1	383	11.1	400	6.8	369
	1977	ND		6.2	334	3.2	362	5.4	389	6.2	348	3.6	344	5.8	326	5.2	343	6.5	381	5.5	353
	1978	ND		4.0	342	4.0	325	5.0	359	5.1	383	5.2	341	3.8	322	3.6	358	3.1	395	4.3	355
	1979	ND		3.5	367	2.0	372	4.3	350	5.6	368	3.8	372	3.5	327	2.6	367	3.5	393	3.7	365
	1980	ND		4.0	371	2.9	375	3.3	346	6.1	342	4.8	350	5.0	336	2.5	354	4.2	390	4.3	357
	1981	ND		4.2	357	3.3	355	3.0	384	4.8	358	4.4	375	4.8	364	3.1	357	5.5	388	4.2	369
	1982	ND		6.2	346	6.2	354	3.7	372	5.1	360	4.5	366	5.1	338	3.5	363	5.9	381	5.0	360
	1983	ND		6.0	350	6.2	341	4.0	378	5.3	352	3.9	396	5.8	356	3.0	362	5.5	399	4.9	367
	1984	ND		6.5	364	5.7	379	4.4	369	3.9	362	3.8	399	4.2	347	3.1	373	4.2	406	4.6	373
	1985	ND		7.1	335	4.5	366	3.7	380	4.2	376	3.3	396	4.0	358	3.4	362	4.6	390	4.6	364
	1986	2.6	395	6.0	349	4.4	390	4.6	379	4.7	408	4.0	378	5.3	347	2.2	381	5.2	404	4.6	377
	1987	2.2	430	5.8	334	4.7	390	5.0	323	5.2	428	3.3	391	4.9	353	1.6	406	4.6	444	4.4	374
	1988	2.2	394	6.8	363	5.2	387	4.3	361	5.8	393	4.3	382	5.0	358	3.1	396	5.7	410	5.2	374
	1989	2.4	401	5.2	343	4.9	387	4.2	345	5.5	399	4.5	400	4.5	398	2.7	433	4.5	431	4.4	384
	1991	3.1	389	5.4	341	5.4	376	4.9	373	6.0	408	4.9	373	6.3	371	4.0	397	7.6	389	5.5	372
	1992	2.7	439	6.1	356	6.1	439	5.6	366	6.0	408	6.2	419	5.8	377	3.4	425	7.3	399	5.7	391
	1993	2.7	379	6.9	347	7.1	457	5.8	380	7.5	430	6.4	455	7.3	394	6.3	443	7.9	425	6.7	407
	1994	3.1	374	6.4	372	6.7	428	5.2	381	6.4	404	5.4	402	5.5	386	7.0	444	6.1	443	5.9	403
	1995	3.9	391	7.1	332	7.2	417	6.0	361	5.9	374	6.4	383	5.8	403	6.5	443	5.7	435	6.2	383
Blue crab	1983	ND		0.1	136	0.3	153	0.1	151	0.1	138	0.2	146	0.2	146	0.3	146	0.3	146	0.2	144
	1984	ND		0.1	151	0.1	140	0.1	147	0.1	147	0.2	145	0.2	141	0.2	138	0.2	148	0.1	145
	1985	ND		<.1	149	0.1	154	<.1	142	0.1	139	0.1	141	0.1	143	0.2	147	0.1	148	0.1	145
	1986	0.2	150	<.1	146	<.1	144	<.1	161	0.1	146	<.1	138	0.1	144	<.1	147	0.1	149	<.1	147
	1987	0.2	154	0.1	140	0.1	158	0.2	154	0.3	153	<.1	158	0.1	157	0.3	157	0.1	152	0.2	153
	1988	0.2	155	0.1	144	0.2	150	<.1	137	0.1	138	0.1	145	0.1	147	<.1	129	<.1	152	0.1	147
	1989	0.1	157	<.1	136	<.1	144	<.1	139	<.1	133	<.1	148	<.1	159	0.0	129	<.1	152	<.1	143
	1990	0.2	146	0.1	149	0.1	144	0.2	144	0.1	144	<.1	149	0.1	138	0.1	129	0.2	142	0.1	144
	1991	0.1	152	<.1	151	0.1	152	0.1	131	0.2	150	<.1	136	0.1	153	0.1	139	0.2	148	0.1	146
	1992	0.1	161	<.1	143	0.1	156	0.1	153	<.1	136	0.1	140	<.1	148	0.2	138	0.1	152	0.1	144
	1993	0.1	169	<.1	145	0.1	150	<.1	156	<.1	146	<.1	160	0.1	155	<.1	157	<.1	142	<.1	153
	1994	0.1	163	<.1	152	0.1	151	<.1	155	<.1	150	<.1	154	<.1	147	<.1	140	<.1	125	<.1	151
	1995	0.1	158	<.1	146	0.1	151	<.1	127	0.1	154	<.1	144	<.1	150	<.1	124	0.1	136	<.1	142

Table 3. Annual mean catch rate (No./ha) and mean total lengths (mm) of selected fishes and shellfishes caught with 18.3-m bag seines by bay system during 1977-95. Blank indicated no measurement taken; ND = no data.

Species	Sabine Lake		Galveston		Past		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide		
	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	
FINFISHES																					
Red drum																					
1977*	ND	35	20	35	ND	8	51	85	51	14	44	1	41	0	0	1	39	18	46		
1978	ND	67	3	67	ND	4	43	13	51	4	94	3	67	11	58	17	52	7	58		
1979	ND	17	62	17	62	ND	6	92	11	5	92	18	85	27	66	15	64	14	70		
1980	ND	59	74	59	74	ND	8	68	28	50	5	88	16	75	4	82	15	72	23		
1981	ND	26	52	26	52	ND	9	86	29	53	30	38	40	46	5	46	45	56	26		
1982	ND	53	62	53	62	ND	9	76	19	102	26	103	21	62	1	55	16	89	24		
1983	ND	47	67	47	67	11 <sup>b</sup>	66	7	99	12	98	7	88	2	59	41	92	20 <sup>b</sup>	78		
1984	ND	13	66	13	66	6	70	12	56	4	100	4	80	2	52	4	73	6	69		
1985	ND	3	131	3	131	10	106	7	114	19	82	9	67	1	61	17	69	9	86		
1986	19	87	8	86	2	78	6	105	1	117	1	117	4	98	3	84	22	94	7		
1987	6	99	45	58	47	61	16	88	15	89	9	59	7	71	2	117	32	63	21		
1988	13	78	8	78	27	79	3	114	6	89	10	78	9	49	4	66	21	63	10		
1989	61	44	3	59	24	4	92	10	82	8	65	4	83	1	54	18	69	9	63		
1990	5	62	17	53	27	50	14	51	19	50	43	40	20	57	4	32	25	39	20		
1991	6	97	14	73	30	53	30	81	36	79	25	61	28	72	5	88	43	61	24		
1992	5	71	9	85	23	56	15	77	22	83	7	74	14	14	14	51	13	70	14		
1993	11	73	28	58	40	48	11	66	17	58	13	60	12	73	9	53	23	62	18		
1994	7	72	22	59	31	55	6	58	13	69	29	58	53	55	11	61	26	70	21		
1995	20	69	12	48	18	54	7	69	15	76	15	62	26	62	7	86	29	62	15		
Spotted seatrout																					
1977*	ND	87	34	87	ND	39	84	50	73	1	99	7	84	16	83	5	85	23	82		
1978	ND	52	35	52	ND	6	86	11	69	8	50	4	59	14	93	2	52	14	61		
1979	ND	79	37	79	ND	3	83	12	70	7	68	12	53	13	80	2	86	14	75		
1980	ND	17	72	17	72	ND	3	84	21	71	11	74	11	79	3	56	<1	60	10		
1981	ND	16	85	16	85	ND	7	110	9	68	13	70	12	65	4	73	6	84	10		
1982	ND	37	82	37	82	ND	7	99	19	62	15	76	4	75	5	78	3	76	15		
1983	ND	26	84	26	84	4 <sup>b</sup>	101	7	73	8	72	14	4	79	5	101	4	80	11 <sup>b</sup>		
1984	ND	7	71	7	71	2	85	1	83	10	74	1	54	1	88	5	98	4	77		
1985	ND	5	80	5	80	24	73	4	64	24	61	3	50	9	70	2	78	9	70		
1986	2	67	2	85	17	66	5	71	5	78	12	60	4	68	1	72	2	58	5		
1987	2	92	22	73	14	68	3	82	19	70	13	69	10	76	1	104	3	63	11		
1988	7	88	6	88	14	75	5	96	7	67	28	68	7	65	5	65	3	87	9		
1989	5	63	6	79	14	80	6	69	20	61	16	71	6	71	4	50	2	56	8		
1990	3	56	5	56	10	74	8	66	8	61	14	61	13	65	2	54	<1	86	7		
1991	1	67	16	63	13	71	15	70	34	59	20	65	8	72	6	63	2	59	14		
1992	2	73	6	73	4	82	10	59	42	52	12	64	8	69	18	50	2	54	12		
1993	5	84	6	61	19	71	6	62	15	54	12	68	7	69	14	59	7	49	9		
1994	3	73	5	64	13	78	13	68	12	66	28	76	7	69	5	79	1	51	10		
1995	17	70	5	77	23	76	17	63	16	77	33	68	5	89	19	73	3	75	14		
Black drum																					
1977*	ND		0		ND	11	147	6	179	1	142	1	150	0	106	0	3	156			
1978	ND		36	95	ND	9	112	22	110	2	165	1	122	4	106	0	13	102			



Table 3 (Cont'd.)

Species Year	East				Corpus Christi				Upper Laguna Madre		Lower Laguna Madre		Coastwide					
	Sabine Lake No./ha	Galveston No./ha	Matagorda No./ha	Matagorda Length	San Antonio No./ha	Aransas No./ha	Christi No./ha	Upper Laguna No./ha	Upper Laguna Length	Lower Laguna No./ha	Lower Laguna Length	Coastwide No./ha		Coastwide Length				
<b>Southern flounder (Cont'd.)</b>																		
1982	ND	9	67	ND	3	82	6	56	18	37	2	62	1	53	13	39	8	51
1983	ND	9	46	1 <sup>b</sup>	2	54	3	58	6	39	1	34	0	0	2	45	4 <sup>b</sup>	46
1984	ND	2	83	2	1	78	1	67	3	62	3	45	1	86	1	64	3	69
1985	ND	4	58	5	2	112	1	43	7	55	5	55	<1	71	2	67	3	64
1986	2	83	4	83	6	70	19	66	4	64	2	54	1	79	12	44	6	63
1987	2	47	21	51	9	54	3	44	1	103	1	37	<1	69	3	56	6	53
1988	15	66	14	61	3	76	3	85	5	48	1	65	<1	60	5	60	6	63
1989	10	74	3	62	10	60	3	67	10	51	8	53	<1	106	2	62	7	50
1990	12	68	22	59	12	55	15	48	3	55	12	47	4	67	9	51	12	54
1991	7	58	5	34	7	56	3	53	1	55	2	46	<1	27	2	60	3	49
1992	7	66	3	41	3	67	3	48	1	41	5	44	<1	22	<1	56	2	46
1993	4	95	6	56	6	46	3	47	2	57	3	69	<1	130	2	54	4	55
1994	2	94	4	62	3	58	3	46	5	54	6	42	<1	34	2	78	3	56
1995	4	65	4	59	5	63	3	42	6	41	5	46	1	58	<1	93	3	53
<b>Atlantic croaker</b>																		
1977 <sup>a</sup>	ND	20	96	ND	0	59	0	100	1	36	11	50	1	181	4	83	6	88
1978	ND	320	61	ND	239	59	10	58	37	73	1	30	11	86	29	38	121	61
1979	ND	463	52	ND	109	74	52	49	7	76	25	65	3	92	221	44	162	53
1980	ND	1,085	55	ND	82	69	17	89	16	56	24	49	1	40	198	42	290	54
1981	ND	528	57	ND	24	94	26	73	26	42	20	55	1	112	32	46	136	58
1982	ND	1,812	61	ND	165	74	67	67	142	61	32	54	0	0	49	53	471	62
1983	ND	888	55	56 <sup>b</sup>	79	236	66	80	63	62	6	61	2	86	49	51	254 <sup>b</sup>	58
1984	ND	815	59	210	64	483	60	25	83	155	68	1,160	4	102	133	59	404	60
1985	ND	242	64	121	63	299	72	13	88	46	78	4	76	11	87	87	42	122
1986	126	74	148	77	198	68	2,138	52	17	72	12	78	<1	89	62	57	364	55
1987	79	335	54	110	56	207	78	33	47	9	4	40	<1	60	10	62	113	61
1988	154	485	53	160	51	60	80	13	66	3	8	50	0	0	15	63	125	56
1989	111	36	77	190	45	22	56	9	49	18	10	61	0	0	9	38	27	59
1990	97	316	51	117	46	82	68	24	32	58	14	59	2	78	46	62	103	55
1991	208	57	635	52	343	47	1,035	58	156	63	35	66	11	36	169	46	353	55
1992	225	56	505	47	450	47	626	48	430	44	95	50	13	54	157	44	326	47
1993	232	64	358	50	421	44	216	47	25	66	25	53	2	67	195	40	165	48
1994	255	52	229	49	186	58	302	43	59	46	74	39	6	44	123	46	143	46
1995	357	57	112	48	247	50	110	57	36	69	24	35	2	58	179	41	95	50
<b>Sand seatrout</b>																		
1977 <sup>a</sup>	ND	0	0	ND	11	61	0	0	0	0	0	0	0	0	0	0	2	61
1978	ND	13	58	ND	3	59	0	0	0	0	<1	54	0	0	0	0	4	58
1979	ND	35	58	ND	14	70	2	75	<1	33	1	77	0	0	0	10	61	0
1980	ND	8	61	ND	7	82	<1	64	<1	89	0	0	0	0	0	0	3	69
1981	ND	21	60	ND	2	72	0	0	0	0	1	76	0	0	<1	78	5	61
1982	ND	47	57	ND	12	67	<1	35	<1	76	<1	73	0	0	<1	65	13	58
1983	ND	47	53	10 <sup>b</sup>	59	64	<1	47	1	70	2	53	0	0	0	0	15 <sup>b</sup>	56
1984	ND	49	55	7	66	22	54	0	0	0	0	0	0	0	8	41	15	54
1985	ND	11	60	8	59	12	71	0	<1	67	1	82	0	<1	60	0	5	65



Table 3 (Cont'd.)

Species Year	Sabine Lake		Galveston		East Matagorda		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha
Sand seatrout (Cont'd.)																				
1986	6	71	9	50	4	60	9	64	0	0	0	0	0	57	0	0	0	0	3	57
1987	4	63	16	58	11	61	14	65	1	61	0	0	0	0	0	0	0	0	6	61
1988	5	54	5	53	38	40	6	66	<1	69	0	0	0	0	0	0	0	0	3	52
1989	9	54	43	55	7	66	4	68	<1	31	0	0	106	0	0	0	0	0	10	56
1990	24	52	75	46	10	59	13	56	1	36	0	0	0	0	0	0	0	0	19	47
1991	7	48	76	55	25	59	39	56	<1	76	3	50	2	42	0	0	<1	65	23	55
1992	7	54	30	53	10	52	36	54	0	<1	81	0	0	61	0	0	0	12	53	51
1993	7	58	53	48	19	53	88	53	4	64	<1	96	1	57	0	2	54	26	26	51
1994	4	61	34	48	16	70	29	56	<1	70	<1	64	0	0	0	<2	53	12	52	48
1995	14	50	59	44	52	38	37	59	<1	56	<1	64	0	0	0	1	57	20	20	48
Gulf menhaden																				
1977*	ND	21	76	76	ND	ND	0	0	0	0	0	0	1	58	0	0	0	0	5	76
1978	ND	533	31	53	ND	3,963	47	169	64	3,310	44	44	41	41	44	42	71	29	1,249	44
1979	ND	122	53	53	ND	867	43	817	0	817	38	38	335	38	6	37	1	31	312	41
1980	ND	14,717	46	46	ND	115	50	24	52	48	30	7	49	4	40	54	54	31	3,343	46
1981	ND	196	45	45	ND	348	51	52	41	355	48	8	41	721	42	11	38	246	45	47
1982	ND	4,788	50	50	ND	820	48	1,008	37	137	33	1,068	36	31	130	32	130	32	1,466	47
1983	ND	4,971	66	66	1,324 <sup>b</sup>	44	809	44	67	42	16	34	619	33	2	30	5	47	1,312 <sup>b</sup>	62
1984	ND	1,839	44	44	470	48	1,260	45	1,084	42	866	39	553	52	128	49	69	56	928	44
1985	ND	486	42	243	43	3,819	50	868	45	48	39	122	37	62	44	20	49	36	819	48
1986	3,049	48	3,024	38	1,502	37	10,076	53	612	36	27	34	11	46	36	44	12	36	2,333	48
1987	633	47	264	50	755	49	3,550	60	35	40	68	11	34	32	63	18	27	637	57	57
1988	600	40	2,625	45	438	41	363	60	<1	43	80	30	<1	44	14	31	81	35	660	45
1989	526	48	781	42	386	51	187	45	53	37	43	37	11	43	2	45	71	39	245	43
1990	774	49	5,106	43	640	44	527	56	797	71	943	35	869	32	21	38	<1	38	1,487	44
1991	270	41	4,298	40	1,258	42	3,044	42	296	42	569	41	244	38	123	36	0	0	1,533	41
1992	593	45	6,025	37	291	36	1,919	38	1,810	35	259	33	43	46	4	30	21	40	1,815	37
1993	1,878	48	7,341	40	509	36	492	46	191	38	634	66	158	38	13	39	108	35	1,827	42
1994	72	51	5,203	48	222	41	418	57	138	39	263	33	15	45	32	33	<1	49	1,212	48
1995	399	46	6,155	39	2,381	50	718	36	140	48	373	37	83	40	133	34	4	48	1,570	39
Pinfish																				
1977*	ND	0	114	55	ND	32	114	24	105	22	105	22	105	66	93	167	102	13	39	103
1978	ND	116	73	75	ND	24	61	77	75	54	74	54	74	133	69	41	84	7	64	65
1979	ND	151	38	38	ND	16	50	363	57	167	66	250	61	17	88	153	59	107	47	77
1980	ND	270	55	55	ND	68	69	131	70	107	85	267	67	40	84	132	75	59	152	55
1981	ND	144	67	67	ND	34	66	590	55	448	67	265	62	100	73	349	57	260	61	66
1982	ND	138	65	65	61 <sup>b</sup>	79	115	80	510	49	642	68	533	66	25	82	211	68	279 <sup>b</sup>	64
1983	ND	247	59	59	180	64	107	71	172	66	471	62	214	54	146	79	120	77	214	64
1984	ND	362	55	55	401	65	209	71	396	55	274	66	234	67	133	68	261	66	280	62
1985	64	74	183	61	676	64	117	58	161	66	696	59	304	58	245	62	329	63	287	61
1986	8	72	50	64	227	57	44	442	63	321	67	463	58	42	56	339	64	206	63	60
1987	7	84	128	61	373	62	43	77	246	63	589	62	983	54	312	59	660	60	357	60
1988	24	75	80	62	359	58	308	53	607	61	300	63	361	57	60	70	251	61	254	60
1989	37	75	182	58	499	61	251	52	552	52	609	55	566	57	392	62	660	60	415	58

Table 3 (Cont'd.)

Species Year	Sabine Lake		Galveston		Past Matagorda		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length
<b>Pinfish (Cont'd.)</b>																				
1991	8	79	138	58	307	60	39	68	248	65	119	61	435	63	240	69	696	57	243	61
1992	12	73	96	46	371	56	67	49	431	53	545	59	475	50	174	59	531	58	293	55
1993	27	78	309	49	139	59	150	53	368	60	564	56	482	54	307	59	452	60	344	56
1994	9	71	164	50	285	66	125	57	174	58	463	58	411	58	102	56	358	56	237	57
1995	12	86	159	49	284	52	73	51	308	59	333	64	326	54	165	59	247	60	212	57
<b>Spot</b>																				
1977*	ND	ND	56	100	ND	23	118	0	0	2	170	12	100	12	0	0	1	125	18	105
1978	ND	ND	407	52	ND	182	49	361	48	80	55	310	47	227	59	149	52	253	51	51
1979	ND	ND	352	42	ND	21	64	201	44	58	60	210	55	103	70	57	59	156	49	49
1980	ND	ND	269	57	ND	76	56	256	51	101	61	95	58	86	59	165	48	160	55	55
1981	ND	ND	331	52	ND	154	57	135	64	97	54	121	61	115	63	220	67	185	58	58
1982	ND	ND	404	62	ND	143	58	467	52	623	54	225	60	180	58	340	66	350	58	58
1983	ND	ND	459	57	50 <sup>a</sup>	64	95	58	169	47	350	56	135	55	57	60	526	63	273 <sup>b</sup>	58
1984	ND	ND	238	53	96	61	146	58	247	46	659	56	564	58	493	66	948	67	433	60
1985	ND	ND	179	62	158	59	216	59	274	44	254	64	227	55	80	77	169	54	197	58
1986	118	65	135	68	319	56	825	51	102	58	258	51	160	60	114	55	614	54	314	54
1987	19	80	264	60	383	60	83	58	203	49	476	58	359	49	17	70	307	47	239	55
1988	44	62	229	69	210	66	116	64	361	59	158	65	212	54	270	59	209	62	209	62
1989	96	52	87	63	256	58	173	59	264	62	253	53	158	62	271	50	151	64	183	58
1990	16	70	222	62	525	54	330	57	691	51	566	52	831	49	684	57	854	55	525	54
1991	22	65	270	56	304	59	131	49	198	69	295	53	279	52	174	53	950	51	314	54
1992	27	70	211	55	89	61	63	53	194	59	164	53	387	45	219	58	347	54	204	54
1993	35	80	164	56	288	55	123	53	149	50	185	59	281	58	221	62	341	53	197	56
1994	55	78	369	49	161	61	99	61	127	56	310	62	250	59	66	60	369	54	231	55
1995	15	104	171	50	199	56	254	50	76	65	191	59	302	53	145	59	217	57	184	55
<b>Striped mullet</b>																				
1977*	ND	ND	31	140	ND	129	106	129	117	27	132	179	156	15	158	62	103	74	126	74
1978	ND	ND	56	120	ND	26	124	126	66	68	103	121	76	53	94	105	81	74	90	90
1979	ND	ND	135	89	ND	93	99	273	66	152	103	202	135	16	102	383	53	174	81	81
1980	ND	ND	90	117	ND	15	107	41	121	61	102	49	88	57	70	95	85	61	100	100
1981	ND	ND	229	57	ND	41	92	249	84	205	81	79	85	31	63	161	98	152	76	76
1982	ND	ND	128	66	ND	553	118	179	77	177	85	29	110	23	86	43	94	174	98	98
1983	ND	ND	85	94	62 <sup>b</sup>	104	26	136	57	64	110	106	37	61	15	99	44	84	57 <sup>b</sup>	94
1984	ND	ND	52	95	33	110	34	69	73	102	57	142	52	154	68	255	96	106	77	77
1985	ND	ND	75	110	199	89	49	22	134	95	58	22	62	70	53	119	81	72	84	84
1986	84	103	34	134	20	144	23	37	93	22	91	62	67	23	57	41	66	35	92	92
1987	48	98	244	75	60	89	33	96	63	115	127	73	141	56	94	37	72	103	116	76
1988	42	80	115	115	69	90	44	64	16	84	50	189	49	64	62	27	125	74	80	80
1989	61	68	41	96	40	61	24	82	10	147	77	47	131	49	33	78	58	55	61	61
1990	43	88	194	71	151	81	21	71	100	156	41	322	44	226	59	114	89	144	63	63
1991	83	94	234	80	162	60	79	65	73	97	40	88	138	41	283	50	126	133	71	71
1992	23	94	149	79	97	78	52	78	72	81	132	80	141	50	70	53	44	99	95	75
1993	74	84	105	83	84	74	41	77	62	71	67	86	133	49	36	39	78	74	71	71
1994	56	75	102	66	29	70	59	75	35	92	53	66	137	48	62	47	221	42	91	57
1995	63	99	45	71	73	84	23	70	122	57	46	19	59	20	57	29	49	35	68	68

Table 3 (Cont'd.)

Species Year	Sabine Lake		Galveston		East Matagorda		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide		
	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	
Total finfishes																					
1977*	ND		959	59	ND		489	88	3,106	52	1,383	64	2,788	60	1,780	67	830	59	1,464	61	
1978	ND		4,103	53	ND		4,855	67	1,671	65	5,038	64	1,515	66	1,282	62	908	54	3,030	61	
1979	ND		3,149	60	ND		1,635	71	3,375	57	3,096	60	2,191	70	1,354	69	2,368	72	2,518	64	
1980	ND		18,543	86	ND		632	77	1,879	67	1,407	68	1,490	67	2,116	63	1,070	59	5,241	82	
1981	ND		3,334	63	ND		1,093	83	1,781	61	2,020	66	2,213	64	1,792	54	1,267	70	2,028	65	
1982	ND		9,007	68	ND		2,077	78	4,321	56	5,021	57	2,596	66	1,355	58	1,342	61	4,194	65	
1983	ND		8,725	71	2,078 <sup>b</sup>		1,857	80	2,147	55	4,059	63	2,160	59	734	61	1,378	68	3,528 <sup>b</sup>	68	
1984	ND		4,644	59	1,617		2,625	62	2,687	58	3,574	62	3,353	52	1,817	60	1,906	71	3,044	60	
1985	ND		1,995	63	1,921		5,152	82	2,200	65	2,514	60	1,389	56	1,534	55	1,458	60	2,383	68	
1986	3,776		69	3,916	71	3,329	63	14,493	73	1,849	60	2,294	57	841	62	1,554	51	1,672	61	4,146	69
1987	1,153		67	2,231	64	2,484	63	4,312	79	1,344	65	2,030	58	1,357	55	1,012	46	1,575	60	2,073	66
1988	1,153		62	4,347	71	2,024	63	913	83	1,391	58	3,150	54	2,344	56	2,271	50	2,144	65	2,464	63
1989	1,243		62	2,157	67	2,097	59	1,362	69	1,997	62	2,079	55	2,006	56	2,360	45	2,341	56	2,010	59
1990	1,319		67	7,186	58	2,951	59	2,106	68	3,470	57	3,968	55	3,913	52	5,385	48	2,993	59	4,209	57
1991	719		62	7,525	62	3,452	63	4,982	69	3,090	63	2,300	59	2,273	60	2,971	50	4,012	54	4,138	61
1992	1,143		56	7,886	54	1,924	57	3,414	57	4,687	53	2,622	57	2,373	52	4,251	47	2,893	55	4,188	54
1993	2,526		62	9,393	64	2,536	54	1,700	60	2,284	54	2,839	65	2,393	53	4,103	48	2,752	54	4,050	60
1994	617		62	6,845	54	1,538	63	1,985	63	1,183	60	2,496	57	1,972	58	2,941	44	2,846	50	3,126	54
1995	1,350		65	7,390	59	5,029	69	1,949	58	1,555	64	2,937	59	1,815	57	3,602	48	3,002	49	3,559	57
SHELLFISHES																					
Blue crab																					
1977*	ND		103	43	ND		31	46	51	46	95	56	56	38	16	58	8	63	56	47	
1978	ND		66	52	ND		10	38	52	51	57	62	33	43	98	61	19	60	48	55	
1979	ND		106	52	ND		27	51	76	49	84	62	152	43	90	48	61	54	83	51	
1980	ND		122	54	ND		24	56	119	45	65	52	80	38	65	40	176	46	95	48	
1981	ND		58	53	ND		43	44	51	54	85	45	86	40	42	58	167	35	74	44	
1982	ND		101	48	ND		31	51	107	42	193	48	52	49	35	54	175	42	102	46	
1983	ND		148	43	15		35	34	105	40	145	43	48	40	36	59	112	33	94	41	
1984	ND		88	58	58		58	42	42	46	63	50	62	42	37	61	80	46	64	51	
1985	ND		144	49	107		54	56	41	42	141	38	184	37	73	52	152	34	113	42	
1986	37		79	90	86		57	53	62	46	30	48	77	40	23	45	91	41	63	49	
1987	23		68	163	41		38	51	64	55	35	35	80	47	50	59	72	44	77	45	
1988	44		64	160	46		31	36	48	42	54	35	89	44	38	43	78	37	78	42	
1989	50		45	85	48		30	25	74	31	56	34	72	43	22	41	31	35	59	38	
1990	67		47	141	44		46	31	98	30	83	35	150	42	37	51	68	40	94	39	
1991	46		56	165	47		44	37	198	38	107	35	158	40	49	45	107	43	117	42	
1992	36		55	90	54		45	26	117	30	140	34	164	38	105	58	129	35	103	37	
1993	36		59	116	35		89	23	89	35	652	41	176	42	67	55	78	36	166	39	
1994	28		51	89	38		176	22	27	34	91	27	210	39	113	47	130	32	102	34	
1995	43		46	59	32		194	22	32	30	56	34	122	37	62	40	97	31	71	32	
Brown shrimp																					
1977*	ND		139	46	ND		64	52	200	49	229	54	99	58	9	63	200	53	137	51	
1978	ND		540	50	ND		167	63	102	63	152	60	258	56	188	68	120	53	245	56	
1979	ND		482	58	ND		194	66	69	63	438	63	499	61	53	59	155	59	285	61	
1980	ND		495	52	ND		143	68	553	60	386	60	183	62	64	64	234	56	314	58	
1981	ND		719	57	ND		157	74	310	64	355	60	679	53	102	76	1,008	58	490	59	

Table 3 (Cont'd.)

Species Year	East								Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide					
	Sabine Lake No./ha	Lake Length No./ha	Galveston Length No./ha	Matsigorda Length No./ha	Matsigorda Length No./ha	San Antonio Length No./ha	Aransas Length No./ha	Christi Length No./ha	Upper Laguna Madre Length No./ha	Lower Laguna Madre Length No./ha	Coastwide Length No./ha	Coastwide Length No./ha								
Brown shrimp (Cont'd.)																				
1982	ND		915	64	ND	207	64	599	51	505	54	428	57	63	565	61	510	60		
1983	ND		484	60	99	248	66	310	57	530	60	295	56	57	532	50	360	58		
1984	ND		628	64	294	65	56	244	66	740	66	291	58	82	389	63	396	64		
1985	ND		522	60	413	59	63	306	56	755	61	370	55	288	70	1,007	56	525	59	
1986	605	74	1,166	58	558	63	524	337	65	231	63	204	58	193	66	627	54	318	62	
1987	401	70	1,162	58	387	56	445	64	158	464	62	293	60	417	56	961	58	610	59	
1988	248	61	516	62	570	57	208	61	206	53	357	58	64	756	73	461	62	416	63	
1989	110	59	519	59	889	56	369	54	739	55	726	51	522	54	167	58	411	59	493	56
1990	127	69	356	56	723	61	477	61	482	56	1,005	60	592	62	77	74	2,128	59	694	59
1991	14	68	601	57	790	61	453	60	624	56	511	67	660	70	248	56	1,064	63	591	61
1992	245	71	708	57	455	55	270	52	455	62	455	62	629	58	328	62	926	55	565	57
1993	102	63	541	58	560	54	232	55	321	54	568	64	636	58	279	62	891	59	482	59
1994	302	62	515	60	480	56	403	61	165	57	513	62	713	63	239	58	841	59	477	60
1995	83	68	331	54	392	50	344	57	290	57	359	57	498	60	477	59	728	59	406	57
Pink shrimp																				
1977*	ND		0		ND	0	0	12	41	0	0	0	0	48	77	0	0	7	69	
1978	ND		0		ND	0	0	<1	100	<1	63	0	0	26	77	0	0	3	77	
1979	ND		0		ND	0	0	0	0	0	0	58	51	12	78	<1	106	7	57	
1980	ND		0		ND	0	0	6	51	13	50	58	55	10	60	2	75	10	55	
1981	ND		0		ND	0	0	28	54	87	44	67	67	8	62	5	49	24	49	
1982	ND		0		ND	0	0	0	0	124	47	67	46	7	61	3	52	25	48	
1983	ND		0		ND	0	0	9	51	50	56	31	47	12	54	0	0	12	53	
1984	ND		0		ND	0	0	0	73	16	48	26	48	14	65	<1	79	6	53	
1985	ND		0		ND	0	0	0	0	17	59	7	49	8	76	0	0	4	61	
1986	0		0		ND	0	0	<1	68	15	39	25	49	6	43	3	65	5	46	
1987	0		0		ND	0	0	0	0	11	52	60	52	14	50	0	0	8	52	
1988	0		0		ND	0	0	<1	38	135	49	106	50	<1	55	6	54	28	50	
1989	0		0		ND	0	0	1	52	45	42	64	46	20	59	0	0	14	47	
1990	0		0		ND	0	0	<1	36	99	49	106	48	4	48	15	51	25	49	
1991	0		0		ND	0	0	<1	110	61	52	25	46	31	42	1	52	14	49	
1992	0		<1	59	0	0	0	1	40	32	53	77	54	38	55	176	59	38	57	
1993	0		<1	34	0	0	0	<1	44	58	47	53	50	32	55	140	56	34	53	
1994	0		2	40	52	56	5	<1	35	103	49	150	53	9	39	235	59	61	54	
1995	0		1	37	16	41	3	4	46	88	50	53	50	7	45	179	57	42	53	
White shrimp																				
1977*	ND		1,586	55	ND	1,054	102	115	47	26	63	84	57	36	85	23	57	553	69	
1978	ND		858	66	ND	554	70	130	61	92	49	62	52	21	55	130	53	335	65	
1979	ND		1,720	61	ND	543	70	212	56	99	64	817	52	5	53	143	47	608	61	
1980	ND		571	64	ND	522	68	291	57	133	61	141	69	62	71	18	45	288	64	
1981	ND		1,393	62	ND	805	59	66	64	183	50	173	51	19	56	264	61	527	60	
1982	ND		3,560	58	ND	1,750	64	650	51	297	43	369	54	14	51	326	50	1,276	58	
1983	ND		1,524	50	348	70	394	65	135	64	129	135	42	7	67	218	52	478	53	
1984	ND		1,557	59	409	65	1,438	71	166	56	415	311	63	17	58	625	58	759	628	
1985	ND		307	61	552	61	584	63	37	44	239	44	33	6	73	204	54	241	58	
1986	308	73	1,389	62	173	65	173	66	140	66	287	44	101	58	2	48	175	49	491	61
1987	682	68	972	53	577	61	579	67	90	54	111	65	152	61	7	37	121	61	386	58
1988	796	63	482	66	429	66	341	68	168	52	425	47	155	61	73	51	534	73	361	63

Table 3 (Cont'd.)

Species Year	Sabine Lake		Galveston		East Matagorda		Matagorda		San Antonio		Aransas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide		
	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	No./ha	Length No./ha	
White shrimp (Cont'd.)																					
1989	615	61	559	55	76	59	384	78	145	52	631	60	372	59	2	68	194	54	356	60	
1990	425	65	1,698	54	690	57	451	63	335	58	821	50	537	67	35	40	368	49	704	55	
1991	385	71	1,723	50	273	51	624	58	236	55	361	71	445	62	77	49	381	61	645	55	
1992	463	68	924	54	264	62	643	60	115	68	211	71	167	66	32	58	85	52	383	58	
1993	324	68	526	56	449	62	585	61	132	68	96	56	876	69	137	58	750	60	437	61	
1994	510	73	985	53	618	55	512	62	327	63	447	64	395	71	55	55	200	59	483	59	
1995	789	70	563	53	613	57	607	60	368	75	218	57	268	70	19	51	378	57	401	60	

<sup>a</sup>Data for October - December only.

<sup>b</sup>East Matagorda Bay data are only for February-September 1983. Coastwide values do not include East Matagorda Bay data.

Table 4. Annual mean catch rate (No./h) and mean total lengths (mm) of select fishes and shellfishes caught with 6.1-m trawls in Texas bay systems during 1982-95. Blank indicates no measurement taken; ND = no data.

Species	Year	East												Corpus Christi				Upper Laguna Madre		Lower Laguna Madre		Coastwide <sup>b</sup>					
		Sabine Lake			Galveston			Matagorda			Matagorda			San Antonio			Arkansas			Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide <sup>b</sup>	
		No./h	Length	No./h Length	No./h	Length	No./h Length	No./h	Length	No./h Length	No./h	Length	No./h Length	No./h	Length	No./h Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h
Atlantic croaker	1982*	ND						102	ND	10	ND	87	75	110	ND	37	ND	28	ND	62	75						
	1983	ND	131	ND	30	117	31	117	ND	18	110	44	106	43	149	15	157	32	154	30	127						
	1984	ND	126	ND	15	104	30	104	87	22	87	52	83	120	121	15	137	44	138	35	112						
	1985	ND	124	ND	20	110	41	110	ND	17	105	33	101	42	138	13	151	24	148	27	119						
	1986	10	157	ND	31	123	ND	52	114	44	105	57	96	83	125	14	139	28	153	43	117						
	1987	25	139	17 <sup>c</sup>	26	117	133	126	103	146	96	87	100	50	129	7	152	44	122	70	106						
	1988	45	135	98	13	131	43	121	90	109	102	100	102	38	125	5	137	21	138	55	109						
	1989	45	145	36	116	4	98	75	120	88	102	71	99	40	127	2	158	19	131	52	115						
	1990	40	113	36	109	12	113	79	118	50	97	45	92	55	125	12	129	66	123	50	112						
	1991	31	115	41	106	8	120	135	106	175	93	223	93	74	125	14	127	34	132	94	103						
	1992	40	139	54	107	4	120	211	100	155	84	238	87	54	114	17	140	37	140	112	98						
	1993	70	131	90	104	15	128	120	104	48	104	123	98	36	131	2	141	27	141	79	106						
	1994	34	144	73	111	17	148	99	116	146	78	55	106	18	135	2	137	27	147	72	106						
	1995	22	117	47	100	24	123	108	103	88	99	87	117	42	124	2	162	51	146	64	106						
	Black drum	1982*	ND					0			<1	221	<1	166	2	235	<1	264	0		<1	238					
		1983	ND	259	ND	<1	274	ND	<1	199	<1	192	<1	201	1	347	1	266	<1	440	<1	283					
1984		ND	168	ND	<1	242	ND	0	0	0	0	<1	251	<1	341	1	202	<1	544	<1	258						
1985		ND	233	ND	<1	246	0 <sup>c</sup>	0	0	0	0	0	403	<1	315	1	280	0		<1	268						
1986		<1	226	ND	<1	246	0	0	0	0	200	0	0	0	186	<1	236	<1	335	<1	250						
1987		<1	278	<1	271	<1	192	<1	170	<1	154	<1	204	<1	299	1	197	<1	160	<1	231						
1988		1	271	<1	274	<1	192	0	0	<1	267	<1	170	170	356	2	212	<1	418	<1	258						
1989		2	260	<1	272	<1	146	<1	930	<1	114	<1	173	<1	560	97	109	<1	169	5	115						
1990		1	272	<1	254	<1	218	0	0	194	<1	194	<1	247	<1	170	71	152	1	229	4	160					
1991		2	268	<1	313	1	235	0	0	212	<1	282	<1	183	<1	359	10	225	1	236	1	236					
1992		2	320	<1	210	<1	309	0	0	212	<1	223	<1	223	<1	379	3	291	<1	357	<1	297					
1993		3	283	<1	275	<1	291	1	259	<1	184	<1	259	<1	401	1	360	<1	408	<1	301						
1994		2	324	<1	291	1	257	0	0	221	<1	221	<1	346	1	325	3	235	0		1	256					
1995		1	306	<1	229	1	257	0	0	24	11	ND	3	ND	1	138	1	193	0		2	141					
Gafftop-sail catfish		1982*	ND					4	ND		3	ND	3	ND	1	175	1	131	0		1	133					
		1983	ND	137	ND	<1	132	1	132	2	123	2	135	2	109	<1	218	<1	196	<1	126	1	126				
	1984	ND	139	ND	<1	144	1	144	5	121	2	128	3	128	1	150	0	<1	210	1	134						
	1985	ND	154	ND	<1	137	2	137	2	128	3	128	3	128	1	150	0	<1	210	1	134						
	1986	0	126	ND	1	126	2	134	2	128	5	128	2	121	<1	92	<1	158	0		1	128					
	1987	<1	174	<1	145	1 <sup>c</sup>	143	2	138	9	122	2	124	<1	132	<1	183	<1	175	2	127						
	1988	0	149	1	135	3	14	3	14	3	131	3	127	<1	14	0		0		1	124						
	1989	<1	126	<1	139	1	139	1	134	4	136	4	139	4	156	<1	156	0		1	137						
	1990	0	218	1	127	1	137	4	136	4	130	2	143	0	173	0	0	0		1	159						
	1991	0	145	1	142	2	145	2	145	5	127	3	141	3	206	0	0	0		2	137						
	1992	<1	144	<1	161	2	128	2	125	5	132	10	117	1	126	0	<1	<1	203	2	127						
1993	0	139	<1	118	2	145	2	145	4	123	4	118	4	183	0	<1	<1	185	2	133							
1994	0	127	<1	197	2	129	2	129	3	119	2	145	2	180	<1	181	0		2	131							
1995	<1	275	2	139	<1	176	1	137	4	129	3	142	1	200	0	<1	<1	207	2	141							
Gulf menhaden	1982*	ND					10	ND		11	ND	24	ND	2	ND	<1	ND	<1	ND	10	ND						
	1983	ND	103	ND	7	103	10	109	17	76	3	89	3	89	3	104	1	87	0		8	96					
	1984	ND	98	ND	3	98	3	93	23	58	45	44	4	82	4	82	6	76	<1	59	9	61					
	1985	ND	112	ND	18	112	10	109	27	79	12	92	2	119	4	106	0		0		14	101					
	1986	<1	121	17	95	ND	4	79	64	18	64	8	55	1	156	<1	92	0		0	9	84					
	1987	3	101	20	95	15 <sup>c</sup>	84	12	101	34	77	22	62	1	128	<1	92	0		0	16	88					

Table 4. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide					
		Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Arkansas	Christi	Upper Laguna Madre	Lower Laguna Madre	Coastwide	No./h	Length	No./h	Length	No./h	Length	No./h	Length				
Gulf menhaden (Cont'd.)	1988	3	94	22	80	1	96	16	96	11	99	4	106	1	124	1	58	<1	110	13	88		
	1989	3	79	14	107	7	97	3	111	21	103	3	65	7	115	<1	60	<1	78	9	105		
	1990	5	68	11	94	2	94	4	121	24	85	19	102	2	97	2	85	<1	111	10	95		
	1991	6	83	21	87	4	82	17	98	34	92	16	88	2	128	1	73	<1	98	17	91		
	1992	2	95	22	103	7	71	31	103	17	94	38	87	3	102	1	108	1	107	20	100		
	1993	2	79	39	84	5	44	10	104	12	68	16	75	4	98	<1	119	<1	136	18	85		
	1994	4	84	30	91	4	46	7	120	13	74	3	117	3	91	<1	187	<1	132	14	93		
	1995	2	68	23	102	2	48	3	105	19	45	6	71	2	131	<1	108	<1	126	11	90		
	Pinfish	1982*	ND	ND	1	ND	ND	7	ND	7	ND	5	ND	2	ND	85	ND	44	ND	39	ND	19	ND
		1983	ND	ND	1	121	ND	6	110	6	110	14	106	38	106	119	124	20	133	45	109	24	119
1984		ND	ND	1	121	ND	6	107	6	107	7	96	39	96	25	113	67	108	73	111	15	107	
1985		ND	ND	1	120	ND	9	111	9	110	23	104	53	110	48	118	18	133	48	110	18	113	
1986		4	117	2	118	ND	10	101	10	101	18	98	55	103	100	116	32	109	95	108	27	109	
1987		<1	126	1	122	5 <sup>c</sup>	113	103	32	91	83	106	106	106	121	12	131	12	131	56	113	32	112
1988		4	126	2	114	5	107	18	111	92	104	139	100	100	272	115	20	112	65	100	59	109	
1989		1	117	2	121	9	98	16	113	53	103	82	103	82	463	117	16	110	81	104	75	114	
1990		3	109	5	107	5	103	34	109	64	101	109	101	109	104	107	104	77	282	101	61	102	
1991		1	111	4	120	8	100	6	116	26	102	32	109	32	109	247	111	81	105	278	52	109	
1992		1	98	2	127	1	112	5	112	10	103	23	101	23	101	159	110	48	117	130	109	31	110
1993		3	119	4	114	3	110	5	103	34	102	91	106	106	136	117	46	105	139	103	37	110	
1994		1	128	9	109	2	125	9	101	27	103	39	118	77	127	127	42	114	97	106	26	115	
1995		<1	122	2	137	2	119	6	106	38	97	69	114	114	138	127	25	115	116	105	34	117	
Red drum		1982*	ND	0	0	0	ND	<1	ND	<1	230	<1	102	<1	102	<1	649	<1	619	0	<1	<1	402
	1983	ND	0	0	0	ND	0	0	<1	319	<1	224	<1	224	0	81	<1	241	<1	280	<1	242	
	1984	ND	<1	583	0	ND	<1	305	<1	344	<1	142	<1	142	<1	276	<1	475	<1	90	<1	292	
	1985	ND	0	0	0	ND	<1	56	0	0	<1	54	<1	78	0	399	<1	630	<1	340	<1	289	
	1986	<1	212	0	0	ND	0	0	0	35	<1	78	<1	78	0	0	<1	399	0	0	<1	154	
	1987	<1	405	<1	34	0 <sup>c</sup>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	<1	72	
	1988	<1	272	<1	53	0	0	0	42	0	0	0	0	0	<1	525	0	40	<1	68	<1	72	
	1989	<1	254	<1	44	0	0	<1	0	0	0	0	0	0	<1	264	<1	383	<1	256	<1	268	
	1990	0	0	<1	320	0	0	0	0	53	<1	0	0	0	<1	117	0	40	<1	342	<1	207	
	1991	0	0	<1	135	0	0	0	0	75	<1	0	0	0	<1	264	<1	383	<1	256	<1	211	
	1992	0	0	<1	197	0	0	<1	63	<1	349	<1	369	<1	117	0	0	415	<1	271	<1	306	
	1993	<1	575	0	433	<1	360	0	0	250	<1	412	<1	412	0	0	<1	415	<1	271	<1	277	
	1994	0	0	<1	433	<1	72	0	0	170	<1	70	<1	70	<1	188	<1	320	<1	382	<1	306	
	1995	<1	246	0	0	0	0	0	0	281	<1	0	0	0	<1	0	<1	320	<1	382	<1	306	
	Sand seatrout	1982*	ND	4	4	4	ND	5	185	5	141	3	126	3	126	14	147	1	201	6	164	5	161
1983		ND	3	3	134	ND	4	132	4	108	3	111	3	111	9	158	<1	196	1	164	3	140	
1984		ND	2	2	147	ND	1	121	1	115	1	107	4	141	4	141	0	0	1	161	1	138	
1985		ND	4	4	127	ND	3	126	3	136	1	119	7	144	7	144	1	160	1	117	3	131	
1986		1	152	3	141	ND	2	117	2	112	<1	133	5	148	5	148	0	0	<1	154	2	137	
1987		2	121	2	110	2	112	5	114	1	99	1	94	1	94	9	134	<1	156	<1	160	3	118
1988		1	140	3	107	1	117	2	126	<1	123	2	107	2	107	3	125	<1	109	<1	128	2	115
1989		2	102	10	96	<1	81	3	111	1	110	4	85	4	85	12	143	0	2	152	6	109	
1990		1	110	5	109	1	96	3	119	1	117	1	113	1	113	3	124	0	2	102	3	113	
1991		1	118	7	130	1	103	2	123	1	119	4	113	4	113	5	143	0	2	140	4	130	
1992		2	113	6	113	<1	150	6	113	2	104	4	128	4	128	2	142	<1	209	2	155	4	117
1993		6	108	6	110	3	107	4	119	1	109	5	103	5	103	5	125	1	146	1	126	4	113
1994		1	76	8	107	3	124	3	119	<1	123	2	130	2	130	3	143	<1	253	2	157	4	114
1995		1	101	8	121	2	104	8	111	1	105	4	126	4	126	3	148	0	0	2	160	5	120

Table 4. (Cont'd.)

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide <sup>b</sup>		
		Sabine Lake	Galveston	Matagorda	Matagorda	Matagorda	San Antonio	Aransas	Christi	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	Upper Laguna Madre	Lower Laguna Madre	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Sheeps-head	1982*	ND	<1	295	ND	0	<1	119	<1	85	<1	345	1	366	1	241	<1	290		
	1983	ND	<1	344	ND	0	<1	113	<1	138	<1	365	1	358	<1	248	<1	323		
	1984	ND	<1	339	ND	<1	147	0	<1	157	<1	342	<1	402	<1	300	<1	314		
	1985	ND	<1	341	ND	<1	102	<1	112	<1	143	<1	259	<1	412	<1	80	<1	242	
	1986	1	<1	451	ND	0	0	0	<1	122	<1	288	<1	356	1	160	<1	228		
	1987	<1	<1	279	0 <sup>c</sup>	<1	111	<1	124	<1	115	<1	299	<1	377	<1	156	<1	255	
	1988	<1	<1	332	0	<1	112	<1	80	<1	95	<1	155	<1	247	<1	152	<1	238	
	1989	1	<1	252	<1	104	<1	120	<1	116	<1	251	<1	518	<1	366	<1	240		
	1990	3	<1	248	0	0	0	<1	89	<1	99	0	0	0	0	<1	234	<1	274	
	1991	2	<1	339	<1	192	0	<1	145	<1	145	<1	229	0	0	<1	136	<1	295	
	1992	3	<1	267	0	<1	65	1	121	<1	149	<1	203	<1	465	<1	187	<1	242	
	1993	5	<1	311	1	286	0	1	134	<1	101	<1	164	<1	473	1	156	<1	227	
	1994	2	<1	281	1	309	0	<1	187	<1	133	<1	165	<1	509	1	225	<1	256	
	1995	3	<1	244	<1	301	<1	123	1	177	<1	158	<1	181	1	161	<1	221		
	Southern flounder	1982*	ND	<1	158	ND	<1	169	1	155	1	186	1	181	2	203	<1	296	1	176
1983		ND	<1	175	ND	<1	196	<1	120	1	180	<1	242	<1	203	<1	161	<1	180	
1984		ND	<1	193	ND	<1	194	<1	153	2	148	<1	175	1	145	<1	168	<1	160	
1985		ND	<1	234	ND	<1	202	1	147	1	152	1	221	1	197	<1	261	<1	191	
1986		<1	1	161	ND	<1	165	1	141	1	144	1	184	1	262	<1	212	1	166	
1987		<1	<1	231	<1 <sup>c</sup>	154	<1	191	<1	160	<1	171	<1	183	<1	181	<1	181		
1988		<1	<1	195	<1	132	<1	148	<1	118	<1	214	<1	226	<1	205	<1	157		
1989		<1	<1	166	<1	181	<1	194	<1	130	<1	193	<1	348	<1	211	<1	168		
1990		<1	<1	174	<1	161	<1	166	<1	121	<1	167	1	190	<1	170	<1	145		
1991		<1	<1	160	<1	147	<1	242	<1	191	<1	228	<1	266	<1	229	<1	180		
1992		<1	<1	184	<1	186	<1	210	<1	135	0	0	<1	0	<1	205	<1	188		
1993		<1	<1	155	<1	177	<1	142	<1	126	<1	232	<1	416	<1	391	<1	154		
1994		<1	<1	160	<1	230	<1	162	<1	226	<1	323	<1	236	<1	223	<1	186		
1995		1	<1	192	0	<1	224	1	170	<1	220	<1	111	<1	332	0	<1	<1		
Spotted seatrout		1982*	ND	9	ND	ND	26	ND	5	112	68	ND	33	ND	10	ND	4	ND	19	ND
	1983	ND	6	120	ND	17	122	5	112	18	118	36	140	2	163	6	135	12	127	
	1984	ND	8	115	ND	34	107	35	84	131	91	74	112	82	118	10	108	39	103	
	1985	ND	13	121	ND	20	118	13	110	60	116	215	132	24	137	19	129	41	126	
	1986	6	120	14	120	ND	29	121	21	99	92	115	129	6	118	5	135	35	119	
	1987	9	134	11	127	12	119	38	115	34	86	117	122	125	4	158	13	112	37	
	1988	24	113	14	117	5	107	42	127	116	108	235	127	4	140	18	118	66	120	
	1989	19	130	11	123	6	111	85	118	73	105	240	136	6	129	18	119	68	125	
	1990	6	130	8	117	12	95	44	119	117	96	164	113	71	110	104	104	78	109	
	1991	6	124	9	120	6	108	94	124	39	105	206	116	24	130	82	117	50	116	
	1992	10	137	19	125	2	125	71	128	25	119	66	130	9	149	25	133	40	123	
	1993	32	119	16	135	4	131	86	112	30	101	167	118	4	134	11	134	54	115	
	1994	25	129	24	116	4	128	23	122	39	103	56	135	1	101	9	137	30	120	
	1995	6	127	17	127	4	119	38	119	53	110	41	146	1	141	15	139	32	124	
	Spotted seatrout	1982*	ND	<1	173	ND	0	ND	<1	232	<1	163	<1	187	1	166	<1	142	<1	171
1983		ND	<1	288	ND	<1	155	<1	168	2	207	<1	327	2	188	<1	200	<1	212	
1984		ND	<1	418	ND	<1	174	<1	252	<1	237	<1	385	<1	351	<1	236	<1	329	
1985		ND	<1	286	ND	<1	171	<1	156	1	156	<1	171	1	146	<1	218	<1	188	
1986		<1	<1	259	ND	<1	193	<1	170	<1	162	1	176	<1	151	1	196	<1	201	
1987		<1	<1	134	<1 <sup>c</sup>	162	<1	143	1	166	1	164	<1	163	1	206	<1	198	<1	167
1988		<1	<1	189	<1	166	<1	249	<1	159	2	166	<1	175	<1	176	<1	95	<1	172
1989		<1	<1	142	<1	128	<1	174	<1	190	1	168	<1	214	1	186	1	139	<1	173
1990		<1	<1	118	0	<1	128	<1	119	<1	176	<1	123	<1	114	0	139	<1	150	
1991		<1	<1	165	<1	184	<1	134	<1	136	1	154	<1	161	1	124	1	177	<1	155



Table 4. (Cont'd.)

Species	Year	Sabine Lake		Galveston		East		San Antonio		Arkansas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Spotted seatrout (Cont'd.)	1992	<1	194	<1	155	<1	150	<1	149	2	182	<1	219	2	175	1	185	1	167	
	1993	<1	196	<1	133	<1	130	<1	149	1	181	<1	239	2	174	<1	216	<1	170	
	1994	<1	142	<1	0	<1	220	<1	127	<1	199	<1	235	1	175	<1	247	<1	159	
	1995	<1	172	<1	182	<1	190	<1	156	1	193	<1	218	1	167	1	157	<1	168	
	1982*	ND		<1		ND			1	2		2	212	1	311	<1		1	232	
Striped mullet	1983	ND		<1	131	<1	131	2	137	3	209	1	211	2	323	1	331	1	210	
	1984	ND		<1	204	<1	204	<1	174	1	192	1	209	6	287	1	307	1	250	
	1985	ND		<1	163	<1	163	<1	136	7	158	<1	168	1	243	<1	254	2	181	
	1986	<1	187	<1	116	<1	116	<1	157	<1	158	1	226	<1	278	<1	266	1	250	
	1987	1	168	<1	158	<1	158	4	145	1	171	1	192	0		0		1	210	
	1988	2	239	<1	138	<1	138	1	130	<1	156	3	185	<1	334	0		0	1	243
	1989	5	183	<1	237	<1	237	1	188	<1	187	2	206	0		<1	365	2	234	
	1990	<1	234	<1	141	<1	141	<1	136	1	155	<1	239	<1	292	0		0	<1	180
	1991	4	174	<1	114	<1	114	3	145	2	141	<1	216	2	279	<1	276	2	181	
	1992	6	232	0		<1	129	3	145	4	143	1	213	<1	215	<1	178	3	206	
	1993	1	209	<1	157	<1	157	1	148	1	168	1	219	<1	334	<1	244	1	214	
	1994	4	261	<1	141	<1	141	1	144	3	184	1	260	<1	354	<1	240	1	204	
	1995	1	190	<1	333	0		<1	134	2	191	3	196	0		<1	220	2	234	
	Total finfishes	1982*	ND		193	139	ND	193	48	179	270	119	371	166	313	232	152	183	171	167
		1983	ND		162	99	ND	162	107	93	174	108	308	139	170	115	143	139	139	116
	1984	ND		111	104	ND	111	104	82	312	86	294	124	197	123	169	130	134	108	
	1985	ND		115	114	ND	115	96	101	236	99	380	129	96	127	149	128	143	117	
	1986	28	151	127	112	ND	127	118	97	261	104	378	132	86	109	188	132	151	117	
	1987	53	136	121	64 <sup>c</sup>	117	122	302	100	354	101	370	131	64	117	157	126	200	112	
	1988	101	131	138	101	49	122	302	100	354	101	370	131	64	117	157	126	200	112	
	1989	98	137	111	119	44	105	295	106	347	109	857	133	53	103	197	121	272	122	
	1990	85	122	94	116	41	108	304	102	381	106	464	123	368	88	564	119	259	113	
	1991	72	127	176	106	41	109	347	97	423	102	614	122	208	125	524	123	318	109	
	1992	94	152	166	121	23	102	268	98	443	97	335	121	106	130	305	129	281	111	
	1993	156	142	201	107	55	132	297	108	365	110	433	131	87	113	269	128	252	114	
	1994	82	152	194	111	54	150	295	92	266	104	268	123	85	106	203	130	218	110	
	1995	59	128	216	103	69	130	329	101	338	120	331	134	53	122	278	132	273	109	
SHELLFISHES																				
Blue crab	1982*	ND		28	91	ND	28	17	81	29	66	7	97	9	148	10	100	17	89	
	1983	ND		24	88	ND	24	21	80	40	81	2	96	7	113	12	97	18	86	
	1984	ND		19	92	ND	19	8	82	31	81	8	88	24	106	50	86	15	90	
	1985	ND		30	79	ND	30	19	76	23	72	5	115	21	103	36	86	21	81	
	1986	6	132	28	79	ND	28	13	85	25	78	14	88	8	100	15	85	19	83	
	1987	5	135	19	78	28 <sup>c</sup>	87	40	93	18	84	6	95	8	108	19	88	17	86	
	1988	5	137	9	71	13	91	89	75	57	63	7	88	7	98	18	84	22	74	
	1989	9	135	25	66	51	63	50	74	24	68	2	94	2	107	9	77	19	72	
	1990	6	98	31	72	15	79	69	69	17	71	14	96	5	93	33	91	21	76	
	1991	7	117	10	64	26	76	6	58	51	58	7	102	5	105	35	89	20	65	
1992	7	139	8	77	2	102	6	105	54	38	56	10	81	26	110	27	98	24	65	
1993	5	131	16	70	6	93	14	82	35	78	10	96	10	114	22	88	20	81		
1994	4	146	16	74	3	90	23	95	71	26	72	3	66	20	83	25	93	24	67	
1995	2	133	8	58	3	111	8	25	55	11	67	4	69	11	76	17	84	10	64	

Table 4. (Cont'd.)

Species	Year	Sabine Lake		Galveston		East Matagorda		San Antonio		Arkansas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide <sup>b</sup>				
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length			
Brown shrimp	1982*	ND		23	90	ND		25	94	17	101	54	80	40	90	40	101	6	61	27	91	
	1983	ND		12	99	ND		26	100	31	99	56	91	8	99	8	102	9	66	21	97	
	1984	ND		13	102	ND		7	102	58	96	107	80	50	103	25	108	6	74	30	94	
	1985	ND		33	75	ND		24	89	27	90	67	81	24	96	16	108	11	63	30	83	
	1986	<1		15	94	ND		29	99	69	98	111	96	42	95	7	108	15	64	34	96	
	1987	4		92	24	7 <sup>c</sup>		47	91	93	85	101	88	66	94	8	100	5	70	46	89	
	1988	3		85	24	84	10	91	32	100	124	91	139	86	17	89	6	93	3	73	44	90
	1989	8		84	29	84	47	97	39	91	156	90	105	90	17	88	5	92	9	63	49	89
	1990	1		113	11	98	40	100	26	96	104	92	78	90	28	88	12	91	27	79	34	92
	1991	1		93	13	87	63	96	21	86	51	89	158	91	29	91	19	97	8	80	32	90
	1992	3		83	38	82	9	90	23	82	64	81	64	81	30	92	40	110	7	73	37	84
	1993	9		79	18	85	14	69	43	94	45	82	95	88	22	87	13	103	5	67	32	88
	1994	9		83	29	99	3	69	51	95	101	88	37	85	10	88	18	103	54	58	41	91
	1995	1		91	12	83	9	65	54	84	100	75	108	84	22	90	15	91	14	70	41	81
	Pink shrimp	1982*	ND		<1	94	ND		<1	113	<1	96	7	89	2	100	1	96	0		1	94
		1983	ND		<1	95	ND		1	112	5	95	9	94	2	103	1	113	1	88	2	99
1984		ND		0		ND		<1	76	<1	72	3	86	3	109	<1	94	<1	71	1	98	
1985		ND		<1	88	ND		<1	104	3	98	4	100	5	96	4	107	1	98	2	99	
1986		0		<1	118	ND		2	114	4	103	11	101	10	103	1	109	<1	70	3	104	
1987		0		<1	111	2 <sup>c</sup>		5	95	2	92	6	84	12	101	1	107	2	72	3	95	
1988		0		1	79	<1		2	89	6	86	20	82	8	93	<1	76	2	77	4	85	
1989		0		<1	90	<1		1	102	8	91	14	91	8	95	<1	85	1	80	3	93	
1990		0		<1	84	0		<1	106	1	97	23	88	4	97	3	71	3	85	3	90	
1991		0		<1	101	1		1	115	2	102	8	84	27	88	8	103	4	79	5	90	
1992		0		<1	58	<1		<1	87	<1	70	7	77	10	95	9	103	20	82	3	89	
1993		0		<1	87	0		<1	100	<1	86	5	76	4	91	1	98	4	79	1	85	
1994		0		<1	92	<1		3	104	5	78	6	85	5	89	4	93	15	62	3	84	
1995		0		<1	88	2		80	10	97	6	82	4	90	13	100	4	95	7	70	5	94
White shrimp		1982*	ND		88	93	ND		39	86	14	99	16	95	26	101	17	110	4	61	46	92
		1983	ND		78	93	ND		20	102	13	96	18	100	14	111	6	112	2	86	36	95
	1984	ND		60	98	ND		15	99	8	99	38	106	24	106	11	126	10	109	32	101	
	1985	ND		62	99	ND		21	110	23	91	17	106	22	104	6	120	1	105	33	101	
	1986	14	105	45	95	ND		60	98	15	96	13	101	19	98	3	108	5	57	34	97	
	1987	23	101	37	97	22 <sup>c</sup>		16	97	42	87	10	94	15	99	2	105	2	76	24	95	
	1988	39	107	21	91	8		16	98	41	93	16	91	12	95	3	102	<1	79	20	94	
	1989	29	87	29	89	11		9	98	43	99	7	98	9	100	3	97	<1	114	20	93	
	1990	50	90	14	98	14		16	115	47	97	13	108	22	98	21	100	1	113	21	100	
	1991	17	91	76	97	7		11	95	27	94	30	89	24	121	14	113	1	107	37	98	
	1992	37	88	59	93	5		31	96	24	95	53	93	5	111	6	114	1	104	35	94	
	1993	11	81	38	91	31		17	97	18	88	21	95	10	90	14	96	2	97	23	92	
	1994	45	96	95	80	15		9	107	44	87	6	101	34	91	10	109	2	94	45	84	
	1995	4	93	55	90	34		87	11	101	28	86	9	98	6	103	7	104	8	93	26	92

<sup>a</sup>Values include May-Dec only.<sup>b</sup>1986 values include Sabine Lake; 1987 values include East Matagorda.<sup>c</sup>Values include Apr-Dec only.

Table 5. Annual mean catch rates (No./h) and mean total lengths (mm) of select fishes and shellfishes caught with 6.1-m trawls in the Texas Territorial Sea during 1985-95. Blank indicates no measurement taken; ND = no data.

Species	Year	Sabine		Galveston		Port O'Connor		Port Aransas		Port Isabel		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
<b>FINFISHES</b>														
Atlantic croaker	1985 <sup>a</sup>	ND		22	145	42	139	17	145	9	149	23	142	
	1986	44 <sup>b</sup>	134	45	126	98	136	43	130	9	132	49	132	
	1987	9	114	110	119	65	131	28	134	<1	157	44	124	
	1988	79	122	78	118	122	132	23	130	2	128	55	125	
	1989	64	115	117	117	75	128	28	128	6	137	60	121	
	1990	175	117	139	111	69	135	65	131	4	119	91	119	
	1991	272	111	153	114	201	121	87	129	4	162	145	117	
	1992	229	110	228	116	153	116	81	106	6	126	142	113	
	1993	437	111	200	110	74	123	91	121	10	144	162	113	
	1994	140	115	109	126	75	141	2	114	14	143	69	126	
	1995	80	116	33	123	112	119	20	133	1	121	50	119	
	Black drum	1985 <sup>a</sup>	ND <sup>b</sup>		0		0		<1	825	0		<1	825
		1986	0		0		<1	900	0		0		<1	900
		1987	<1	851	<1	760	<1	680	<1	680	0		<1	741
		1988	0		<1	752	0		0		0		<1	752
		1989	<1	698	0		<1	506	0		0		<1	631
1990		0		<1	528	0		0		0		<1	538	
1991		0		<1	970	0		0		0		<1	970	
1992		0		0		<1	889	0		0		<1	889	
1993		<1	146	<1	825	0		0		0	780	<1	632	
1994		0		<1	843	0		0		0		<1	843	
1995	<1	871	<1	142	0		0		0		<1	495		
Gafftopsail catfish	1985 <sup>a</sup>	ND <sup>b</sup>		<1	165	<1	156	<1	136	0		<1	160	
	1986	13	121	<1	118	<1	115	<1	176	0		3	121	
	1987	3	116	0		<1	158	<1	134	0		1	118	
	1988	2	118	<1	169	<1	168	0		<1	180	<1	126	
	1989	2	144	1	123	<1	546	<1	187	0		<1	143	
	1990	3	119	<1	123	0		0		0		1	119	
	1991	1	145	<1	170	<1	181	<1	178	0		<1	150	
	1992	12	125	1	148	<1	148	<1	209	0		3	127	
	1993	6	123	<1	129	<1	182	<1	145	0		1	127	
	1994	6	131	2	152	<1	239	1	204	0		2	143	
	1995	5	131	1	141	0		1	155	0		1	135	
Gulf menhaden	1985 <sup>a</sup>	ND <sup>b</sup>		2	150	1	159	1	151	0		1	152	
	1986	4	125	2	147	<1	180	<1	197	0		1	135	
	1987	3	132	5	135	1	146	<1	159	0		2	136	
	1988	5	124	10	57	6	107	<1	122	0		87	87	
	1989	1	137	1	144	<1	131	<1	177	<1	51	1	138	
	1990	2	133	4	136	1	122	<1	162	0		1	134	
	1991	7	134	1	144	1	130	<1	148	0		2	135	
	1992	4	141	14	116	1	139	1	145	0		4	123	
	1993	5	142	1	129	<1	159	0		0		1	141	
	1994	6	131	3	132	3	117	<1	116	<1	157	2	129	
1995	5	137	4	137	1	159	<1	163	<1	131	<1	140		

Table 5. (Cont'd.)

Species	Year	Sabine		Galveston		Port O'Connor		Port Aransas		Port Isabel		Coastwide	
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
King mackerel	1985 <sup>a</sup>	ND <sup>b</sup>		<1	173	0		<1	124	0		<1	142
	1986	0		<1	159	0		0		0		<1	159
	1987	0		0		<1	120	<1	200	0		<1	131
	1988	0		0		0		0		0		0	
	1989	0		0		<1	161	<1	164	0		<1	162
	1990	0		<1	201	<1	223	0		0		<1	210
	1991	0		<1	172	<1	157	<1	99	0		<1	132
	1992	0		<1	149	<1	152	1	136	<1	192	<1	144
	1993	0		0		0		<1	169	0		<1	169
	1994	0		0		0		<1	167	0		<1	167
	1995	0		0		0	70	<1	88	<1	89	<1	84
	Pinfish	1985 <sup>a</sup>	ND <sup>b</sup>		<1	124	3	109	4	110	1	135	2
1986		<1	98	2	104	2	105	4	107	2	103	2	105
1987		0		<1	100	3	111	3	115	<1	112	1	113
1988		<1	93	<1	112	8	105	8	110	3	105	4	107
1989		<1	100	1	108	3	116	7	110	6	105	3	109
1990		<1	86	1	111	4	110	18	105	2	98	5	105
1991		<1	121	1	132	2	116	18	113	2	118	4	114
1992		<1	115	2	121	3	110	6	103	3	107	3	108
1993		<1	72	<1	102	3	105	6	110	2	111	3	108
1994		<1	131	1	111	5	107	4	107	6	107	3	107
1995		0		1	117	3	101	4	121	7	115	3	114
Red drum		1985 <sup>a</sup>	ND <sup>b</sup>		0		0		<1	84	0		<1
	1986	0		0		0		0		0		0	
	1987	0		0		<1	948	0		<1	42	<1	520
	1988	0		0		0		0		0		0	
	1989	0		<1	1,110	0		0		0		<1	1,110
	1990	0		<1	61	0		0		0		<1	61
	1991	0		0		0		0		0		0	
	1992	0		0		0		0		<1	95	<1	95
	1993	0		<1	1,013	0		0		0		<1	1,013
	1994	0		0		0		0		0		0	
	1995	0		0		0	<1	811	<1	1,037	0	<1	922
	Red snapper	1985 <sup>a</sup>	ND <sup>b</sup>		0		0		2	85	7	89	2
1986		0		0		<1	152	1	95	<1	103	<1	100
1987		0		0	68	<1	88	1	122	<1	83	<1	107
1988		0		0		0		1	111	1	106	<1	109
1989		0		<1	74	2	87	4	87	3	90	2	88
1990		0		0		<1	94	3	105	2	113	1	106
1991		0		0		0		9	80	2	106	2	84
1992		0		0		0	79	6	77	2	99	2	81
1993		0		<1	126	1	76	2	77	3	98	1	88
1994		0		0		3	89	3	103	5	97	2	96
1995		0		0		4	70	6	88	8	89	4	84

Table 5. (Cont'd.)

Species	Year	Sabine		Galveston		Port O'Connor		Port Aransas		Port Isabel		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Sand seatrout	1985 <sup>a</sup>	ND <sub>b</sub>		10	141	6	168	3	140	<1	221	5	150	
	1986	5	164	4	141	3	151	1	174	0		3	154	
	1987	7	131	6	133	5	134	2	162	<1	108	4	135	
	1988	3	148	5	114	11	129	1	184	<1	137	4	130	
	1989	22	133	41	110	16	127	7	155	2	123	18	122	
	1990	50	136	8	126	7	139	2	130	1	118	14	135	
	1991	28	130	12	143	7	146	12	129	1	153	12	135	
	1992	41	132	11	138	6	148	5	131	<1	161	13	135	
	1993	45	129	7	131	15	116	10	112	2	121	16	124	
	1994	82	132	3	149	5	148	2	125	1	130	18	134	
	1995	23	126	18	129	111	106	16	125	6	88	36	112	
	Southern flounder	1985 <sup>a</sup>	ND <sub>b</sub>		0		<1	280	<1	137	0		<1	199
		1986	1	162	<1	255	<1	184	<1	311	0		<1	173
		1987	<1	256	<1	197	0		<1	179	<1	168	<1	191
		1988	<1	204	0		<1	214	<1	225	0		<1	214
1989		0		0		<1	210	<1	298	0		<1	239	
1990		<1	187	0		<1	212	<1	164	<1	250	<1	197	
1991		<1	286	<1	260	<1	194	<1	188	0		<1	220	
1992		<1	143	<1	240	0		<1	284	<1	418	<1	270	
1993		<1	124	0				<1	279	<1		<1	201	
1994		<1	171	<1	180	<1	215	0		<1	286	<1	205	
1995		0		0		<1	262	0		0		<1	262	
Spanish mackerel		1985 <sup>a</sup>	ND <sub>b</sub>		0		0		0		0		0	
		1986	<1	200	0		0		0		0		<1	200
		1987	<1	93	<1	183	0		<1	258	0		<1	203
		1988	<1	166	<1	178	<1	182	<1	110	<1	200	<1	180
	1989	<1	206	<1	172	<1	175	<1	175	0		<1	182	
	1990	<1	174	1	176	<1	225	<1	192	0		<1	180	
	1991	1	184	1	163	<1	144	<1	134	0		<1	168	
	1992	<1	158	<1	175	<1	181	<1	164	<1		<1	168	
	1993	1	167	<1	188	0		<1	237	0		<1	190	
	1994	0		0		<1	170	<1	170	0		<1	170	
	1995	<1	194	<1	186	<1	135	<1	242	0		<1	192	
	Spot	1985 <sup>a</sup>	ND <sub>b</sub>		3	132	20	130	21	141	1	142	11	136
		1986	3	124	8	128	7	124	25	123	2	125	9	124
		1987	5	140	9	126	4	125	22	129	<1	170	8	129
		1988	4	115	7	116	23	128	23	122	3	110	12	123
1989		6	120	27	108	18	124	48	121	4	121	21	118	
1990		9	123	25	121	102	125	93	117	4	112	47	125	
1991		18	117	4	125	67	122	37	127	1	129	26	123	
1992		5	127	12	126	6	122	10	126	2	117	7	125	
1993		4	122	14	119	4	126	19	125	4	138	9	124	
1994		13	125	4	131	13	125	4	131	4	129	8	127	
1995		7	115	3	135	30	132	7	150	4	125	10	132	

Table 5. (Cont'd.)

Species	Year	Sabine		Galveston		Port O'Connor		Port Aransas		Port Isabel		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Spotted seatrout	1985 <sup>a</sup>	ND <sup>b</sup>		0		0		<1	140	0		<1	140	
	1986	<1 <sup>b</sup>	163	<1	172	<1	165	0		0		<1	165	
	1987	<1	178	0		0		0		0		<1	178	
	1988	0		<1	65	<1	110	0		0		<1	88	
	1989	<1	98	0		<1	173	0		0		<1	137	
	1990	<1	110	<1	160	<1	122	<1	144	0		<1	132	
	1991	0		0		<1	148	0		0		<1	148	
	1992	<1	112	0		0		0		0		<1	112	
	1993	0		0		<1	160	0		0		<1	160	
	1994	<1	187	<1	54	0		0		0		<1	67	
	1995	<1	226	0		0		0		0		<1	226	
	Total finfishes	1985 <sup>a</sup>	ND <sup>b</sup>		148	119	188	118	227	114	130	101	174	114
		1986	159 <sup>b</sup>	122	207	118	215	123	292	119	72	110	190	120
1987		158	98	289	111	229	118	226	114	80	96	199	110	
1988		153	120	273	104	379	114	291	106	52	103	234	110	
1989		178	114	301	111	350	118	354	113	106	108	261	114	
1990		477	121	355	113	464	138	337	115	80	103	346	122	
1991		427	117	322	125	666	115	458	108	124	102	404	115	
1992		524	115	499	116	523	111	332	103	128	96	406	111	
1993		651	117	324	116	376	102	381	104	135	106	377	110	
1994		408	121	253	121	560	110	447	99	151	105	367	111	
1995		231	119	165	109	900	98	394	109	174	97	380	104	
Blue crab		1985 <sup>a</sup>	ND <sup>b</sup>		<1	105	1	134	1	127	<1	144	<1	127
		1986	4	96	6	105	1	141	1	145	1	123	3	110
	1987	3	96	1	112	2	105	<1	142	<1	140	1	106	
	1988	2	85	<1	104	1	113	1	128	<1	160	1	105	
	1989	4	61	2	72	1	130	<1	134	<1	146	1	78	
	1990	15	80	4	63	1	118	1	126	1	127	4	84	
	1991	19	72	6	58	1	102	2	114	<1	121	6	73	
	1992	7	58	1	104	<1	85	1	95	<1	123	2	69	
	1993	5	78	1	83	2	116	1	130	1	102	2	95	
	1994	9	77	2	123	1	115	2	66	1	128	3	87	
	1995	8	65	1	61	<1	120	1	122	<1	122	2	70	
	Brown shrimp	1985 <sup>a</sup>	ND <sup>b</sup>		7	103	7	125	47	109	18	106	19	109
		1986	10	107	13	99	6	114	10	105	6	110	9	105
1987		7	104	24	104	9	108	14	106	1	118	11	106	
1988		15	102	5	109	24	103	28	106	<1	116	15	104	
1989		33	103	50	96	56	105	140	95	12	98	59	98	
1990		34	101	10	108	55	107	58	114	20	106	36	108	
1991		12	90	2	102	12	93	9	101	17	123	10	104	
1992		9	91	20	103	4	96	19	92	2	115	11	97	
1993		23	100	21	97	13	105	9	97	4	109	14	100	
1994		6	100	10	101	5	99	16	94	7	106	9	99	
1995		49	102	5	97	49	101	31	94	2	112	28	100	

## SHELLFISHES

Table 5. (Cont'd.)

Species	Year	Sabine		Galveston		Port O'Connor		Port Aransas		Port Isabel		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Pink shrimp	1985 <sup>a</sup>	ND <sup>b</sup>		<1	120	<1	130	1	119	1	108	1	116	
	1986	0		<1	124	2	110	4	105	3	118	2	111	
	1987	0		0		1	114	5	102	1	124	1	108	
	1988	<1	87	0		1	108	7	103	1	125	2	106	
	1989	0		<1	105	1	103	7	100	4	117	2	105	
	1990	0		<1	104	1	101	3	118	3	117	1	114	
	1991	<1	101	<1	99	1	109	6	112	2	118	2	112	
	1992	<1	88	<1	79	<1	114	4	102	<1	122	1	104	
	1993	0		<1	104	4	99	5	104	9	112	4	107	
	1994	<1	90	<1	116	1	109	10	98	8	116	4	106	
	1995	<1	78	0		6	102	6	109	3	112	3	106	
	White shrimp	1985 <sup>a</sup>	ND <sup>b</sup>		53	110	26	124	11	126	1	105	24	115
		1986	41	101	53	101	15	120	8	124	2	137	24	105
		1987	26	105	14	109	16	112	8	119	1	121	13	110
		1988	14	105	17	100	19	110	9	116	<1	133	12	107
1989		21	102	25	106	22	108	14	113	1	122	17	107	
1990		18	104	11	115	15	118	6	136	2	136	10	115	
1991		28	105	10	117	30	106	6	127	1	122	15	109	
1992		51	98	31	108	11	112	10	118	1	145	21	105	
1993		61	101	10	108	11	121	5	134	1	133	17	106	
1994		17	109	8	109	15	114	9	116	1	128	10	112	
1995		10	110	20	113	14	117	12	125	3	134	12	117	

<sup>a</sup> Values include Feb-Dec only off Port Aransas and Aug-Dec only off all other areas.

<sup>b</sup> Values include Jun-Dec only.

Table 6. Annual mean catch rates (No./h) and mean total lengths (mm) by size class<sup>a</sup> of Eastern oyster caught with 46.0-cm wide dredges on "reef" stations in Texas bay systems during 1984-95. Blank indicates no measurement taken; ND = no data.

Size class	Year	Galveston		Matagorda		San Antonio		Arkansas		Coastwide		
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	
Spat	1984	491		ND		ND		ND		491		
	1985	891		ND		ND		ND		891		
	1986	1,010	764	654	66	439	1,864	3,071	1,685	1,022	1,440	
	1987	1,054	654	938	2,019	1,117	894	410	860	1,194		
	1988	1,440	938	2,019	1,117	894	410	860	1,194			
	1989	1,322	1,289	718	1,458	1,991	1,458	82	0	1,440		
	1990	2,147	1,289	718	1,458	1,991	1,458	82	0	1,440		
	1991	1,458	1,289	718	1,458	1,991	1,458	82	0	1,440		
	1992	3,083	454	268	122	546	719	1,836				
	1993	3,194	139	122	546	719	1,836					
	1994	1,263	329	546	719	1,836						
	1995	718	1,311	1,493								
	Small	1984	1,705	47	ND		ND		ND		1,705	47
		1985	2,096	54	ND		ND		ND		2,095	54
		1986	1,316	54	382	51	565	58	1,273	51	1,001	54
		1987	1,070	51	555	51	240	55	2,499	50	1,077	51
		1988	1,500	53	580	52	235	42	2,187	52	1,208	52
1989		1,086	47	706	48	1,995	50	2,278	49	1,463	48	
1990		2,196	45	417	48	1,401	53	1,495	45	1,971	46	
1991		4,927	48	1,040	51	538	54	1,016	48	2,615	49	
1992		4,601	51	622	52	92	48	263	54	2,168	51	
1993		3,895	54	396	54	500	51	296	59	1,926	54	
1994		3,002	52	805	48	573	47	1,010	46	1,749	50	
1995		2,656	53	1,193	49	987	52	4,192	52	2,354	52	
Market		1984	447	91	ND		ND		ND		447	91
		1985	674	88	ND		ND		ND		674	88
		1986	617	88	212	92	444	92	191	86	438	89
		1987	370	91	167	91	258	93	411	86	323	90
		1988	397	89	201	91	23	89	402	87	284	88
	1989	232	90	177	90	414	90	282	85	275	89	
	1990	179	88	114	89	445	88	99	83	215	88	
	1991	502	87	216	89	377	91	65	84	349	88	
	1992	796	87	164	88	24	93	40	83	384	87	
	1993	1,346	88	204	92	74	87	161	87	664	87	
	1994	1,214	90	313	95	287	93	355	93	708	91	
	1995	760	89	453	92	415	93	1,056	92	690	91	

<sup>a</sup> Spat (5-25 mm), small (26-75 mm), market ( $\geq 76$  mm). Mean total length not calculated for spat.



Table 7. Seasonal (May-Nov) mean catch rates (No./ha) and mean total lengths (mm) of select fishes and shellfishes caught with 60.9-m beach seines in 5 Texas gulf shoreline areas during 1987-95. Blank indicates no measurement taken; ND = no data.

Species	Year	Gulf-17		Gulf-18		Gulf-19		Gulf-20		Gulf-21		Coastwide		
		No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	
<b>PINFISHES</b>														
Atlantic croaker	1987 <sup>a</sup>	2	267	<1	306	<1	239	0	0	0	0	0	1	267
	1988	1	264	1	252	<1	260	0	<1	292	<1	<1	<1	262
	1989	2	257	<1	263	<1	205	0	0	0	0	<1	<1	255
	1990	1	260	<1	250	0	0	0	<1	230	<1	<1	<1	259
	1991	2	257	<1	224	<1	251	<1	238	0	0	<1	<1	256
	1992	<1	307	<1	233	<1	255	0	0	0	0	<1	<1	264
	1993	1	255	0	0	<1	289	<1	286	1	290	<1	<1	270
	1994	2	238	0	0	0	0	0	<1	229	<1	<1	<1	238
	1995	2	256	<1	277	<1	263	0	0	<1	283	<1	<1	258
	Black drum	1987 <sup>d</sup>	1	344	<1	215	1	287	<1	249	<1	236	1	293
		1988	1	240	1	226	1	281	<1	272	0	0	1	253
		1989	1	286	4	262	2	249	1	236	<1	216	2	256
1990		2	318	2	243	2	300	2	276	1	280	2	292	
1991		3	264	3	231	1	257	11	240	1	233	3	245	
1992		1	258	3	254	2	305	2	287	<1	340	2	286	
1993		1	334	2	303	1	354	1	340	1	394	1	339	
1994		2	257	1	240	1	463	1	416	0	0	1	333	
1995		3	284	2	265	1	291	1	318	1	240	1	281	
Gulf menhaden		1987 <sup>d</sup>	0	0	0	0	0	0	0	0	0	0	0	0
		1988	7	158	1	166	<1	197	<1	197	<1	226	2	159
		1989	0	0	<1	158	<1	63	0	0	0	0	<1	69
	1990	0	0	<1	214	0	0	<1	237	<1	234	<1	232	
	1991	0	0	<1	211	<1	187	<1	213	0	0	<1	206	
	1992	0	0	0	197	0	0	0	0	0	0	<1	197	
	1993	0	0	<1	209	<1	161	0	0	0	0	<1	198	
	1994	<1	253	1	236	0	42	0	0	0	0	<1	221	
	1995	7	190	<1	261	0	0	0	0	0	0	<1	190	
	Red drum	1987 <sup>d</sup>	0	0	0	0	1	337	<1	340	<1	345	<1	338
		1988	<1	460	<1	324	<1	528	<1	305	<1	702	<1	459
		1989	<1	552	<1	501	<1	370	<1	547	<1	352	<1	485
1990		0	0	<1	320	<1	391	<1	344	<1	356	<1	384	
1991		4	321	1	317	2	318	2	318	<1	375	2	320	
1992		<1	436	<1	496	1	415	<1	395	<1	365	1	417	
1993		<1	438	<1	337	<1	498	1	330	<1	330	<1	397	
1994		<1	652	<1	281	1	431	<1	500	<1	267	<1	448	
1995		1	447	<1	384	<1	476	<1	356	<1	338	<1	442	
Sand seatrout		1987 <sup>d</sup>	1	328	0	0	0	0	0	0	0	0	<1	328
		1988	<1	322	<1	276	<1	298	0	0	<1	286	<1	297
		1989	0	0	<1	353	0	0	0	0	0	0	<1	353
	1990	<1	291	<1	284	0	287	0	0	0	0	<1	287	
	1991	0	0	<1	251	<1	319	0	0	<1	0	<1	307	
	1992	0	0	<1	301	0	0	0	0	0	0	<1	301	

Table 7. (Cont'd.)

Species	Year	Gulf-17		Gulf-18		Gulf-19		Gulf-20		Gulf-21		Coastwide	
		No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length
Sand seatrout (Cont'd.)	1993	0		0		0		<1	360	0		<1	360
	1994	<1	332	<1	279	0		<1	215	<1	65	<1	262
	1995	0		<1	286	<1	356	0		0		<1	344
Sheepshead	1987 <sup>d</sup>	0		0		0		0		0		0	
	1988	<1	416	<1	445	<1	292	<1	288	0		<1	366
	1989	0		0		0		<1	370	0		<1	370
	1990	0		<1	375	<1	312	<1	322	<1	298	<1	344
	1991	0		<1	270	<1	328	0		<1	460	<1	314
	1992	0		<1	458	<1	327	0		<1	441	<1	382
	1993	0		<1	361	<1	413	0		0		<1	372
	1994	0		<1	365	0		0		<1	275	<1	341
	1995	0		<1	352	<1	241	<1	319	<1	319	<1	287
	Southern flounder	1987 <sup>a</sup>	0		1	250	0		<1	313	0		<1
1988		<1	279	1	261	<1	203	<1	207	<1	434	<1	265
1989		<1	375	<1	276	0		<1	270	0		<1	319
1990		<1	264	1	220	<1	226	<1	193	<1	217	<1	231
1991		<1	308	1	267	<1	267	<1	265	0		<1	279
1992		<1	465	<1	270	<1	307	<1	309	<1	192	<1	303
1993		<1	381	1	338	<1	324	0		<1	177	<1	347
1994		<1	274	<1	226	<1	371	<1	357	0		<1	302
1995		<1	282	1	268	<1	384	0		<1	494	<1	287
Spanish mackerel		1987 <sup>d</sup>	0		0		0		0		0		0
	1988	0		0		0		<1	392	<1	392	<1	392
	1989	0		<1	606	0		0		0		<1	606
	1990	0		0		<1	415	<1	477	<1	521	<1	486
	1991	0		<1	264	<1	353	0		0		<1	303
	1992	0		0		<1	54	<1	518	<1	518	<1	135
	1993	0		<1	415	<1	143	0		0		<1	209
	1994	<1	502	<1	465	<1	452	0		0		<1	475
	1995	<1	499	<1	425	0		0		<1	139	<1	417
	Spot	1987 <sup>d</sup>	2	244	2	248	<1	248	2	214	0		1
1988		3	245	1	235	<1	225	1	243	<1	237	1	242
1989		<1	210	1	230	<1	277	<1	230	2	236	<1	237
1990		<1	319	<1	246	<1	246	1	212	1	238	<1	227
1991		<1	238	1	231	<1	210	1	217	<1	230	<1	220
1992		<1	231	1	227	<1	227	1	241	<1	257	<1	236
1993	1	229	<1	228	<1	231	2	229	2	267	1	240	
1994	1	230	1	239	<1	259	1	231	<1	244	1	236	
1995	<1	238	1	227	<1	243	2	217	0		1	225	
Spotted seatrout	1987 <sup>a</sup>	<1	408	<1	403	<1	397	<1	516	0		<1	417
	1988	3	410	2	431	1	397	<1	440	<1	469	2	414
	1989	1	419	3	431	1	419	1	428	<1	445	1	426

Table 7. (Cont'd.)

Species	Year	Gulf-17		Gulf-18		Gulf-19		Gulf-20		Gulf-21		Coastwide	
		No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length
Spotted seatrout (Cont'd.)	1990	2	440	2	417	<1	431	<1	457	1	473	1	437
	1991	3	406	2	441	1	421	1	399	<1	424	1	415
	1992	<1	432	2	428	2	423	1	431	<1	489	1	426
	1993	1	430	1	432	1	447	1	420	<1	501	1	438
	1994	1	432	1	444	1	434	3	402	<1	510	2	424
	1995	2	404	2	400	2	405	2	408	1	433	2	406
Striped mullet	1987 <sup>a</sup>	13	393	5	358	1	351	5	343	17	349	7	368
	1988	19	362	32	342	7	344	14	356	5	346	14	351
	1989	39	370	28	344	3	334	1	360	8	341	15	358
	1990	44	350	52	336	5	333	6	349	6	376	21	344
	1991	23	345	65	338	34	320	25	326	13	326	32	330
	1992	34	343	51	341	42	341	25	355	10	344	34	343
	1993	22	350	24	341	14	334	10	357	13	355	16	345
	1994	36	349	24	339	28	336	37	356	6	369	29	345
	1995	40	363	49	349	51	336	15	379	8	366	37	349
	Total finfishes	1987 <sup>a</sup>	23	327	9	305	6	266	10	295	18	332	12
1988		54	322	44	326	43	141	40	189	11	343	41	237
1989		52	341	48	288	20	218	39	100	15	298	34	254
1990		59	337	63	314	16	309	18	269	13	323	32	319
1991		50	322	80	309	45	293	46	284	20	324	48	304
1992		37	338	65	316	55	322	35	316	13	334	44	323
1993		30	338	32	324	21	326	22	283	23	314	25	321
1994		70	328	36	294	35	337	155	154	14	245	62	247
1995		61	335	65	315	57	337	27	320	15	335	49	330
SHELLFISHES		1987 <sup>a</sup>	<1	118	<1	159	0		0		0		<1
	1988	2	117	<1	143	<1	137	<1	138	<1	126	1	125
	1989	2	137	2	135	<1	140	0		<1	153	1	137
	1990	5	139	7	136	<1	129	<1	132	<1	128	2	137
	1991	7	143	20	137	5	127	1	123	1	131	6	136
	1992	3	133	3	126	2	142	<1	88	1	132	2	133
	1993	1	133	3	132	1	136	<1	132	<1	127	1	133
	1994	3	145	15	148	3	144	1	137	<1	129	4	146
	1995	3	124	6	132	1	134	<1	156	<1	151	2	130

<sup>a</sup> Values include Oct-Nov only.

Table 8. Seasonal (May-Nov) mean catch rates (No./ha) and mean total lengths (mm) of select fishes and shellfishes caught with 18.3-m bag seines in 5 Texas gulf shoreline areas during 1987-95. Blank indicates no measurement taken.

Species	Year	Gulf-17		Gulf-18		Gulf-19		Gulf-20		Gulf-21		Coastwide		
		No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	
<b>FINFISHES</b>														
Atlantic croaker	1987 <sup>a</sup>	0		0		0		0		0		0		
	1988	30	37	1	62	1	64	0	0	0	0	0	7	
	1989	3	32	0		0		0	0	0	0	0	32	
	1990	1	171	1	84	0		<1	22	2	157	1	127	
	1991	50	31	0		1	150	5	31	0		13	33	
	1992	0		1	91	1	61	0	0	0	0	1	70	
	1993	1	193	4	113	0		0	0	2	25	1	119	
	1994	1	28	0		0		0	0	0	0	<1	28	
	1995	0		<1	115	3	131	0	0	27	94	4	104	
	1987 <sup>a</sup>	0		0		0		0	0	0	0	0	0	
1988	1	111	1	104	0	168	0	0	0	1	182	0	132	
1989	2	170	0		0		0	0	0	0	<1	170		
1990	4	154	6	142	1	114	0	0	0	0	0	2	143	
1991	3	151	2	151	1	193	0	0	0	0	0	1	162	
1992	4	159	1	179	0		0	0	0	0	0	1	160	
1993	2	122	0		0		0	0	0	0	0	1	122	
1994	1	125	8	63	<1	55	<1	0	0	0	0	2	74	
1995	0		<1	161	0		0	0	0	0	0	<1	161	
Gulf menhaden	1987 <sup>a</sup>	0		0		4	48	0	0	0	0	0	1	48
	1988	2	93	22	87	5	87	28	37	2	74	10	63	
	1989	2	86	6	76	9	100	0	0	2	81	5	92	
	1990	3	59	0		5	57	1	83	17	5	5	68	
	1991	0		3	46	2	71	0	0	0	0	1	62	
	1992	1	91	9	72	4	50	0	0	0	0	3	63	
	1993	13	39	9	87	7	70	0	0	7	0	7	59	
	1994	0		2	56	6	32	1	33	6	82	3	45	
	1995	1	65	1	40	114	28	1	31	0	0	39	28	
	King mackerel	1987 <sup>a</sup>	0		0		0		0	0	0	0	0	0
1988		0		0		0		0	0	2	67	<1	67	
1989		0		0		0		0	0	0	0	0	0	
1990		0		0		3	48	0	0	0	0	1	48	
1991		0		0		0		0	0	0	0	0	0	
1992		0		0		0		0	0	0	0	0	0	
1993		0		0		<1	45	0	0	0	0	<1	45	
1994		0		0		0		0	0	0	0	0	0	
1995		0		0		1	47	5	56	0	0	1	54	
1987 <sup>a</sup>		0		0		1	85	0	0	0	0	<1	85	
1988	0		1	100	1	122	0	0	103	105	12	105		
1989	0		0		<1	69	0	0	0	0	<1	69		
1990	0		0		1	97	<1	52	2	133	1	101		
1991	0		2	98	26	71	1	55	27	67	12	70		
1992	1	68	1	125	0		0	0	1	62	<1	77		

Table 8. (Cont'd.)

Species	Year	Gulf-17		Gulf-18		Gulf-19		Gulf-20		Gulf-21		Coastwide	
		No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length
Pinfish (Cont'd.)	1993	1	66	5	89	1	127	0	0	6	79	2	88
	1994	0		1	75	1	53	1	77	0		1	65
	1995	0		3	82	0		1	72	10	111	2	100
	1987 <sup>a</sup>	0		0		0		0		0		0	
	1988	0		12	48	1	61	0		0		2	50
Sand seatrout	1989	11	44	0		0		0		0		3	44
	1990	0		0		<1	124	0		0		<1	124
	1991	5	31	2	40	6	86	0		0		54 (Cont'd.)	
	1992	2	34	<1	42	0		0		0		1	35
	1993	74	65	0		<1	96	0		0		17	65
	1994	7	33	0		50	33	15	41	1	71	22	34
	1995	1	43	4	106	10	29	1	32	0		4	38
	1987 <sup>a</sup>	0		0		0		0		0		0	
	1988	0		5	107	1	126	0		0		0	112
	1989	1	114	10	91	0		0		0		2	95
Southern flounder	1990	0		2	107	1	183	0		0		1	151
	1991	0		0		0		2	102	0		<1	102
	1992	1	134	11	120	0		4	90	2	162	3	116
	1993	4	135	11	110	0		0		0		2	119
	1994	6	106	6	128	0		0		0		2	114
	1995	1	81	2	128	0		0		0		<1	106
	1987 <sup>a</sup>	41	50	0		0		0		0		9	50
	1988	0		1	59	2	53	0		2	110	1	64
	1989	0		6	37	0		8	60	0		2	51
	1990	0		1	25	2	35	0		0		1	34
Spanish mackerel	1991	0		<1	40	0		0		0		<1	40
	1992	0		0		0		1	55	0		<1	55
	1993	0		0		1	54	14	25	0		3	27
	1994	0		4	53	2	42	0		0		1	48
	1995	0		0		<1	35	25	41	24	101	7	63
	1987 <sup>a</sup>	0		0		0		0		0		0	
	1988	0		1	80	0		0		52	92	6	91
	1989	0		0		1	78	0		2	104	<1	89
	1990	1	182	0		1	86	<1	66	0		1	119
	1991	0		<1	182	0		<1	64	0		<1	122
Spot	1992	1	109	0		0		1	26	0		<1	81
	1993	0		0		9	87	0		4	81	3	87
	1994	1	78	17	74	1	68	0		0		3	74
	1995	0		<1	44	0		2	34	15	112	2	98
	1987 <sup>a</sup>	7	26	0		0		2	100	14	146	4	84
	1988	50	97	36	115	22	59	1	88	0		24	88
	1989	253	86	42	90	15	187	1	93	3	191	69	95
1990	49	66	86	79	3	170	10	32	5	155	27	75	

Table 8. (Cont'd.)

Species	Year	Gulf-17		Gulf-18		Gulf-19		Gulf-20		Gulf-21		Coastwide			
		No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length	No./ha	Length		
Striped mullet (Cont'd.)	1991	18	173	141	130	23	140	1	144	2	106	32	138		
	1992	11	70	10	138	4	73	2	53	3	127	6	89		
	1993	5	160	5	62	5	159	4	64	0		4	128		
	1994	90	94	2	59	27	181	35	23	1	52	37	103		
	1995	5	112	25	122	3	126	6	35	5	55	7	104		
Total finfishes	1987 <sup>a</sup>	344	66	449	60	475	73	668	45	2,142	83	659	69		
	1988	1,046	65	6,271	96	2,351	58	1,702	48	3,164	84	2,572	74		
	1989	2,413	95	2,794	75	3,590	68	9,527	59	1,159	80	4,009	69		
	1990	1,168	76	1,125	71	1,292	55	3,075	46	1,081	105	1,538	61		
	1991	1,140	84	1,625	83	4,006	64	7,512	54	2,140	73	3,439	63		
	1992	1,312	84	1,029	65	1,090	59	2,514	47	923	58	1,371	61		
	1993	1,545	82	4,223	51	2,267	92	4,671	46	1,158	69	2,685	66		
	1994	1,881	69	3,633	56	1,458	62	7,244	52	2,260	56	2,990	57		
	1995	1,176	62	8,173	50	2,305	59	6,955	48	4,322	85	3,926	56		
	SHELLFISHES	Blue crab	1987 <sup>a</sup>	0	0	0	0	0	0	0	0	3	22	<1	22
1988			14	101	1	25	4	83	0		0		5	93	
1989			33	95	65	34	2	108	0		0		17	60	
1990			11	85	52	90	1	113	1	24	0		10	89	
1991			42	107	72	69	24	117	1	91	0		28	96	
1992			30	92	49	84	9	116	2	21	1	174	17	92	
1993			20	104	26	85	4	109	1	157	0		10	98	
1994			19	119	26	105	17	142	2	74	0		14	124	
1995			43	78	25	101	2	90	1	33	0		14	84	
Brown shrimp			1987 <sup>a</sup>	0	0	0	0	0	0	0	0	0	0	0	0
			1988	7	52	0	96	3	76	0		1	46	3	60
			1989	7	56	0		0		0		0		2	56
			1990	0		47	76	0		0		0		7	76
			1991	9	44	1	54	<1	58	0		0		2	45
			1992	27	66	10	52	0	31	1		0		8	63
	1993	13	59	1	39	1	92	0		0		3	61		
	1994	12	66	6	68	4	47	0		0		5	61		
	1995	0		4	60	1	56	0		0		1	58		
	White shrimp	1987 <sup>a</sup>	11	78	16	71	71	69	2	72	0		29	70	
		1988	35	64	6	77	2	61	<1	45	1	69	10	65	
		1989	38	58	4	70	20	65	2	52	0		16	61	
		1990	8	75	9	57	0	59	<1	59	0		3	67	
		1991	664	53	4	70	1	69	0		0		154	53	
		1992	285	75	12	86	2	81	0		0		68	75	
1993		49	57	7	61	<1	60	0		1	38	12	57		
1994		43	68	2	67	2	71	0		0		11	68		
1995		626	66	44	60	12	52	13	64	0		157	66		

<sup>a</sup> Values include Oct-Dec only.

Table 9. Annual mean catch rate (No./h) and mean total lengths (mm) of selected fishes and shellfishes caught with 6.1-m trawls within the Intracoastal Waterway in Texas bay systems during 1992-95.

Species	Year	East										Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		Sabine Lake No./h Length	Galveston No./h Length	Mataforda No./h Length	Mataforda No./h Length	San Antonio No./h Length	Aransas No./h Length	Christi No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length	Madre No./h Length		
FINFISHES																					
Atlantic croaker	1992	88	133	306	125	38	109	130	96	1,181	83	684	82	26	114	27	119	45	122	241	98
	1993	190	132	135	119	57	131	270	108	355	108	155	109	23	147	5	174	37	134	130	119
	1994	107	122	122	117	36	117	23	130	172	89	72	92	6	107	7	150	100	142	79	117
	1995	62	128	98	139	86	94	343	75	216	92	81	100	38	117	9	142	60	120	97	104
	1992	1	234	<1	250	1	180	0	0	0	0	0	0	0	0	1	240	<1	282	<1	235
Black drum	1993	6	197	<1	142	<1	173	<1	190	0	0	0	0	0	<1	264	0	0	0	1	199
	1994	2	180	1	193	<1	148	<1	179	<1	81	<1	140	0	<1	328	0	0	0	1	188
	1995	<1	215	1	172	<1	187	0	0	0	0	0	0	0	1	240	0	0	0	<1	207
	1992	32	110	11	153	2	151	8	129	81	135	44	130	<1	176	0	0	<1	186	18	129
Gafftop-sail catfish	1993	13	121	18	145	2	135	31	123	80	146	36	118	<1	167	0	1	170	17	135	
	1994	18	125	5	149	7	152	3	137	92	123	20	134	<1	134	0	134	2	166	14	128
	1995	4	117	6	148	10	134	15	138	46	124	19	132	1	192	0	0	2	148	9	131
	1992	1	120	5	93	5	89	13	113	19	90	25	102	1	138	<1	119	1	139	6	101
menhaden	1993	16	77	9	102	<1	75	2	124	6	105	8	85	7	62	0	0	1	126	6	88
	1994	1	114	1	97	1	116	<1	131	6	97	1	133	1	138	0	0	<1	115	1	106
	1995	1	103	<1	103	<1	105	1	134	1	93	<1	127	1	161	0	0	<1	148	<1	114
	1992	0	117	2	142	2	121	6	106	5	95	31	113	206	105	57	119	18	112	31	113
Pinfish	1993	2	117	2	113	2	108	4	109	13	91	93	110	101	115	25	131	19	97	20	113
	1994	0	127	31	104	7	117	3	118	7	137	3	98	3	145	<1	245	17	157	11	121
	1995	0	119	22	114	9	120	47	98	10	129	7	114	9	128	1	184	25	145	15	120
	1992	0	113	17	127	4	134	4	110	32	114	12	133	4	149	1	138	11	157	11	127
Red drum	1993	0	115	22	129	15	129	7	121	43	120	9	112	6	120	1	221	3	152	17	122
	1994	<1	257	0	0	0	0	0	0	0	0	<1	42	0	0	0	0	0	0	<1	218
	1995	0	0	0	0	0	0	0	0	0	0	0	0	0	0	<1	33	<1	283	<1	148
	1992	9	113	17	127	4	134	4	110	32	114	12	133	4	149	1	138	11	157	11	127
Sand seatrout	1993	36	115	22	129	15	129	7	121	43	120	9	112	6	120	1	221	3	152	17	122
	1994	6	127	31	104	7	117	3	118	7	137	3	98	3	145	<1	245	17	157	11	121
	1995	7	119	22	114	9	120	47	98	10	129	7	114	9	128	1	184	25	145	15	120
	1992	<1	160	<1	155	<1	132	<1	137	<1	96	<1	96	<1	121	<1	405	<1	115	<1	172
Sheeps-head	1993	<1	134	<1	190	<1	144	0	116	1	116	<1	100	0	0	0	0	<1	309	<1	171
	1994	0	0	<1	187	1	168	<1	139	<1	157	<1	108	0	0	<1	139	<1	157	<1	152
	1995	0	0	1	144	<1	147	<1	136	<1	86	<1	97	<1	92	<1	111	0	0	<1	126
Southern flounder	1992	1	256	5	239	1	220	3	209	1	211	1	193	<1	198	<1	398	<1	248	2	236
	1993	2	252	2	256	3	183	3	167	3	149	1	190	<1	161	<1	217	1	204	2	208
	1994	1	199	3	229	3	195	2	222	3	161	1	169	<1	177	1	210	1	225	2	208
	1995	2	246	6	223	2	231	3	237	2	206	1	184	<1	208	<1	262	<1	364	2	227

Table 9. (Cont'd.)

Species	Year	Sabine Lake		Galveston		East		Matagorda		Matagorda		San Antonio		Arkansas		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide			
		No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
Spot	1992	3	134	149	124	10	122	44	110	150	102	55	103	38	111	12	148	32	127	57	118				
	1993	17	119	33	121	9	124	117	115	97	112	80	94	231	120	6	150	15	151	45	116				
	1994	5	134	17	126	10	115	18	112	77	110	15	126	222	136	2	163	99	118	37	122				
	1995	16	84	16	129	7	123	26	104	154	107	32	124	107	151	3	146	52	155	35	124				
Spotted seatrout	1992	<1	184	3	144	2	145	<1	165	3	115	5	120	<1	158	2	200	3	205	2	158				
	1993	2	188	1	159	4	169	2	154	2	138	1	125	<1	100	1	202	2	279	2	175				
	1994	2	172	2	145	2	175	1	162	4	153	2	115	<1	166	1	251	1	259	2	175				
	1995	1	164	2	170	4	174	2	134	6	152	5	143	<1	189	5	166	1	235	3	164				
Striped mullet	1992	<1	136	3	183	1	147	0		2	125	3	188	<1	281	<1	291	<1	250	1	181				
	1993	10	215	<1	202	<1	338	<1	216	0		<1	222	<1	136	<1	294	0	216	2	216				
	1994	1	138	<1	254	0		<1	146	<1	161	0		<1	299	0		0	162	<1	162				
	1995	0		<1	316	<1	304	0		<1	124	0		<1	280	0		<1	271	<1	259				
Total finfish	1992	291	131	585	125	83	115	322	98	1,670	94	972	94	333	111	157	119	200	134	461	110				
	1993	453	127	266	122	432	127	560	105	699	109	669	107	452	121	85	126	177	146	302	118				
	1994	196	127	270	114	104	126	210	108	523	107	234	111	433	134	67	112	353	139	243	121				
	1995	139	120	198	131	156	115	692	96	729	104	405	116	368	140	79	131	345	150	291	119				
SHELLFISHES																									
Blue crab	1992	40	74	67	73	64	78	41	62	222	55	238	57	14	94	66	95	26	72	77	68				
	1993	59	69	47	77	57	93	76	81	63	74	129	78	10	109	44	98	59	90	59	82				
	1994	52	71	39	77	84	73	57	72	54	64	81	70	12	76	49	66	49	86	52	73				
	1995	40	63	51	82	108	76	48	82	71	66	53	68	29	72	28	80	28	93	47	76				
Brown shrimp	1992	44	79	209	79	21	84	19	84	269	81	340	82	34	83	92	99	33	84	117	83				
	1993	135	81	74	84	48	78	39	89	323	78	301	86	27	90	33	94	48	82	104	83				
	1994	13	94	107	92	12	79	19	98	180	89	103	86	20	94	74	106	80	85	70	89				
	1995	4	82	42	77	19	79	85	88	242	81	175	84	53	90	49	92	69	89	69	85				
Pink shrimp	1992	0		0		<1	91	<1	87	2	74	40	73	18	86	48	104	57	90	20	93				
	1993	0		0		2	89	1	110	2	90	19	71	13	91	3	103	29	87	7	86				
	1994	0		<1	98	8	98	6	105	12	83	24	88	18	87	6	96	29	88	9	90				
	1995	0		<1	94	7	82	6	98	67	86	21	86	87	97	10	98	62	96	22	93				
White shrimp	1992	35	100	77	90	8	82	5	92	28	85	42	91	5	94	4	104	5	106	28	92				
	1993	75	85	28	91	62	89	50	86	26	85	30	88	3	107	12	102	15	102	35	89				
	1994	12	91	79	72	33	92	7	91	53	82	38	93	2	112	8	105	31	97	32	83				
	1995	16	79	25	84	36	87	24	83	33	76	45	86	7	114	22	94	62	100	30	89				






Figure 1. Texas gulf shoreline and Texas Territorial Sea (TTS).




Figure 2. Spring gill net mean catch rates (No./h  $\pm$  1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1976-95.




Figure 3. Fall gill net mean catch rates (No./h  $\pm$  1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1975-95.




Figure 4. Spring gill net mean total lengths (mm  $\pm$  1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1976-95.




Figure 5. Fall gill net mean total lengths (mm  $\pm$  1SE) for red drum, black drum, spotted seatrout and Atlantic croaker during 1975-95.




Figure 6. Seasonal bag seine mean catch rates (No./ha) for juvenile red drum (Nov-Mar), black drum (Jun-Jul), spotted seatrout (Jul-Nov) and Atlantic croaker (Feb-May) during 1978-95. Red drum 35-75 mm, spotted seatrout 20-75 mm, black drum 35-110 mm and Atlantic croaker 30-85 mm are considered to be young-of-the-year.




Figure 7. Seasonal bag seine mean total lengths (mm + 1SE) for juvenile red drum (Nov-Mar), black drum (Jun-Jul), spotted seatrout (Jul-Nov) and Atlantic croaker (Feb-May) during 1978-95. Red drum 35-75 mm, spotted seatrout 20-75 mm, black drum 35-110 mm and Atlantic croaker 30-85 mm are considered to be young-of-the-year.




Figure 8. Seasonal bag seine mean catch rates (No./ha) for juvenile brown shrimp (Apr-Jul), white shrimp (Jul-Nov) and blue crab (Mar-Jun) during 1978-95. Brown and white shrimp 33-82 mm and blue crab 13-42 mm are considered to be young-of-the year.






Figure 9. Seasonal bag seine mean total lengths (mm  $\pm$  1SE) for brown shrimp (Apr-Jul), white shrimp (Jul-Nov) and blue crab (Mar-Jun) during 1978-95. Brown and white shrimp 33-82 mm and blue crab 13-42 mm are considered to be young-of-the-year.




Figure 10. Annual bay trawl catch rates (No./h  $\pm$  1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95.




Figure 11. Annual bay trawl mean total lengths (mm  $\pm$  1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95.




Figure 12. Annual gulf trawl mean catch rates (No./h + 1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95.




Figure 13. Annual gulf trawl mean total lengths (mm ± 1SE) for brown shrimp, white shrimp, blue crab and Atlantic croaker during 1982-95.




Figure 14. Annual mean catch rates (No./h) for Eastern oyster spat ( $\leq 25$  mm), small oysters (26-75 mm) and market oysters ( $> 76$  mm) during 1984-95.




Figure 15. Annual mean total lengths (mm) for small and market Eastern oysters during 1984-95.

Appendix A. Summary of historical sampling dates, gear description, procedures, dates, number of samples collected, weighting factors, and list of species collected.



Table A.1. Historical sampling dates (month/year) by bay system and gear.

GEAR	SABINE	GALVESTON	EAST MATAGORDA	MATAGORDA	SAN ANTONIO	ARANSAS-COPANO	CORPUS CHRISTI	UPPER LAGUNA	LOWER LAGUNA
GILL NET	April 1986- Present	Nov. 1975- Present	Oct. 1976- Present	Nov. 1975- Present	Nov. 1975- Present	Nov. 1975- Present	Nov. 1975- Present	Nov. 1975- Present	Nov. 1975- Present
GULF TRAWL	Jul. 1986- Present	Aug. 1985- Present	Not used.	Not used.	Aug 1985- Present	Not used.	Feb. 1985- Present	Not used.	Aug. 1985- Present
BAY TRAWL	Jan. 1986- Present	Jan. 1982- Present	April 1987- Present	May 1982- Present	Jan. 1982- Present	Jan. 1982- Present	May 1982- Present	May 1982- Present	May 1982- Present
ICWW TRAWL	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995	Jan. 1992- 1995
BEACH SEINE	Oct. 1987- 1995	Oct. 1987- 1995	Oct. 1987- 1995	Not used.	Oct. 1987- 1995	Oct. 1987- 1991	Not used.	Oct. 1987- 1995	Oct. 1987- 1995
BEACH BAG SEINE	Oct. 1987- 1995	Oct. 1987- 1995	Oct. 1987- 1995	Not used.	Oct. 1987- 1995	Oct. 1987- 1991	Not used.	Oct. 1987- 1995	Oct. 1987- 1995
BAY BAG SEINE	Jan. 1986- Present	Oct. 1977- Present	Feb. 1983- Present	Oct. 1977- Present	Oct. 1977- Present	Oct. 1977- Present	Oct. 1977- Present	Oct. 1977- Present	Oct. 1977- Present
OYSTER REEF DREDGE	Jan. 1986- 1991	Oct. 1984- Present	Jan. 1986- 1991	Jan 1986- Present	Jan. 1986- Present	Jan. 1986- Present	Jan. 1986- 1991	Not used.	Jan. 1986- 1991
NON-REEF DREDGE	1986-1989	1985-1989	1986-1989	1986-1989	1986-1989	1986-1989	1986-1989	1986-1988	1986-1988

Table A. 2. Gear descriptions.

GEAR	GEAR DESCRIPTION
Gill Net	Monofilament, 183 m long; 1.2 m deep with separate 45.7-m sections of 7.6-, 10.2- (#12 monofilament), 12.7- and 15.2-cm (#18 monofilament) stretched mesh tied together in ascending mesh size.
Trawl	6.1 m wide at mouth with 3.8-cm stretched nylon multifilament mesh throughout, and doors 1.2 m long and 0.6 m tall.
Beach Seine	60.9-m long; 1.8-m deep with 7.6-cm stretched #12 monofilament mesh.
Bag Seine	18.3 m long; 1.8 m deep with 1.3-cm stretched nylon multifilament mesh in the 1.8 m wide central bag with remaining webbing 1.9-cm stretched mesh.
Oyster Dredge	Louisiana style 8-tooth: 46 cm wide, 25 cm tall with a 36-cm deep bag. 6 bottom rows of linked metal rings 5 cm in diameter; four top rows of 7.6-cm mesh webbing made of 0.8-cm nylon rope.

Table A.3. Historical sampling procedures by gear.

GEAR	HISTORICAL SAMPLING PROCEDURES
GILL NET	<p>Monofilament gill nets have been systematically used in 7 Texas bay systems since November 1975; East Matagorda Bay was added in fall 1976 and Sabine Lake in April 1986 (Figure 1). Prior to September 1984, sites for setting gill nets during spring (Ten week period, generally, 15 April-15 June) and fall (Ten week period, generally, 15 September-15 November) were randomly selected from about 100 stations in each bay system (McEachron and Green 1985). Beginning September 1984 current site selection methods were adopted.</p> <p>Prior to fall 1981, no less than one nor more than 18 overnight gill net sets occurred in each season in each bay system. Since fall 1981, 45 gill nets were set overnight during each season in each bay system except East Matagorda Bay. In East Matagorda Bay, from fall 1981 to spring 1984 not less than six nor more than 12 gill nets were set during each season; since fall 1984, 20 sets were set in each season.</p>
GULF TRAWLS	<p>Trawls have been systematically used in 4 Gulf areas of Texas Territorial Seas since August 1985, and a total of 5 areas since July 1986.</p>
BAY TRAWLS	<p>Trawls have been systematically used in 3 Texas bays since January 1982 and in 7 bays since May 1982; Sabine Lake was added January 1986 and East Matagorda Bay April 1987. From January 1982 to present, 20 monthly samples were collected in the Galveston, San Antonio and Aransas systems. Beginning in May 1982, 20 monthly samples were collected in Matagorda and Corpus Christi Bay systems, and 10 in upper and lower Laguna Madre. Beginning in January 1986 in the Sabine system, and in April 1987 in the East Matagorda system, 10 monthly samples were collected.</p>
ICWW TRAWLS	<p>From 1992 through 1995, 6 monthly samples were collected in each of the 9 bay systems along the Texas coast.</p>

Table A.3. (Cont'd.)

BEACH SEINE	Beach seines have been systematically used on Texas Gulf beaches since October 1987. From October 1987 to 1995, 6 beach seine samples were collected each month along Gulf beach shoreline areas.
BEACH BAG SEINE	Beach bag seine samples have been systematically used on Texas Gulf beaches since October 1987. From October 1987 to 1995, 6 beach bag seine samples were collected each month along Gulf beach shoreline areas.
BAY BAG SEINE	<p>Bay bag seine samples have been systematically collected in 7 Texas bay systems since October 1977; sample collection began in the East Matagorda system February 1983, and in the Sabine system in January 1986. Bay bag seine samples were collected by pulling the seine 15.2-30.5 m parallel to shore prior to September 1984; since then it has been pulled 15.2 m. Prior to September 1984, sites for sampling with bag seines (monthly) were randomly selected from about 100 stations in each bay system (McEachron and Green 1985). Prior to October 1981, six bag seine samples were collected each month in each bay system (except during June 1978 when no samples were collected). During October 1981 through August 1984 10 bag seine samples were collected each month in each bay system; half of the samples were collected during each of the first and last two fullest weeks of each month (McEachron and Green 1985). Beginning September 1984, half of the monthly samples were collected during the 1st-15th and half during the 16th-31st of each month. From April 1988 through December 1989 12 bag seine samples were collected each month in each bay system. Beginning January 1990, 16 bag seine samples were collected each month in each bay system. Beginning January 1992, 20 samples were collected in each bay system each month, except in East Matagorda where only 10 samples were collected per month.</p>

Table A.3. (Cont'd.)

<p>OYSTER REEF DREDGE</p>	<p>Oyster dredges have been systematically used in Texas bays since January 1986. The number of monthly samples collected in the Galveston system were: 20 in 1984; 80 in 1985; and 56 in 1986-1991. Monthly samples collected in the Aransas system were: 56 in 1986-1989; and 26 in 1990-1991. From 1986 to 1991 10 samples per month were collected in Sabine Lake and the Lower Laguna Madre and 26 monthly samples were collected in the Matagorda, San Antonio, Corpus Christi and East Matagorda systems. Beginning January 1992 30 samples were collected each month in the Galveston system, and 20 samples in the Matagorda, San Antonio and Aransas systems. Sampling in other systems was discontinued beginning January 1992.</p>
<p>NON-REEF DREDGE</p>	<p>Non-reef dredge samples were systematically collected in Texas bays from 1985-1989. In 1985 10 monthly samples were collected in the Galveston system. From 1986-1989 10 monthly samples were collected in all bay systems.</p>





Table A.6. Number of samples collected during routine monitoring in 5 Texas Gulf surf zones by gear and year.

	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
Beach Bag	9	15	25	21	12	82
Seine	28	56	101	67	42	294
	29	55	91	74	42	291
	30	54	98	70	42	294
	26	58	97	71	42	294
	27	57	84	42	42	252
	28	56	84	42	42	252
	27	57	84	42	42	252
Beach	9	15	26	22	12	84
Seine	28	56	100	68	42	294
	29	55	91	74	42	291
	30	54	98	70	42	294
	26	58	97	71	42	294
	27	57	84	42	42	252
	28	56	84	42	42	252
	27	57	84	42	42	252

Table A.7. Number of Gulf trawl samples collected during routine monitoring in 5 Gulf zones, by Gulf area and year.

	Sabine Lake	Galveston	Port O'Connor	Port Aransas	Port Isabel	Coastwide
Gulf Trawl	0	80	80	176	80	416
	112	192	192	192	192	880
	192	192	192	192	192	960
	192	192	192	184	184	952
	192	192	192	184	189	949
	192	192	192	192	192	960
	192	192	192	184	192	952
	192	192	192	184	192	952
	192	192	192	192	192	960
	192	192	187	192	192	955



Table A.8. Weighting factors used in calculating coastwide average catch rates.

Area	Gill net and <sup>a</sup> bay bag seine	Bay <sup>b</sup> trawl	ICWW <sup>c</sup> trawl	Oyster <sup>d</sup> dredge	Gulf <sup>e</sup> trawl
<b>BAY SYSTEM</b>					
Sabine	75.6	1.220	57.6		
Galveston	411.2	9.408	61.8	126	
East Matagorda	64.4	0.101	23.4		
Matagorda	284.8	6.288	27.4	42	
San Antonio	225.2	3.680	27.0	66	
Aransas	263.5	2.251	25.2	55	
Corpus Christi	171.3	3.357 <sup>f</sup>	13.8		
Upper Laguna Madre	222.3	1.534	55.1		
Lower Laguna Madre	252.1	1.153	46.6		
Total	1,970.4	28.992	337.9	289	
<b>GULF AREA</b>					
Sabine					266
Galveston					273
Port O'Connor					277
Port Aransas					257
Port Isabel					248
Total					1,317

<sup>a</sup> Equals miles of shoreline (Matlock and Osborn 1982. Shallow-water surface areas and shoreline distances on the Texas coast).

<sup>b</sup> Equals total bay surface area (divided by 10,000) minus 1977 estimate of shallow water area (<=1.2 m)(for the Lagunas Madre) or minus the mean of 1972 and 1977 estimates (for other bays)(Matlock and Osborn 1982).

<sup>c</sup> Equals nautical miles of ICWW.

<sup>d</sup> Equals total number of grids containing oyster reef.

<sup>e</sup> Equals total number of Gulf trawlable grids.

<sup>f</sup> No estimate was available for 1977 shallow water area, so 1977 area was estimated as proportion of sampling grid zones that are designated as trawls grids, times the total surface area of the bay.

Table A.9. Species caught (alphabetical by scientific name; Robins et al. 1991) in Texas marine waters by TPW sampling gear during 1975-1995. Common names in ( ) are assigned common names by TPW for identification purposes.

<u>Scientific Name</u>	<u>Common Name</u>
<b>Finfish</b>	
<u>Abudefduf saxatilis</u>	Sergeant major
<u>Achirus lineatus</u>	Lined sole
<u>Adinia xenica</u>	Diamond killifish
<u>Aetobatis narinari</u>	Spotted eagle ray
<u>Agonostomus monticola</u>	Mountain mullet
<u>Albula vulpas</u>	Bonefish
<u>Alectis ciliaris</u>	African pompano
<u>Alosa chrysochloris</u>	Skipjack herring
<u>Aluterus heudeloti</u>	Dotterel filefish
<u>Aluterus schoepfi</u>	Orange filefish
<u>Aluterus scriptus</u>	Scrawled filefish
<u>Ambloplites rupestris</u>	Rock bass
<u>Ameiurus melas</u>	Black bullhead
<u>Ameiurus natalis</u>	Yellow bullhead
<u>Amia calva</u>	Bowfin
<u>Anchoa hepsetus</u>	Striped anchovy
<u>Anchoa lyolepis</u>	Dusky anchovy
<u>Anchoa mitchilli</u>	Bay anchovy
<u>Anchoa nasuta</u>	Longnose anchovy
<u>Ancylopsetta dilecta</u>	Three-eye flounder
<u>Ancylopsetta quadrocellata</u>	Ocellated flounder
<u>Anquilla rostrata</u>	American eel
<u>Antennarius radiosus</u>	Singlespot frogfish
<u>Antennarius striatus</u>	Striated frogfish
<u>Aplodinotus grunniens</u>	Freshwater drum
<u>Archosargus probatocephalus</u>	Sheepshead
<u>Arius felis</u>	Hardhead catfish
<u>Astroscopus y-graecum</u>	Southern stargazer
<u>Bagre marinus</u>	Gafftopsail catfish
<u>Bairdiella chrysoura</u>	Silver perch
<u>Balistes capriscus</u>	Gray triggerfish
<u>Bascanichthys bascanium</u>	Sooty eel
<u>Bascanichthy scuticaris</u>	Whip eel
<u>Bathygobius soporator</u>	Frillfin goby
<u>Bellator militaris</u>	Horned searobin
<u>Bodianus pulchellus</u>	Spotfin hogfish
<u>Bollmannia communis</u>	Ragged goby
<u>Bothus robinsi</u>	Twospot flounder
<u>Brevoortia patronus</u>	Gulf menhaden
<u>Brevoortia gunteri</u>	Finescale menhaden
<u>Brotula barbata</u>	Bearded brotula
<u>Calamus leucosteus</u>	Whitebone porgy
<u>Cantherhines pullus</u>	Orangespotted filefish
<u>Canthidermis maculata</u>	Rough triggerfish
<u>Caranx bartholomaei</u>	Yellow jack

Table A.9. (Cont'd.)

Scientific Name	Common Name
<b>Finfish (Cont'd.)</b>	
<u>Caranx crysos</u>	Blue runner
<u>Caranx hippos</u>	Crevalle jack
<u>Caranx latus</u>	Horse-eye jack
<u>Caranx ruber</u>	Bar jack
<u>Carassius auratus</u>	Goldfish
<u>Carcharhinus acronotus</u>	Blacknose shark
<u>Carcharhinus brevipinna</u>	Spinner shark
<u>Carcharhinus falciformis</u>	Silky shark
<u>Carcharhinus isodon</u>	Finetooth shark
<u>Carcharhinus leucas</u>	Bull shark
<u>Carcharhinus limbatus</u>	Blacktip shark
<u>Carcharhinus obscurus</u>	Dusky shark
<u>Carcharhinus plumbeus</u>	Sandbar shark
<u>Carcharhinus porosus</u>	Smalltail shark
<u>Centropomus parallelus</u>	Fat snook
<u>Centropomus undecimalis</u>	Common snook
<u>Centropristis ocyurus</u>	Bank sea bass
<u>Centropristis philadelphica</u>	Rock sea bass
<u>Chaetodipterus faber</u>	Atlantic spadefish
<u>Chaetodon ocellatus</u>	Spotfin butterflyfish
<u>Chasmodes bosquianus</u>	Striped blenny
<u>Chilomycterus schoepfi</u>	Striped burrfish
<u>Chloroscombrus chrysurus</u>	Atlantic bumper
<u>Citharichthys macrops</u>	Spotted whiff
<u>Citharichthys spilopterus</u>	Bay whiff
<u>Conodon nobilis</u>	Barred grunt
<u>Ctenopharyngodon idella</u>	Grass carp
<u>Cyclopsetta chittendeni</u>	Mexican flounder
<u>Cyclopsetta fimbriata</u>	Spotfin flounder
<u>Cynoscion arenarius</u>	Sand seatrout
<u>Cynoscion nebulosus</u>	Spotted seatrout
<u>Cynoscion nothus</u>	Silver seatrout
<u>Cyprinodon variegatus</u>	Sheepshead minnow
<u>Cyprinus carpio</u>	Common carp
<u>Dasyatis americana</u>	Southern stingray
<u>Dasyatis sabina</u>	Atlantic stingray
<u>Dasyatis say</u>	Bluntnose stingray
<u>Decapterus punctatus</u>	Round scad
<u>Diapterus auratus</u>	Irish pompano
<u>Dibranchius atlanticus</u>	Atlantic batfish
<u>Diodon hystrix</u>	Porcupinefish
<u>Diplectrum bivittatum</u>	Dwarf sand perch
<u>Diplectrum formosum</u>	Sand perch
<u>Diplodus holbrooki</u>	Spottail pinfish
<u>Dormitator maculatus</u>	Fat sleeper
<u>Dorosoma cepedianum</u>	Gizzard shad
<u>Dorosoma petenense</u>	Threadfin shad
<u>Echeneis naucrates</u>	Sharksucker

Table A.9. (Cont'd.)

<u>Scientific Name</u>	<u>Common Name</u>
<b>Finfish (Cont'd.)</b>	
<u>Echiophis intertinctus</u>	Spotted spoon-nose eel
<u>Elagatis bipinnulata</u>	Rainbow runner
<u>Elops saurus</u>	Ladyfish
<u>Epinephelus nigritus</u>	Warsaw grouper
<u>Epinephelus niveatus</u>	Snowy grouper
<u>Equetus umbrosus</u>	Cubbyu
<u>Erotelis smaragdus</u>	Emerald sleeper
<u>Etropus crossotus</u>	Fringed flounder
<u>Etrumeus teres</u>	Round herring
<u>Eucinostomus argenteus</u>	Spotfin mojarra
<u>Eucinostomus gula</u>	Silver jenny
<u>Eucinostomus lefroyi</u>	Mottled mojarra
<u>Eucinostomus melanopterus</u>	Flagfin mojarra
<u>Evorthodus lyricus</u>	Lyre goby
<u>Fundulus chrysotus</u>	Golden topminnow
<u>Fundulus grandis</u>	Gulf killifish
<u>Fundulus pulvereus</u>	Bayou killifish
<u>Fundulus similis</u>	Longnose killifish
<u>Gadella maraldi</u>	(Barbelless codlet)
<u>Gambusia affinis</u>	Western mosquitofish
<u>Gerres cinereus</u>	Yellowfin mojarra
<u>Gnathagnus egregius</u>	Freckled stargazer
<u>Gobiesox punctulatus</u>	Stippled clingfish
<u>Gobiesox strumosus</u>	Skilletfish
<u>Gobioides broussoneti</u>	Violet goby
<u>Gobiomorus dormitor</u>	Bigmouth sleeper
<u>Gobionellus boleosoma</u>	Darter goby
<u>Gobionellus hastatus</u>	Sharptail goby
<u>Gobionellus shufeldti</u>	Freshwater goby
<u>Gobiosoma bosc</u>	Naked goby
<u>Gobiosoma robustum</u>	Code goby
<u>Gonioplectrus hispanus</u>	Spanish flag
<u>Gunterichthys longipenis</u>	Gold brotula
<u>Gymnachirus texae</u>	Fringed sole
<u>Gymnothorax nigromarginatus</u>	Blackedge moray
<u>Gymnura micrura</u>	Smooth butterfly ray
<u>Haemulon aurolineatum</u>	Tomtate
<u>Halieutichthys aculeatus</u>	Pancake batfish
<u>Harengula jaguana</u>	Scaled sardine
<u>Hemicaranx amblyrhynchus</u>	Bluntnose jack
<u>Hemipteronotus novacula</u>	Pearly razorfish
<u>Hemiramphus balao</u>	Balao
<u>Hemiramphus brasiliensis</u>	Ballyhoo
<u>Hildebrandia flava</u>	Yellow conger
<u>Hippocampus erectus</u>	Lined seahorse
<u>Hippocampus zosterae</u>	Dwarf seahorse
<u>Histrio histrio</u>	Sargassumfish
<u>Holacanthus bermudensis</u>	Blue angelfish

Table A.9. (Cont'd.)

Scientific Name	Common Name
<b>Finfish (Cont'd.)</b>	
<u>Hoplostethus mediterraneus</u>	Armorhead
<u>Hypleurochilus geminatus</u>	Crested blenny
<u>Hyporhamphus unifasciatus</u>	Silverstripe halfbeak
<u>Hypsoblennius hentz</u>	Feather blenny
<u>Hypsoblennius ionthas</u>	Freckled blenny
<u>Ictalurus furcatus</u>	Blue catfish
<u>Ictalurus punctatus</u>	Channel catfish
<u>Ictiobus bubalus</u>	Smallmouth buffalo
<u>Ictiobus cyprinellus</u>	Bigmouth buffalo
<u>Isurus oxyrinchus</u>	Shortfin mako
<u>Jenkinsia lamprotaenia</u>	Dwarf herring
<u>Kyphosus incisor</u>	Yellow chub
<u>Kyphosus sectatrix</u>	Bermuda chub
<u>Labrisomus nuchipinnis</u>	Hairy blenny
<u>Lactophrys quadricornis</u>	Scrawled cowfish
<u>Lagocephalus laevigatus</u>	Smooth puffer
<u>Lagodon rhomboides</u>	Pinfish
<u>Larimus fasciatus</u>	Banded drum
<u>Leiostomus xanthurus</u>	Spot
<u>Lepisosteus oculatus</u>	Spotted gar
<u>Lepisosteus osseus</u>	Longnose gar
<u>Lepisosteus platostomus</u>	Shortnose gar
<u>Lepisosteus spatula</u>	Alligator gar
<u>Lepomis cyanellus</u>	Green sunfish
<u>Lepomis gulosus</u>	Warmouth
<u>Lepomis macrochirus</u>	Bluegill
<u>Lepomis megalotis</u>	Longear sunfish
<u>Lepomis microlophus</u>	Redear sunfish
<u>Lepophidium brevibarbe</u>	Blackedge cusk-eel
<u>Lobotes surinamensis</u>	Tripletail
<u>Lucania parva</u>	Rainwater killifish
<u>Lutjanus apodus</u>	Schoolmaster
<u>Lutjanus campechanus</u>	Red snapper
<u>Lutjanus griseus</u>	Gray snapper
<u>Lutjanus jocu</u>	Dog snapper
<u>Lutjanus synagris</u>	Lane snapper
<u>Lutjanus vivanus</u>	Silk snapper
<u>Megalops atlanticus</u>	Tarpon
<u>Membras martinica</u>	Rough silverside
<u>Menidia beryllina</u>	Inland silverside
<u>Menidia clarkhubbsi</u>	Texas silverside
<u>Menidia peninsulae</u>	Tidewater silverside
<u>Menticirrhus americanus</u>	Southern kingfish
<u>Menticirrhus littoralis</u>	Gulf kingfish
<u>Menticirrhus saxatilis</u>	Northern kingfish
<u>Microgobius gulosus</u>	Clown goby
<u>Microgobius thalassinus</u>	Green goby
<u>Micropogonias undulatus</u>	Atlantic croaker

Table A.9. (Cont'd.)

<u>Scientific Name</u>	<u>Common Name</u>
<b>Finfish (Cont'd.)</b>	
<u>Morone chrysops</u>	White bass
<u>Morone mississippiensis</u>	Yellow bass
<u>Morone saxatilis</u>	Striped bass
<u>Morone X</u>	Hybrid bass (striped x white)
<u>Mugil cephalus</u>	Striped mullet
<u>Mugil curema</u>	White mullet
<u>Mullus auratus</u>	Red goatfish
<u>Mustelus canis</u>	Smooth dogfish
<u>Mycteroperca bonaci</u>	Black grouper
<u>Mycteroperca microlepis</u>	Gag
<u>Mycteroperca phenax</u>	Scamp
<u>Mycteroperca rubra</u>	Comb grouper
<u>Myrophis punctatus</u>	Speckled worm eel
<u>Narcine brasiliensis</u>	Lesser electric ray
<u>Negaprion brevirostris</u>	Lemon shark
<u>Neomerinthe hemingwayi</u>	Spinycheek scorpionfish
<u>Ogcocephalus nasutus</u>	Shortnose batfish
<u>Ogcocephalus pantostictus</u>	Spotted batfish
<u>Ogcocephalus parvus</u>	Roughback batfish
<u>Ogcocephalus radiatus</u>	Polka-dot batfish
<u>Ogcocephalus sp.</u>	(Batfish-unidentified)
<u>Oligoplites saurus</u>	Leatherjacket
<u>Ophichthus gomesi</u>	Shrimp eel
<u>Ophichthus ophis</u>	Spotted snake eel
<u>Ophichthus puncticeps</u>	Palespotted eel
<u>Ophidion grayi</u>	Blotched cusk-eel
<u>Ophidion holbrookii</u>	Bank cusk-eel
<u>Ophidion marginatum</u>	Striped cusk-eel
<u>Ophidion welshi</u>	Crested cusk-eel
<u>Opisthonema oglinum</u>	Atlantic thread herring
<u>Opsanus beta</u>	Gulf toadfish
<u>Opsanus pardus</u>	Leopard toadfish
<u>Orthopristis chrysoptera</u>	Pigfish
<u>Parablennius marmoratus</u>	Seaweed blenny
<u>Paraconger caudilimbatus</u>	Margintail conger
<u>Paralichthys albigutta</u>	Gulf flounder
<u>Paralichthys lethostigma</u>	Southern flounder
<u>Paralichthys sp.</u>	(Flounder-unidentified)
<u>Paralichthys squamilentus</u>	Broad flounder
<u>Parasudis truculenta</u>	Longnose greeneye
<u>Peprilus alepidotus</u>	Harvestfish
<u>Peprilus burti</u>	Gulf butterflyfish
<u>Phaeoptyx conklini</u>	Freckled cardinalfish
<u>Physiculus fulvus</u>	Metallic codling
<u>Platybelone argalus</u>	Keeltail needlefish
<u>Poecilia formosa</u>	Amazon molly
<u>Poecilia latipinna</u>	Sailfin molly
<u>Pogonias cromis</u>	Black drum

Table A.9. (Cont'd.)

<u>Scientific Name</u>	<u>Common Name</u>
<b>Finfish (Cont'd.)</b>	
<u>Polydactylus octonemus</u>	Atlantic threadfin
<u>Pomacentrus fuscus</u>	Dusky damselfish
<u>Pomacentrus variabilis</u>	Cocoa damselfish
<u>Pomadasys crocro</u>	Burro grunt
<u>Pomatomus saltatrix</u>	Bluefish
<u>Pomoxis annularis</u>	White crappie
<u>Pomoxis nigromaculatus</u>	Black crappie
<u>Pontinus longispinis</u>	Longspine scorpionfish
<u>Porichthys plectrodon</u>	Atlantic midshipman
<u>Priacanthus arenatus</u>	Bigeye
<u>Prionotus longispinosus</u>	Bigeye searobin
<u>Prionotus martis</u>	Barred searobin
<u>Prionotus ophryas</u>	Bandtail searobin
<u>Prionotus paralatus</u>	Mexican searobin
<u>Prionotus roseus</u>	Bluespotted searobin
<u>Prionotus rubio</u>	Blackwing searobin
<u>Prionotus scitulus</u>	Leopard searobin
<u>Prionotus stearnsi</u>	Shortwing searobin
<u>Prionotus tribulus</u>	Bighead searobin
<u>Pristigenys alta</u>	Short bigeye
<u>Pristipomoides aquilonaris</u>	Wenchman
<u>Pristis pectinata</u>	Smalltooth sawfish
<u>Pylodictis olivaris</u>	Flathead catfish
<u>Rachycentron canadum</u>	Cobia
<u>Raja eglanteria</u>	Clearnose skate
<u>Raja texana</u>	Roundel skate
<u>Remora remora</u>	Remora
<u>Rhinobatos lentiginosus</u>	Atlantic guitarfish
<u>Rhinoptera bonasus</u>	Cownose ray
<u>Rhizoprionodon terraenovae</u>	Atlantic sharpnose shark
<u>Rhomboplites aurorubens</u>	Vermilion snapper
<u>Rypticus saponaceus</u>	Greater soapfish
<u>Sardinella aurita</u>	Spanish sardine
<u>Saurida brasiliensis</u>	Largescale lizardfish
<u>Saurida caribbaea</u>	Smallscale lizardfish
<u>Scartella cristata</u>	Molly miller
<u>Sciaenops ocellatus</u>	Red drum
<u>Scomber japonicus</u>	Chub mackerel
<u>Scomberomorus cavalla</u>	King mackerel
<u>Scomberomorus maculatus</u>	Spanish mackerel
<u>Scomberomorus sp.</u>	(Mackerel-unidentified)
<u>Scorpaena brasiliensis</u>	Barbfish
<u>Scorpaena calcarata</u>	Smoothhead scorpionfish
<u>Scorpaena plumieri</u>	Spotted scorpionfish
<u>Scyliorhinus retifer</u>	Chain dogfish
<u>Selar crumenophthalmus</u>	Bigeye scad
<u>Selene setapinnis</u>	Atlantic moonfish
<u>Selene vomer</u>	Lookdown

Table A.9. (Cont'd.)

<u>Scientific Name</u>	<u>Common Name</u>
<b>Finfish (Cont'd.)</b>	
<u>Seriola dumerili</u>	Greater amberjack
<u>Seriola fasciata</u>	Lesser amberjack
<u>Seriola zonata</u>	Banded rudderfish
<u>Serraniculus pumilio</u>	Pygmy sea bass
<u>Serranus atrobranchus</u>	Blackear bass
<u>Serranus phoebe</u>	Tattler
<u>Serranus subligarius</u>	Belted sandfish
<u>Sparisoma radians</u>	Bucktooth parrotfish
<u>Sphoeroides parvus</u>	Least puffer
<u>Sphoeroides spengleri</u>	Bandtail puffer
<u>Sphyraena barracuda</u>	Great barracuda
<u>Sphyraena quachancho</u>	Guaguanche
<u>Sphyrna lewini</u>	Scalloped hammerhead
<u>Sphyrna mokarran</u>	Great hammerhead
<u>Sphyrna tiburo</u>	Bonnethead
<u>Sphyrna tudes</u>	Smalleye hammerhead
<u>Stellifer lanceolatus</u>	Star drum
<u>Stenotomus caprinus</u>	Longspine porgy
<u>Strongylura marina</u>	Atlantic needlefish
<u>Strongylura timucu</u>	Timucu
<u>Syacium gunteri</u>	Shoal flounder
<u>Syacium papillosum</u>	Dusky flounder
<u>Symphurus civitatus</u>	Offshore tonguefish
<u>Symphurus diomedianus</u>	Spottedfin tonguefish
<u>Symphurus parvus</u>	Pygmy tonguefish
<u>Symphurus plagiusa</u>	Blackcheek tonguefish
<u>Symphurus urospilus</u>	Spottail tonguefish
<u>Syngnathus floridae</u>	Dusky pipefish
<u>Syngnathus louisianae</u>	Chain pipefish
<u>Syngnathus pelagicus</u>	Sargassum pipefish
<u>Syngnathus scovelli</u>	Gulf pipefish
<u>Synodus foetens</u>	Inshore lizardfish
<u>Synodus poeyi</u>	Offshore lizardfish
<u>Thunnus thynnus</u>	Bluefin tuna
<u>Tilapia aurea</u>	Blue tilapia
<u>Trachinocephalus myops</u>	Snakefish
<u>Trachinotus carolinus</u>	Florida pompano
<u>Trachinotus falcatus</u>	Permit
<u>Trachinotus goodei</u>	Palometa
<u>Trachurus lathami</u>	Rough scad
<u>Trichiurus lepturus</u>	Atlantic cutlassfish
<u>Trinectes maculatus</u>	Hogchoker
<u>Umbrina coroides</u>	Sand drum
<u>Upeneus parvus</u>	Dwarf goatfish
<u>Urophycis cirrata</u>	Gulf hake
<u>Urophycis floridana</u>	Southern hake
<u>Xanthichthys ringens</u>	Sargassum triggerfish



Table A.9. (Cont'd.)

Scientific Name	Common Name
<b>Invertebrates</b>	
<u>Acetes americanus</u>	(Sergestid shrimp)
<u>Agriopoma texasianum</u>	Texas venus
<u>Albunea gibbesii</u>	Surf mole crab
<u>Albunea paretii</u>	Beach mole crab
<u>Alpheua formosus</u>	Striped snapping shrimp
<u>Alpheus estuariensis</u>	Estuarine snapping shrimp
<u>Amaea mitchelli</u>	Mitchell's wentletrap
<u>Anachis avara</u>	Greedy dovesnail
<u>Anadara brasiliana</u>	Incongruous ark
<u>Anadara floridana</u>	Cut-ribbed ark
<u>Anadara ovalis</u>	Blood ark
<u>Anadara transversa</u>	Transverse ark
<u>Anasimus latus</u>	Stilt spider crab
<u>Anomalocardia auberiana</u>	Pointed-venus
<u>Anomia simplex</u>	Common jingle
<u>Aplysia brasiliana</u>	Sooty seahare
<u>Arbacia punctulata</u>	Red sea urchin
<u>Arca imbricata</u>	Mossy ark
<u>Architectonica nobilis</u>	Common sundial
<u>Arcinella cornuta</u>	Florida spiny jewelbox
<u>Arenaeus cribrarius</u>	Speckled swimming crab
<u>Argopecten gibbus</u>	Atlantic calico scallop
<u>Argopecten irradians</u>	Bay scallop
<u>Armina tigrina</u>	Tiger armina
<u>Astropecten duplicatus</u>	Two-spined starfish
<u>Atrina serrata</u>	Sawtooth pen shell
<u>Aurelia aurita</u>	Moon jellyfish
<u>Barbatia candida</u>	White-beard ark
<u>Beroe ovata</u>	Sea walnut
<u>Brachidontes exustus</u>	Scorched mussel
<u>Brissopsis alta</u>	Heart urchin
<u>Bulla striata</u>	Striate bubble
<u>Bursatella leachii pleii</u>	Ragged seahare
<u>Busycon sinistrum</u>	Lightning whelk
<u>Busycotypus spiratus</u>	Pearwhelk
<u>Calappa flammea</u>	Flame box crab
<u>Calappa ocellata</u>	Ocellate box crab
<u>Calappa sulcata</u>	Yellow box crab
<u>Callianassa louisianensis</u>	Gulf estuarine ghost shrimp
<u>Callinectes marginatus</u>	(Sargassum crab)
<u>Callinectes sapidus</u>	Blue crab
<u>Callinectes similis</u>	Lesser blue crab
<u>Cancellaria reticulata</u>	Common nutmeg
<u>Cantharus cancellarius</u>	Cancellate cantharus
<u>Cerithidea pliculosa</u>	Plicate hornsnail
<u>Cerithium lutosum</u>	Variable cerith
<u>Chasmocarcinus mississippiensis</u>	Roughwrist soft crab
<u>Chione cancellata</u>	Cross-barred venus

Table A.9. (Cont'd.)

Scientific Name	Common Name
<b>Invetebrates (Cont'd.)</b>	
<u>Chione clenchi</u>	Clench venus
<u>Chione intapurpurea</u>	Lady-in-waiting venus
<u>Chiropsalmus quadrumanus</u>	Sea wasp
<u>Chrysaora quinquecirrha</u>	Sea nettle
<u>Clibanarius vittatus</u>	Thinstripe hermit
<u>Crassostrea virginica</u>	Eastern oyster
<u>Crepidula convexa</u>	Convex slippersnail
<u>Crepidula fornicata</u>	Common Atlantic slippersnail
<u>Crepidula plana</u>	Eastern white slippersnail
<u>Cyclinella tenuis</u>	Thin cyclinella
<u>Cyrtopleura costata</u>	Angelwing
<u>Dardanus fucosus</u>	Bareye hermit
<u>Dinocardium robustum</u>	Atlantic giant-cockle
<u>Distorsio clathrata</u>	Atlantic distorsio
<u>Donax variabilis</u>	Variable coquina
<u>Dosinia discus</u>	Disk dosinia
<u>Dromidia antillensis</u>	Hairy sponge crab
<u>Dyspanopeus texana</u>	Gulf grassflat crab
<u>Echinometra lucunter</u>	Rock-boring urchin
<u>Emerita portoricensis</u>	Puerto Rican sand crab
<u>Ensis minor</u>	Minor jackknife
<u>Euceramus praelongus</u>	Olivepit porcelain crab
<u>Eurypanopeus abbreviatus</u>	Lobate mud crab
<u>Eurypanopeus depressus</u>	Flatback mud crab
<u>Exhippolysmata oplophoroides</u>	Redleg humpback shrimp
<u>Fasciolaria liliium liliium</u>	Banded tulip
<u>Glypturus acanthochirus</u>	Ghost shrimp
<u>Haminoea antillarum</u>	Antilles glassy-bubble
<u>Haminoea succinea</u>	Amber glassy-bubble
<u>Hepatus epheliticus</u>	Calico box crab
<u>Hepatus pudibundus</u>	Flecked box crab
<u>Heterocrypta granulata</u>	Smooth elbow crab
<u>Hexapanopeus angustifrons</u>	Smooth mud crab
<u>Hexapanopeus paulensis</u>	Knobbed mud crab
<u>Hypoconcha arcuata</u>	Granulate shellback shrimp
<u>Hypoconcha sabulosa</u>	Shellback crab (Dromiid)
<u>Ischadium recurvum</u>	Hooked mussel
<u>Isocheles wurdemanni</u>	Surf hermit
<u>Laevicardium mortoni</u>	Morton eggcockle
<u>Latreutes fucorum</u>	Slender sargassum shrimp
<u>Latreutes parvulus</u>	Sargassum shrimp
<u>Leander tenuicornis</u>	Brown grass shrimp
<u>Leiolambrus nitidus</u>	White elbow crab
<u>Lepidopa benedicti</u>	(Gulf mole crab)
<u>Libinia dubia</u>	Longnose spider crab
<u>Libinia emarginata</u>	Portly spider crab
<u>Littorina irrorata</u>	Marsh periwinkle
<u>Loligo pealeii</u>	Longfin squid

Table A.9. (Cont'd.)

Scientific Name	Common Name
<b>Invertebrates (Cont'd.)</b>	
<u>Loligo pleii</u>	Arrow squid
<u>Loliguncula brevis</u>	Atlantic brief squid
<u>Lucifer faxoni</u>	Sergestid shrimp
<u>Lucina pectinata</u>	Thick lucine
<u>Luidia alternata</u>	Banded sea star
<u>Luidia clathrata</u>	Large sea star
<u>Lysiosquilla scabricauda</u>	(Giant) mantis shrimp
<u>Lysmata wurdemanni</u>	Peppermint shrimp
<u>Lytechinus variegatus</u>	Short spined sea urchin
<u>Macoma brevifrons</u>	Short macoma
<u>Macrobrachium acanthurus</u>	Cinnamon river shrimp
<u>Macrobrachium ohione</u>	Ohio shrimp
<u>Macrocallista maculata</u>	Calico clam
<u>Mactra fragilis</u>	Fragile Atlantic mactra
<u>Melampus bidentatus</u>	Eastern melampus
<u>Mellita quinquiesperforata</u>	Five-lunuled sand dollar
<u>Menippe adina</u>	Gulf stone crab
<u>Mercenaria campechiensis</u>	Southern quahog
<u>Mercenaria campechiensis texana</u>	Texas quahog
<u>Metoporphaphis calcarata</u>	False arrow crab
<u>Mnemiopsis mccradyi</u>	Phosphorus jelly
<u>Molgula manhattensis</u>	Sea squirt
<u>Mulinia lateralis</u>	Dwarf surf clam
<u>Muricanthus fluvescens</u>	Giant eastern murex
<u>Nassarius vibex</u>	Bruised nassa
<u>Nemopsis bachei</u>	(Hydromedusa)
<u>Neritina virginea</u>	Virgin nerite
<u>Neverita duplicata</u>	Shark eye
<u>Noetia ponderosa</u>	Ponderous ark
<u>Octopus vulgaris</u>	Common octopus
<u>Oculina diffusa</u>	Ivory coral
<u>Ocypode quadrata</u>	Atlantic ghost crab
<u>Oliva sayana</u>	Lettered olive
<u>Ophiolepis elegans</u>	Brittle star
<u>Orchestia grillus</u>	Beach flea (amphipod)
<u>Ostreola equestris</u>	Crested oyster
<u>Ovalipes floridanus</u>	Florida lady crab
<u>Paguristes hummi</u>	(Blue spot hermit crab)
<u>Pagurus annulipes</u>	(Brown-banded hermit crab)
<u>Pagurus brevidactylus</u>	Short-fingered hermit
<u>Pagurus impressus</u>	Dimpled hermit
<u>Pagurus longicarpus</u>	Longwrist hermit
<u>Pagurus pollicaris</u>	Flatclaw hermit
<u>Palaemonetes pugio</u>	Daggerblade grass shrimp
<u>Palaemonetes vulgaris</u>	Marsh grass shrimp
<u>Panopeus simpsoni</u>	Oystershell mud crab
<u>Paranthus rapiformis</u>	Onion anemone
<u>Parthenope serrata</u>	Sawtooth elbow crab

Table A.9. (Cont'd.)

Scientific Name	Common Name
<b>Inveterbrates (Cont'd.)</b>	
<u>Pelia mutica</u>	Cryptic teardrop crab
<u>Penaeus aztecus</u>	Brown shrimp
<u>Penaeus duorarum</u>	Pink shrimp
<u>Penaeus setiferus</u>	White shrimp
<u>Persephona crinita</u>	Pink purse crab
<u>Persephona mediterranea</u>	Mottled purse crab
<u>Petrochirus diogenes</u>	Giant hermit
<u>Petrolisthes armatus</u>	Green porcelain crab
<u>Phalium granulatum</u>	Scotch bonnet
<u>Physalia physalis</u>	Portuguese man-of-war
<u>Pinnotheres maculatus</u>	Squatter pea crab
<u>Pleurobranchaea tarda</u>	(Side-gilled slug)
<u>Pleuroploca gigantea</u>	Horse conch
<u>Podochela riisei</u>	Longfinger neck crab
<u>Podochela sidneyi</u>	Shortfinger neck crab
<u>Polycera hummi</u>	Humm's polycera
<u>Polymesoda maritima</u>	Southern marshclam
<u>Porcellana sayana</u>	Spotted porcelain crab
<u>Porcellana sigsbeiana</u>	Striped porcelain crab
<u>Portunus anceps</u>	Delicate swimming crab
<u>Portunus gibbesii</u>	Iridescent swimming crab
<u>Portunus sayi</u>	Sargassum swimming crab
<u>Portunus spinicarpus</u>	Longspine swimming crab
<u>Portunus spinimanus</u>	Blotched swimming crab
<u>Portunus ventralis</u>	(Portunid swimming crab)
<u>Procambarus clarkii</u>	Red swamp crawfish
<u>Pseudocyphoma intermedium</u>	Intermediate cyphoma
<u>Rangia cuneata</u>	Atlantic rangia
<u>Rangia flexuosa</u>	Brown rangia
<u>Raninoides louisianensis</u>	Gulf frog crab
<u>Renilla mulleri</u>	Sea pansy
<u>Rhithropanopeus harrisii</u>	Harris mud crab
<u>Scyllaea pelagica</u>	Sargassum nudibranch
<u>Sesarma reticulatum</u>	Heavy marsh crab
<u>Sicyonia brevirostris</u>	Brown rock shrimp
<u>Sicyonia dorsalis</u>	Lesser rock shrimp
<u>Sicyonia stimpsoni</u>	Eyespot rock shrimp
<u>Sicyonia typica</u>	Kinglet rock shrimp
<u>Simnialena marferula</u>	Sea-whip simnia
<u>Sinum perspectivum</u>	White baby-ear
<u>Solariorbis blake</u>	(Vitrinella)
<u>Solenocera vioscai</u>	Humpback shrimp
<u>Speocarcinus lobatus</u>	Gulf squareback crab
<u>Spisula lidissima</u>	Atlantic surfclam
<u>Squilla chydrea</u>	(Offshore mantis shrimp)
<u>Squilla empusa</u>	Mantis shrimp
<u>Squilla neglecta</u>	Lesser mantis shrimp
<u>Stenorhynchus seticornis</u>	Yellowline arrow crab
<u>Stomolophus meleagris</u>	Cabbagehead
<u>Strombus alatus</u>	Florida fighting conch

Table A.9. (Cont'd.)

<u>Scientific Name</u>	<u>Common Name</u>
<b>Inveterbrates</b> (Cont'd.)	
Suborder Zygoptera	(Damselfly nymphs)
<u>Synalpheus fritzmuelleri</u>	Speckled snapping shrimp
<u>Tagelus plebeius</u>	Stout tagelus
<u>Tellina alternata</u>	Alternate tellin
<u>Tellina tampaensis</u>	Tampa tellin
<u>Terebra protexta</u>	Fine-ribbed auger
<u>Thais haemastoma floridana</u>	Florida rocksnail
<u>Thyone mexicana</u>	Sea cucumber
<u>Tonna galea</u>	Giant tun
<u>Tozeuma carolinense</u>	Arrow shrimp
<u>Trachycardium muricatum</u>	Yellow pricklycockle
<u>Trachypenaeus constrictus</u>	Roughneck shrimp
<u>Trachypenaeus similis</u>	Roughback shrimp
<u>Uca panacea</u>	Gulf sand fiddler
<u>Upogebia affinis</u>	Coastal mud shrimp
<u>Velella velella</u>	By-the-wind sailor
<u>Xiphopenaeus kroyeri</u>	Seabob

Appendix B. Summary of hydrological data collected for gill net, bay and beach bag seine, oyster dredge, bay and gulf trawl, beach seine, and ICWW trawl samples.

Table B.1. Mean surface salinity (o/oo) at sampled gill net sites by bay system during spring and fall, 1975-95. ND = no data.

Year	Sabine		East		San		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
	Lake	Galveston	Matagorda	Matagorda	Antonio	Aransas	Christi	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Coastwide
	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall
1975	ND	ND	ND	ND	ND	17.6	ND	18.5	ND	20.0	ND	33.3	ND	25.7
1976	ND	ND	ND	18.8	ND	17.9	ND	10.9	ND	14.9	ND	26.0	ND	23.2
1977	ND	15.4	23.2	14.2	18.6	19.2	15.0	9.0	19.1	18.2	30.9	26.1	28.5	30.5
1978	ND	18.5	21.3	20.8	18.4	19.2	15.6	19.0	12.5	26.5	26.2	38.2	39.3	18.2
1979	ND	7.6	13.3	14.0	11.8	11.1	9.6	7.5	12.3	17.4	19.7	30.0	28.2	31.8
1980	ND	11.3	22.6	17.0	24.1	14.3	20.8	18.2	9.4	18.2	23.4	37.3	24.6	26.0
1981	ND	25.8	10.3	26.8	17.5	20.1	13.6	19.0	10.8	20.2	8.4	29.4	30.6	30.8
1982	ND	12.1	20.5	18.3	24.1	12.4	23.0	17.3	26.9	12.1	25.1	23.6	24.0	31.5
1983	ND	14.8	11.4	17.5	13.4	20.1	12.7	19.5	17.3	21.6	7.8	29.3	32.8	36.0
1984	ND	21.4	19.0	23.1	15.8	23.8	19.0	27.4	29.6	22.1	26.8	30.2	33.7	17.4
1985	ND	18.0	22.3	14.7	23.5	11.0	23.3	12.8	23.7	13.4	24.2	22.3	35.1	31.2
1986	11.7	13.1	15.0	20.9	25.3	14.1	23.9	22.3	21.9	22.9	21.4	24.4	41.7	32.3
1987	8.2	14.3	19.7	21.5	15.8	13.6	16.1	20.4	12.3	16.1	16.7	13.5	32.8	38.2
1988	7.8	12.1	18.3	21.8	24.9	27.3	25.4	32.4	23.8	23.0	21.3	24.8	33.6	34.1
1989	5.5	8.7	15.9	14.8	26.0	26.3	26.5	28.4	26.5	29.9	30.8	34.3	35.3	30.9
1990	2.0	10.4	12.4	19.3	19.2	27.8	19.6	25.3	23.7	24.3	27.0	22.2	31.5	38.3
1991	0.2	5.4	9.4	17.4	11.7	19.4	11.2	19.5	16.3	25.1	16.9	18.4	26.9	31.2
1992	2.0	12.1	10.4	22.4	21.5	23.4	5.7	23.1	2.7	20.9	4.1	17.6	16.7	39.0
1993	2.1	8.3	12.1	21.2	11.5	26.0	10.8	24.5	9.2	17.5	10.2	18.7	26.5	30.5
1994	1.4	5.1	11.3	12.3	21.4	24.3	18.2	18.8	12.2	18.5	17.3	22.4	27.7	32.0
1995	0.2	8.0	9.7	19.7	15.7	24.3	13.5	21.9	17.0	23.5	18.5	22.8	25.8	34.5

Table B.2. Mean surface water temperature (C) at sampled gill net sites by bay system during spring and fall, 1975-95. ND = no data.

Year	Sabine		East		San		Corpus Christi		Upper Laguna Madre		Lower Laguna Madre		Coastwide	
	Lake	Galveston	Matagorda	Matagorda	Antonio	Aransas	Christi	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Coastwide
	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall
1975	ND	ND	20.7	ND	ND	21.2	ND	17.4	ND	23.9	ND	23.0	ND	24.4
1976	ND	30.0	18.2	ND	ND	24.8	ND	24.0	ND	24.2	ND	19.6	ND	20.8
1977	ND	24.9	20.6	25.0	21.3	25.2	23.1	25.6	22.7	25.5	23.3	26.4	21.3	26.6
1978	ND	26.5	21.5	25.6	24.2	25.8	24.1	25.1	24.2	27.3	23.5	26.4	23.2	24.1
1979	ND	26.5	22.8	27.4	23.4	27.3	23.6	27.3	24.2	27.1	24.5	28.1	25.0	24.6
1980	ND	25.9	24.4	25.9	23.5	26.0	25.6	26.8	24.6	27.0	25.2	29.0	27.0	27.4
1981	ND	27.1	25.3	27.3	23.1	26.0	24.6	27.4	25.0	27.3	25.2	26.9	26.9	26.3
1982	ND	26.1	24.6	26.9	25.1	27.2	24.6	25.7	25.6	26.3	24.1	27.8	24.9	25.8
1983	ND	25.7	25.3	25.8	25.9	25.0	25.5	26.2	25.2	26.6	25.3	27.4	27.0	26.8
1984	ND	26.7	25.0	25.7	27.2	25.1	25.3	25.8	25.2	26.2	25.0	27.3	25.9	26.8
1985	ND	27.9	25.5	28.6	25.6	27.4	25.0	27.5	25.8	26.3	26.0	27.6	28.2	27.4
1986	26.8	26.3	26.4	25.1	27.0	26.3	25.4	27.2	25.3	24.5	26.3	26.6	26.6	26.9
1987	25.7	24.0	26.4	24.0	27.1	24.5	26.4	26.7	26.3	24.8	24.8	27.6	26.2	25.2
1988	25.4	26.2	25.3	25.8	26.2	26.3	25.0	26.9	25.9	24.4	25.8	26.6	27.4	25.2
1989	25.0	24.8	25.7	24.0	28.7	25.6	26.4	24.1	26.6	24.1	26.8	26.5	26.4	27.0
1990	23.3	25.7	24.6	23.8	27.8	25.5	26.8	24.7	27.0	24.9	25.7	27.7	27.0	26.3
1991	27.0	24.9	24.3	23.8	27.8	24.1	26.5	23.6	27.0	23.2	28.1	25.0	28.3	25.8
1992	27.5	23.5	26.3	23.0	24.8	24.3	24.6	24.3	27.3	24.7	26.8	28.0	25.3	27.4
1993	25.9	22.9	25.8	22.4	26.9	24.1	25.6	23.2	26.1	24.0	25.8	23.7	28.0	26.7
1994	27.1	25.1	26.7	24.9	27.1	26.3	26.6	26.2	25.4	26.0	27.2	26.5	27.9	27.8
1995	26.2	24.8	26.2	24.7	26.7	25.1	25.9	25.9	26.9	25.5	26.4	26.4	27.3	26.6

Table B.3. Annual mean surface turbidity at sampled gill net sites by bay system during spring and fall 1975-95. ND = no data.

Year	Sabine Lake		Galveston		East Matagorda		Matagorda		San Antonio		Arkansas		Corpus Christi		Upper Laquana Madre		Lower Laquana Madre		Coastwide	
	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall
Jackson Turbidity Units																				
1975	ND	ND	53	ND	ND	ND	30	ND	42	ND	24	ND	27	ND	42	ND	28	ND	37	37
1976	ND	ND	52	ND	157	ND	33	ND	25	ND	63	ND	60	ND	51	ND	38	ND	79	50
1977	ND	ND	80	75	118	46	67	48	13	41	52	169	47	34	39	40	31	64	50	50
1978	ND	ND	44	44	36	15	68	74	20	55	50	61	47	66	67	37	39	54	48	48
1979	ND	ND	153	72	38	28	74	66	80	22	70	42	67	39	32	34	83	80	55	55
1980	ND	ND	99	69	67	49	74	33	17	19	53	40	51	36	57	55	64	71	64	48
1981	ND	ND	68	68	62	64	82	64	81	21	43	58	67	39	185	45	87	66	84	55
1982	ND	ND	66	56	82	55	75	47	35	27	91	33	49	38	63	32	113	79	72	47
1983	ND	ND	57	63	61	27	50	40	41	32	49	38	41	42	50	40	59	72	51	48
1984	ND	ND	43	34	27	25	35	44	47	40	40	39	47	38	69	56	113	90	54	47
1985	ND	ND	26	28	59	37	52	51	57	49	46	39	57	41	72	41	98	56	55	42
1986	43	28	32	35	64	37	60	31	46	32	38	41	57	26	61	85	53	59	48	43
Nephelometric Units																				
1987	30	18	17	42	19	19	28	19	26	15	10	7	22	7	14	11	23	13	21	14
1988	21	11	16	11	29	19	16	19	22	21	13	15	24	10	18	14	26	29	19	17
1989	25	9	12	9	16	22	36	15	30	12	22	8	18	12	12	9	45	13	24	11
1990	16	8	9	13	23	13	26	15	38	15	21	13	16	11	24	11	29	14	22	13
1991	15	6	20	8	52	21	29	15	19	13	23	13	13	12	25	18	13	10	21	12
1992	20	11	10	22	13	46	17	52	14	41	14	14	23	11	25	12	22	16	32	13
1993	24	11	24	16	35	14	46	16	29	11	24	15	17	14	18	15	23	14	27	14
1994	13	13	19	23	23	15	27	26	19	9	21	10	18	9	12	11	25	15	20	16
1995	25	8	39	11	38	20	38	20	23	8	21	13	27	13	16	9	37	12	30	13



Table B.4. Annual mean surface salinity (o/oo) at sampled bag seine sites by bay system during 1977-95. ND = no data.

Year	East					Corpus Christi					Lower Coastwide				
	Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Christi	Laquna Madre	Laquna Madre	Coastwide	Upper	Lower	Laquna Madre	Laquna Madre	Coastwide
1977	ND	21.9	ND	17.6	17.7	20.9	33.8	39.8	33.0	33.0	39.8	33.0	33.0	25.4	
1978	ND	21.8	ND	19.7	20.6	19.9	29.5	39.6	29.2	29.2	39.6	29.2	29.2	25.0	
1979	ND	12.2	ND	11.4	11.8	11.1	23.9	31.9	27.3	27.3	31.9	27.3	27.3	17.4	
1980	ND	20.9	ND	19.9	21.0	19.8	28.1	29.6	28.8	28.8	29.6	28.8	28.8	23.4	
1981	ND	18.2	ND	19.2	15.6	12.1	25.0	26.0	28.3	28.3	26.0	28.3	28.3	20.1	
1982	ND	15.9	ND	18.2	17.0	17.6	27.6	29.8	29.7	29.7	29.8	29.7	29.7	21.3	
1983	ND	12.2	15.4	16.5	17.3	16.8	27.5	36.4	31.7	31.7	36.4	31.7	31.7	21.2	
1984	ND	19.5	17.8	21.6	23.2	22.6	31.8	39.5	29.9	29.9	39.5	29.9	29.9	25.5	
1985	ND	17.0	16.9	19.7	17.5	19.7	28.1	36.7	32.1	32.1	36.7	32.1	32.1	23.2	
1986	10.1	16.1	20.1	19.8	17.0	23.5	32.6	39.7	34.9	34.9	39.7	34.9	34.9	24.2	
1987	7.6	18.1	15.3	15.4	10.8	13.7	28.7	31.4	31.5	31.5	31.4	31.5	31.5	19.9	
1988	7.7	20.2	26.5	27.4	22.6	24.3	35.2	44.9	31.9	31.9	44.9	31.9	31.9	27.4	
1989	6.6	15.1	26.9	26.9	27.4	31.4	35.6	48.6	34.2	34.2	48.6	34.2	34.2	28.5	
1990	6.4	16.9	23.6	23.6	24.8	26.7	32.4	47.7	35.8	35.8	47.7	35.8	35.8	27.2	
1991	2.6	12.4	17.3	16.7	19.3	17.7	30.8	40.0	28.8	28.8	40.0	28.8	28.8	21.1	
1992	5.3	15.2	15.4	13.5	9.4	10.7	22.4	25.3	28.7	28.7	25.3	28.7	28.7	16.8	
1993	4.3	12.6	18.2	17.1	13.9	13.8	27.6	27.7	27.5	27.5	27.7	27.5	27.5	18.3	
1994	4.3	13.5	22.7	18.8	15.4	21.5	30.3	34.7	30.3	30.3	34.7	30.3	30.3	21.5	
1995	3.4	14.6	19.6	18.5	18.8	21.1	28.9	37.5	31.9	31.9	37.5	31.9	31.9	22.3	

Table B.5. Annual mean surface temperature (C) at sampled bag seine sites by bay system during 1977-95. ND = no data.

Year	East					Corpus Christi					Lower Coastwide				
	Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Christi	Laquna Madre	Laquna Madre	Coastwide	Upper	Lower	Laquna Madre	Laquna Madre	Coastwide
1977	ND	20.3	ND	20.9	21.7	20.8	20.4	20.6	20.5	20.5	20.6	20.5	20.5	20.7	
1978	ND	21.4	ND	20.2	21.6	22.3	21.3	22.3	22.4	22.4	22.3	22.4	22.4	21.6	
1979	ND	22.8	ND	22.8	23.3	23.2	23.6	21.8	23.1	23.1	21.8	23.1	23.1	22.9	
1980	ND	23.9	ND	21.9	23.2	23.6	23.4	24.6	24.3	24.3	24.6	24.3	24.3	23.5	
1981	ND	22.5	ND	21.5	22.4	23.7	22.6	24.1	24.6	24.6	24.1	24.6	24.6	23.0	
1982	ND	23.9	ND	23.3	23.1	24.2	23.4	24.1	23.9	23.9	24.1	23.9	23.9	23.7	
1983	ND	24.0	23.6	21.9	21.7	24.3	24.3	25.4	24.9	24.9	25.4	24.9	24.9	23.8	
1984	ND	23.9	22.3	22.5	21.9	24.0	23.3	24.0	24.2	24.2	24.0	24.2	24.2	23.4	
1985	ND	24.4	24.4	23.5	24.0	23.9	23.5	23.5	24.4	24.4	23.5	24.4	24.4	24.0	
1986	23.7	24.2	23.4	23.3	23.5	25.2	23.6	24.5	25.0	25.0	24.5	25.0	25.0	24.2	
1987	22.0	22.8	23.8	23.4	22.2	23.1	24.1	24.2	23.8	23.8	24.2	23.8	23.8	23.2	
1988	21.7	23.4	23.9	23.4	21.1	24.3	23.3	23.9	25.1	25.1	23.9	25.1	25.1	23.5	
1989	21.4	23.1	22.9	22.3	23.0	22.8	24.3	25.0	23.4	23.4	25.0	23.4	23.4	23.4	
1990	21.7	22.6	24.7	23.6	23.0	24.4	24.9	24.9	25.5	25.5	24.9	25.5	25.5	23.9	
1991	22.9	22.3	24.5	22.2	23.2	23.1	24.8	25.0	23.5	23.5	25.0	23.5	23.5	23.5	
1992	22.2	21.7	22.2	21.4	23.3	22.6	23.4	24.3	25.9	25.9	24.3	25.9	25.9	23.0	
1993	22.4	22.2	24.3	22.9	22.5	23.9	23.6	23.4	25.1	25.1	23.4	25.1	25.1	23.2	
1994	23.1	23.1	24.5	23.3	22.8	25.0	25.0	24.6	25.7	25.7	24.6	25.7	25.7	24.1	
1995	22.2	23.3	25.0	23.9	24.4	24.3	25.2	24.7	25.8	25.8	24.7	25.8	25.8	24.3	

Table B.6. Annual mean surface turbidity at sampled bag seine sites by bay system during 1977-95. ND = no data.

Year	East			San Antonio	Aransas	Upper		Lower	
	Sabine Lake	Galveston	Mataforda			Christi	Laguna Madre	Laguna Madre	Coastwide
Jackson Turbidity Units									
1977	ND	94	ND	60	50	40	50	30	55
1978	ND	78	ND	55	41	43	51	34	51
1979	ND	90	ND	70	53	44	47	59	60
1980	ND	90	ND	42	47	52	75	73	61
1981	ND	87	ND	54	65	44	107	95	71
1982	ND	105	ND	50	31	60	69	87	69
1983	ND	96	88	54	51	46	57	48	58
1984	ND	79	42	41	48	41	82	61	57
1985	ND	52	67	45	47	40	108	68	59
1986	46	84	59	46	46	44	60	80	61
Nephelemetric Units									
1987	24	28	39	36	9	26	15	17	24
1988	26	26	28	29	28	20	22	24	26
1989	25	29	26	25	22	20	22	22	26
1990	21	29	26	30	23	21	20	23	26
1991	28	25	32	33	42	17	21	15	26
1992	24	23	34	41	31	21	17	25	29
1993	21	30	27	38	30	23	26	36	30
1994	21	24	31	26	17	17	10	40	24
1995	27	30	33	32	27	17	12	25	26

Table B.7. Annual mean bottom salinity (o/oo) at sampled oyster dredge "reef" sites in Texas bay systems from 1984-95. ND = no data.

Year	East			San Antonio	Aransas	Coastwide	
	Galveston	Mataforda	Mataforda			Aransas	Coastwide
1984	16.7	ND	ND	ND	ND	16.7	16.7
1985	17.6	ND	ND	ND	ND	17.6	17.6
1986	15.5	22.0	18.2	21.0	18.9	18.9	18.9
1987	16.3	16.6	10.9	14.2	14.5	14.5	14.5
1988	19.6	28.1	22.9	25.0	23.7	23.7	23.7
1989	16.0	29.2	27.9	29.7	25.1	25.1	25.1
1990	16.0	24.4	24.1	26.2	22.3	22.3	22.3
1991	12.3	17.4	19.5	18.6	16.7	16.7	16.7
1992	14.9	11.8	9.2	8.7	11.4	11.4	11.4
1993	13.5	15.9	13.2	14.5	14.2	14.2	14.2
1994	13.7	19.4	17.4	19.8	16.8	16.8	16.8
1995	14.7	17.8	18.7	20.2	17.1	17.1	17.1

Table B.8. Annual mean bottom temperature (C) at sampled oyster dredge "reef" sites in Texas bay systems from 1984-94. ND = no data

Year	Galveston	Matagorda	San Antonio	Aransas	Coastwide
1984	21.0	ND	ND	ND	20.9
1985	22.0	ND	ND	ND	22.0
1986	22.8	22.4	22.3	22.1	22.4
1987	21.2	22.2	21.4	19.9	21.3
1988	21.6	21.8	21.6	22.0	21.7
1989	20.9	20.8	21.6	20.4	21.0
1990	21.7	22.6	22.6	23.0	22.4
1991	21.6	21.9	21.8	21.3	21.7
1992	21.8	20.8	22.6	21.4	21.7
1993	21.4	22.2	21.9	21.0	21.6
1994	22.0	22.5	23.3	21.4	22.2
1995	21.4	22.4	23.4	22.6	22.2

Table B.9. Annual mean bottom turbidity at sampled oyster dredge "reef" sites" in Texas bay systems from 1984-95. ND = no data.

Year	Galveston	Matagorda	San Antonio	Aransas	Coastwide
<b>Jackson Turbidity Units</b>					
1984	25	ND	ND	ND	25
1985	47	ND	ND	ND	47
1986	40	51	48	37	45
<b>Nephelometric Units</b>					
1987	14	22	30	8	20
1988	15	21	16	16	17
1989	19	20	27	16	21
1990	14	22	26	16	20
1991	16	23	23	20	21
1992	15	32	37	31	26
1993	21	24	20	22	22
1994	17	18	16	16	17
1995	19	20	16	15	18

Table B.10. Annual mean bottom salinity (o/oo) at sampled bay trawl sites in Texas bay systems from 1977-95. ND = no data.

Year	East						Corpus Christi			Upper Laguna Madre			Lower Laguna Madre			Coastwide
	Sabine Lake	Galveston	Matacorda	Matacorda	San Antonio	Aransas	Christi	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Coastwide	
1977	ND	20.5	ND	17.9	13.9	19.5	ND	ND	ND	ND	ND	ND	ND	ND	18.5	
1978	ND	20.1	ND	19.3	14.7	20.6	ND	ND	ND	ND	ND	ND	ND	ND	19.0	
1979	ND	9.0	ND	10.3	5.7	ND	ND	ND	ND	ND	ND	ND	ND	ND	8.8	
1980	ND	22.8	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	22.8	
1981	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
1982	ND	16.0	ND	22.4	16.3	19.2	30.3	34.1	35.8	34.1	35.8	35.8	35.8	35.8	21.3	
1983	ND	10.7	ND	20.4	16.9	19.6	29.8	36.9	33.0	36.9	33.0	36.9	33.0	36.9	19.1	
1984	ND	18.5	ND	25.2	22.9	25.2	32.5	40.0	31.0	40.0	31.0	40.0	31.0	40.0	24.6	
1985	ND	17.0	ND	21.0	16.2	21.2	29.8	37.3	33.1	37.3	33.1	37.3	33.1	37.3	21.5	
1986	7.8	14.8	ND	24.5	17.3	22.7	31.1	39.6	36.1	39.6	36.1	39.6	36.1	39.6	21.6	
1987	7.3	15.1	16.7	20.6	9.9	18.1	27.5	31.9	33.3	31.9	33.3	33.3	33.3	33.3	18.6	
1988	7.8	19.2	28.7	29.6	21.7	25.7	34.9	45.0	34.8	45.0	34.8	45.0	34.8	45.0	25.6	
1989	6.2	16.4	27.6	30.2	26.8	30.4	35.4	49.3	35.9	49.3	35.9	49.3	35.9	49.3	26.1	
1990	5.7	15.1	25.8	26.1	21.6	27.0	32.0	48.6	36.3	48.6	36.3	48.6	36.3	48.6	23.4	
1991	2.2	11.9	18.7	20.4	17.7	20.0	29.9	41.4	31.5	41.4	31.5	41.4	31.5	41.4	19.2	
1992	5.5	13.6	16.6	15.0	7.9	10.7	22.9	24.6	30.7	24.6	30.7	30.7	30.7	30.7	15.0	
1993	3.1	13.8	18.9	18.5	12.4	16.9	28.6	28.0	30.9	28.0	30.9	30.9	30.9	30.9	17.6	
1994	3.4	13.2	25.2	21.3	15.7	21.0	30.8	35.4	32.9	35.4	32.9	32.9	32.9	32.9	19.5	
1995	4.5	13.6	21.7	22.1	18.8	20.5	29.7	38.3	29.7	38.3	29.7	29.7	29.7	29.7	20.2	

Table B.11. Annual mean bottom temperature (C) at sampled bay trawl sites in Texas bay systems from 1977-95. ND = no data.

Year	East						Corpus Christi			Upper Laguna Madre			Lower Laguna Madre			Coastwide
	Sabine Lake	Galveston	Matacorda	Matacorda	San Antonio	Aransas	Christi	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Laguna Madre	Coastwide	
1977	ND	18.7	ND	17.9	21.1	17.8	ND	ND	ND	ND	ND	ND	ND	ND	18.8	
1978	ND	21.6	ND	23.5	24.2	24.8	ND	ND	ND	ND	ND	ND	ND	ND	22.9	
1979	ND	22.5	ND	21.6	25.5	ND	ND	ND	ND	ND	ND	ND	ND	ND	22.8	
1980	ND	23.8	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	23.8	
1981	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
1982	ND	21.8	ND	24.8	23.3	23.1	25.0	26.1	25.1	26.1	25.1	26.1	25.1	26.1	23.5	
1983	ND	21.5	ND	21.7	21.7	22.3	22.2	21.8	22.7	21.8	22.7	22.7	22.7	22.7	21.8	
1984	ND	22.2	ND	22.8	21.6	23.4	21.8	22.0	22.8	22.0	22.8	22.8	22.8	22.8	22.3	
1985	ND	21.9	ND	22.5	22.5	21.7	21.9	23.0	22.8	23.0	22.8	22.8	22.8	22.8	22.2	
1986	22.1	22.2	ND	23.3	23.1	22.1	21.8	23.3	22.5	23.3	22.5	23.3	22.5	23.3	22.6	
1987	20.0	21.5	24.3	21.9	21.8	21.3	21.1	22.3	22.6	22.3	22.6	22.6	22.6	22.6	21.6	
1988	21.8	21.8	21.1	20.2	22.1	21.3	22.2	22.1	24.5	22.1	24.5	24.5	24.5	24.5	21.6	
1989	20.8	20.4	21.0	20.5	21.1	20.5	21.8	23.8	21.0	23.8	21.0	23.8	21.0	23.8	21.0	
1990	21.2	21.4	22.7	22.6	21.9	22.6	23.4	23.8	24.2	23.8	24.2	24.2	24.2	24.2	22.3	
1991	21.7	21.5	22.0	21.5	22.2	21.7	22.8	23.4	23.2	23.4	23.2	23.4	23.2	23.4	21.9	
1992	20.7	21.6	20.6	22.1	22.6	21.4	21.4	22.9	21.7	22.9	21.7	22.9	21.7	22.9	21.7	
1993	21.0	20.9	22.2	22.2	22.5	21.8	22.1	23.7	23.7	23.7	23.7	23.7	23.7	23.7	21.7	
1994	22.1	22.2	22.6	22.6	22.7	20.9	23.0	23.6	24.8	23.6	24.8	24.8	24.8	24.8	22.5	
1995	22.0	21.8	22.6	23.0	23.3	22.5	23.5	23.6	23.9	23.6	23.9	23.9	23.9	23.9	22.7	

Table B.12. Annual mean bottom turbidity at sampled bay trawl sites in Texas bay systems from 1983-95. ND = no data.

Year	East						Upper		Lower	
	Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Corpus Christi	Laquana Madre	Laquana Madre	Coastwide
Jackson Turbidity Units										
1983	ND	101	ND	25	26	105	77	76	38	67
1984	ND	75	ND	30	30	71	62	70	38	55
1985	ND	41	ND	33	55	42	32	52	59	41
1986	35	37	ND	45	53	41	42	49	67	43
Nephelometric Units										
1987	15	17	19	22	29	7	13	15	12	18
1988	17	14	20	23	17	13	15	14	15	16
1989	16	18	27	19	22	19	15	12	14	18
1990	13	18	20	15	28	17	11	15	13	17
1991	18	16	22	19	22	19	10	10	8	17
1992	19	18	17	24	37	30	12	9	18	21
1993	16	24	17	19	19	22	13	13	47	21
1994	17	19	14	13	17	16	10	9	16	15
1995	17	20	24	18	19	17	14	12	19	18

Table B.13. Annual mean bottom salinity (o/oo) at sampled gulf trawl sites in the Texas Territorial Sea 1985-95. ND = no data.

Year	Port O'Connor			Port Aransas		Port Isabel		Coastwide
	Sabine Lake	Galveston	Port O'Connor	Port Aransas	Port Isabel	Coastwide		
1985	ND	30.6	32.3	30.9	31.7	31.4	31.4	
1986	29.1	29.7	32.4	30.5	32.7	30.9	30.9	
1987	27.4	28.8	33.5	34.4	34.4	31.7	31.7	
1988	27.3	28.3	30.7	32.4	35.0	30.7	30.7	
1989	25.4	29.9	32.9	30.9	33.7	30.6	30.6	
1990	25.3	29.5	30.5	32.4	33.9	30.3	30.3	
1991	23.7	28.5	31.0	31.9	31.2	29.3	29.3	
1992	26.5	29.4	31.5	32.4	30.7	30.1	30.1	
1993	23.1	27.3	28.9	34.5	30.8	28.9	28.9	
1994	21.3	27.2	28.1	31.7	33.9	28.4	28.4	
1995	26.1	28.0	31.5	31.2	33.4	30.0	30.0	

Table B.14. Annual mean bottom temperature (C) at sampled gulf trawl sites in the Texas Territorial Sea 1985-95. ND = no data.

Year	Port O'Connor			Port Aransas		Port Isabel		Coastwide
	Sabine Lake	Galveston	Port O'Connor	Port Aransas	Port Isabel	Coastwide		
1985	ND	23.4	23.6	22.5	25.4	23.7	23.7	
1986	25.6	22.0	22.8	22.3	22.7	23.1	23.1	
1987	21.1	21.7	22.1	22.4	21.9	21.8	21.8	
1988	21.1	21.6	21.2	22.2	21.8	21.6	21.6	
1989	19.8	21.5	21.3	21.7	21.8	21.2	21.2	
1990	21.3	21.9	21.8	22.2	21.8	21.8	21.8	
1991	22.0	22.3	22.1	21.8	21.5	21.9	21.9	
1992	19.9	21.5	20.9	22.5	20.9	21.1	21.1	
1993	21.8	21.7	21.8	21.6	21.3	21.7	21.7	
1994	22.1	22.2	21.9	22.2	22.0	22.1	22.1	
1995	22.2	21.9	21.8	22.3	22.2	22.1	22.1	

Table B.15. Annual mean bottom turbidity at sampled gulf trawl sites in the Texas Territorial Sea 1985-95. ND = no data.

Year	Sabine Lake	Galveston	Port O'Connor	Port Aransas	Port Isabel	Coastwide
<b>Jackson Turbidity Units</b>						
1985	ND	31	37	25	24	30
1986	30	24	29	24	24	26
<b>Nephelometric Units</b>						
1987	10	10	11	4	6	8
1988	6	9	10	4	4	7
1989	7	9	9	7	4	7
1990	9	11	7	8	3	8
1991	11	12	7	8	3	8
1992	13	10	10	10	4	9
1993	12	6	14	7	4	9
1994	10	8	5	5	4	6
1995	13	13	7	5	4	8

Table B.16. Annual mean shoreline salinity (o/oo) at sampled 60.9-m beach seine sites in 5 Texas gulf areas 1987-95.

Year	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
1987	28.0	29.8	30.7	32.9	33.5	30.7
1988	28.6	30.8	31.9	35.8	36.8	32.2
1989	22.6	25.3	31.3	32.9	32.9	28.9
1990	24.2	26.5	31.3	31.5	35.6	29.5
1991	24.1	26.1	28.2	30.9	31.5	27.8
1992	27.0	30.3	30.9	32.1	31.7	30.2
1993	23.4	27.4	28.6	32.3	32.5	28.3
1994	23.3	26.4	28.2	31.4	34.6	28.1
1995	24.8	26.8	27.7	31.1	33.1	28.1

Table B.17. Annual mean shoreline temperature (C) at sampled 60.9-m beach seine sites in 5 Texas gulf areas 1987-95.

Year	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
1987	21.0	21.0	22.2	23.4	22.6	22.0
1988	26.7	26.5	26.9	27.5	26.5	26.8
1989	24.2	26.0	26.3	26.6	26.7	25.9
1990	26.1	26.4	26.3	26.9	27.1	26.5
1991	25.8	26.9	26.6	26.8	27.5	26.6
1992	26.5	26.0	25.1	25.9	26.6	25.9
1993	25.9	26.1	25.9	24.8	26.0	25.7
1994	27.7	26.9	26.9	26.9	27.2	27.1
1995	26.0	26.4	26.7	27.4	27.0	26.6

Table B.18. Annual mean shoreline turbidity (NTU) at sampled 60.9-m beach seine sites in 5 Texas gulf areas 1987-95.

Year	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
1987	51	36	41	16	12	35
1988	43	23	30	9	10	26
1989	131	26	39	13	7	50
1990	48	31	28	14	10	28
1991	73	31	31	12	18	36
1992	71	22	35	18	22	37
1993	68	28	35	19	23	37
1994	61	18	25	13	16	29
1995	49	30	31	12	17	30

Table B.19. Annual mean shoreline salinity (o/oo) at sampled 18.3-m bag seine sites in 5 Texas gulf areas 1987-95.

Year	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
1987	27.7	30.0	30.3	33.1	33.6	30.5
1988	28.6	30.8	31.9	35.8	36.8	32.3
1989	22.5	25.3	31.3	32.9	32.9	28.9
1990	25.2	26.6	31.1	32.2	35.5	29.8
1991	23.9	26.1	28.0	31.2	31.5	27.8
1992	27.2	30.0	30.9	32.0	31.7	30.2
1993	23.3	27.3	28.6	32.2	32.4	28.2
1994	23.2	26.9	28.4	31.3	34.4	24.2
1995	25.1	26.8	27.6	30.9	33.5	28.2

Table B.20. Annual mean shoreline temperature (C) at sampled 18.3-m bag seine sites in 5 Texas gulf areas 1987-95.

Year	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
1987	21.9	21.2	22.3	23.8	22.6	22.4
1988	26.8	26.8	26.9	27.5	26.4	26.9
1989	24.3	26.2	26.4	26.6	26.7	26.0
1990	26.2	26.7	26.3	27.1	27.1	26.6
1991	25.8	27.3	26.7	26.9	27.5	26.7
1992	26.6	26.3	25.2	26.0	26.6	26.0
1993	25.8	26.3	25.8	24.8	26.2	25.8
1994	27.9	27.2	26.9	27.2	27.3	27.3
1995	26.1	26.5	26.8	27.5	27.0	26.8

Table B.21. Annual mean shoreline turbidity (NTU) at sampled 18.3-m bag seine sites in 5 Texas gulf areas 1987-95.

Year	Gulf-17	Gulf-18	Gulf-19	Gulf-20	Gulf-21	Coastwide
1987	56	41	45	16	12	38
1988	38	24	28	9	10	24
1989	134	29	37	13	7	51
1990	44	32	28	14	10	28
1991	73	31	31	12	18	36
1992	69	24	37	14	23	37
1993	92	30	36	21	26	44
1994	59	18	26	15	15	29
1995	51	30	32	11	18	31

Table B.22. Annual mean surface salinity (o/oo) at sampled ICWM trawl sites by bay system during 1992-95. ND = no data.

Year	East				Corpus Christi				Lower				
	Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Christi	Christi	Upper	Upper	Laquna Madre	Laquna Madre	Coastwide
1992	8.9	20.0	15.8	19.1	10.6	15.2	26.3	26.3	30.8	30.1	30.1	30.1	19.5
1993	6.2	18.5	18.4	18.1	15.3	22.1	32.0	32.0	31.9	30.6	30.6	30.6	20.5
1994	5.4	16.4	24.1	20.4	20.1	24.3	31.6	31.6	36.0	33.0	33.0	33.0	22.3
1995	6.9	17.3	21.3	22.4	23.4	23.1	31.0	31.0	36.6	33.1	33.1	33.1	22.4

Table B.23. Annual mean surface temperature (C) at sampled ICWM trawl sites by bay system during 1992-95. ND = no data.

Year	East				Corpus Christi				Lower				
	Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Christi	Christi	Upper	Upper	Laquna Madre	Laquna Madre	Coastwide
1992	20.2	21.5	20.8	20.6	22.1	22.5	21.8	21.8	23.0	23.8	23.8	23.8	21.6
1993	21.9	22.2	23.3	22.3	22.4	21.7	22.1	22.1	22.0	23.2	23.2	23.2	22.2
1994	21.6	22.7	23.1	22.7	22.8	21.7	22.8	22.8	23.9	23.4	23.4	23.4	22.7
1995	21.2	22.7	23.9	22.2	23.8	23.2	23.2	23.2	23.0	23.2	23.2	23.2	22.8

Table B.24. Annual mean surface turbidity (NTU) at sampled ICWM trawl sites by bay system during 1992-95. ND = no data.

Year	East				Corpus Christi				Lower				
	Sabine Lake	Galveston	Matagorda	Matagorda	San Antonio	Aransas	Christi	Christi	Upper	Upper	Laquna Madre	Laquna Madre	Coastwide
1992	29	23	22	27	38	25	9	9	7	22	22	22	28
1993	45	22	20	33	22	16	9	9	6	32	32	32	23
1994	33	17	17	12	16	14	6	6	6	47	47	47	16
1995	41	23	32	21	17	14	10	10	9	36	36	36	20



Appendix C. Summary of SEAMAP samples by year and depth zone for brown shrimp, white shrimp, pink shrimp and blue crab collected off Texas during 1982-95.

Table C.1. Mean catch rates (No./h) and mean size (mm) of select shellfishes caught during SEAMAP\* sampling off Texas during June-July 1982-95. Blanks indicate no measurement taken.

Year	Depth (m)	Samples (No.)	Brown Shrimp		White Shrimp		Pink Shrimp		Blue Crab	
			No./h	Length	No./h	Length	No./h	Length	No./h	Length
1982	0-18	22	1,222	108	15	173	161	136	8	
	19-37	50	1,427	115	0		20	138	1	
	38-55	29	138	145	0		<1	126	0	
	56-73	5	117	179	0		0		0	
74-91	3	79	182	0		0		0		
1983	0-18	28	254	99	20	153	195	127	8	
	19-37	47	1,445	119	1	167	87	121	4	
	38-55	24	304	132	0		1	118	1	
	56-73	8	66	156	0		0		0	
	74-91	2	71	168	0		0		0	
1984	0-18	16	733	116	30	174	4	151	6	
	19-37	40	1,594	116	1	168	3	150	0	
	38-55	16	544	131	0		0		0	
	56-73	12	194	138	0		0		0	
	74-91	5	86	151	0		0		0	
1985	0-18	30	450	98	41	168	15	135	20	
	19-37	40	1,362	112	2	167	10	131	4	
	38-55	14	150	127	0		<1	127	0	
	56-73	5	154	144	0		0		0	
	74-91	1	36	179	0		0		0	
1986	0-18	35	250	98	33	165	18	116	11	
	19-37	43	809	108	0		42	130	10	
	38-55	10	311	124	0		0		0	
	56-73	5	176	136	0		0		0	
	74-91	3	49	147	0		0		0	
1987	0-18	74	189	103	15	159	24	115	3	
	19-37	56	606	107	3	162	19	108	7	
	38-55	17	26	142	0		<1	180	2	
	56-73	8	16	177	0		0		1	
	74-91	7	11	177	0		0		0	
1988	0-18	75	227	106	4	166	22	110	5	
	19-37	50	309	113	0		2	127	2	
	38-55	17	18	126	0		0		0	
	56-73	7	4	180	0		0		0	
	74-91	7	3	198	0		0		0	
1989	0-18	85	556	101	16	167	51	116	6	111
	19-37	54	928	118	4	126	24	116	1	144
	38-55	12	212	129	0		<1	135	0	
	56-73	8	40	140	0		0		0	
	74-91	7	11	159	0		0		0	
1990	0-18	74	279	113	17	171	18	126	5	127
	19-37	48	850	123	1	156	62	122	2	81
	38-55	16	202	136	0		<1	135	1	79
	56-73	10	76	140	0		0		0	
	74-91	8	16	154	0		0		<1	164

Table C.1. (Cont'd.)

Year	Depth (m)	Samples (No.)	Brown Shrimp		White Shrimp		Pink Shrimp		Blue Crab	
			No./h	Length	No./h	Length	No./h	Length	No./h	Length
1991	0-18	92	202	106	31	167	27	125	14	90
	19-37	51	1,153	125	7	173	64	136	4	143
	38-55	20	186	143	0		<1	157	1	135
	56-73	10	76	171	0		0		1	96
74-91	9	41	176	0		0		0	0	
1992	0-18	85	234	100	36	166	15	112	4	114
	19-37	58	217	127	<1	185	6	121	1	148
	38-55	17	22	158	0		0		<1	248
	56-73	10	15	180	0		0		0	0
	74-91	8	10	186	0		0		0	0
1993	0-18	89	121	104	16	171	23	122	10	120
	19-37	55	236	111	2	169	63	121	6	119
	38-55	22	69	139	0	176	19	122	<1	152
	56-73	10	35	152	0		0		<1	161
	74-91	2	34	169	0		0		<1	140
1994	0-18	93	113	109	9	169	58	124	6	125
	19-37	50	850	120	0		27	118	<1	102
	38-55	19	46	151	0		0		0	0
	56-73	11	36	181	0		0		0	0
	74-91	3	12	181	0		0		0	0
1995	0-18	88	343	105	19	165	34	117	6	112
	19-37	54	829	114	8	165	10	109	1	142
	38-55	20	101	125	0		<1	100	<1	143
	56-73	11	106	148	0		0		0	0
	74-91	7	43	177	0		0		0	0

Data presented here were collected by R/V OREGON II (NMFS) in conjunction with TPWD research vessels. The data were made available by the Southeast Area Monitoring and Assessment Program (SEAMAP). Samples collected with 12.2-m trawl, except 6.1-m trawl by TPW vessels since 1987. Data normalized to 12.2-m trawl by NMFS.

Table C.2. Mean catch rates (No./h) and mean size (mm) of select shellfishes caught during SEMAP\* sampling off Texas during November 1986-95. Blanks indicate no measurement taken.

Year	Depth (m)	Samples			Brown shrimp			White shrimp			Pink shrimp			Blue crab		
		(No.)	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length	No./h	Length
1986	0-18	12	71		77		26					0				
	19-37	34	93		15		2				1					
	38-55	26	68		0		0				0					
	56-73	12	41		0		0				0					
	74-91	4	22		0		0				0					
1987	0-18	65	20		89		18				0					
	19-37	40	50		7		2				<1					
	38-55	12	21		0		0				0					
	56-73	2	6		0		0				0					
	74-91	1	0		0		0				0					
1988	0-18	77	21		98		9				0					
	19-37	49	48		15		12				0					
	38-55	16	44		0		1				0					
	56-73	10	15		0		0				0					
	74-91	7	8		0		0				0					
1989	0-18	78	21	100	137	102	16	124	124	16	124	2	45			
	19-37	60	68	140	23	117	10	123	123	10	123	<1	83			
	38-55	20	71	169	<1		1	124	124	1	124	<1	94			
	56-73	7	43	173	0		0			0		<1	74			
	74-91	9	5	185	0		0			0		0				
1990	0-18	64	18	105	56	129	11	137	137	11	137	<1	70			
	19-37	59	69	140	5	159	7	126	126	7	126	<1	87			
	38-55	22	60	168	<1		1	129	129	1	129	1	75			
	56-73	9	34	173	0		0			0		1	74			
	74-91	6	7	190	0		0			0		0				
1991	0-18	88	28	107	31	124	14	108	108	14	108	<1	52			
	19-37	57	120	134	4	166	4	107	107	4	107	<1	133			
	38-55	20	65	161	0		0			0		1	135			
	56-73	12	31	172	0		0			0		0				
	74-91	11	12	181	0		0			0		0				
1992	0-18	89	11	115	135	115	3	131	131	3	131	<1	34			
	19-37	55	80	135	8	157	1	122	122	1	122	<1	141			
	38-55	18	42	164	0		0			0		<1	141			
	56-73	8	49	172	0		0			0		0				
	74-91	4	33	176	0		0			0		0				
1993	0-18	88	11	126	160	119	31	95	95	31	95	3	160			
	19-37	55	91	119	17	134	28	88	88	28	88	1	151			
	38-55	17	60	93	<1	109	<1	76	76	<1	76	<1	108			
	56-73	9	12	106	<1	125	<1	76	76	<1	76	0				
	74-91	5	17	85	<1	119	<1	98	98	<1	98	<1	144			

Table C.2. (Cont'd.)

Year	Depth (m)	Samples (No.)	Brown shrimp		White shrimp		Pink shrimp		Blue crab	
			No./h	Length	No./h	Length	No./h	Length	No./h	Length
1994	0-18	89	13	108	36	124	19	106	1	105
	19-37	54	106	132	5	152	6	110	<1	134
	38-55	17	53	162	0	0	0	0	0	0
	56-73	13	26	177	0	0	0	0	0	0
	74-91	11	8	192	0	0	0	0	0	0
1995	0-18	95	5	118	120	114	10	126	5	54
	19-37	51	131	129	15	162	19	117	<1	75
	38-55	19	49	162	0	0	2	138	0	0
	56-73	9	45	166	0	0	<1	160	0	0
	74-91	9	54	177	0	0	0	0	0	0

\*Data presented here were collected with 12.2-m trawl by R/V OREGON II (NMFS) and with 6.1-m trawl by TPWD research vessels. The data were made available by the Southeast Area Monitoring and Assessment Program (SEAMAP). Data normalized to 12.2-m trawl by NMFS.

PWD RP V3400-461 (11/97)

Dispersal of this publication conforms with Texas State Documents Depository Law, and it is available at Texas State Publications Clearinghouse and/or Texas Depository Libraries.